


KILGORE'S PLANTING GUIDE FOR FLORIDA



THE KILGORE SEED COMPANY

General Offices and Maii Order Department, Plant City, Florida

A DIVISION OF ASSOCIATED SEED GROWERS, INC.





GREAT LAKES 118 LETTUCE

THE KILGORE SEED COMPANY

PLANT CITY, FLORIDA
A DIVISION OF ASSOCIATED SEED GROWERS, INC.



AMOUNT ENCLOSED

Post Office							
					Money Order		-
					Cash		
Ship by	Parcel Post	Express	Truck	Freight	Casii		-
			Ш		Total		_
Exp. or Frt. O	ffice	Date		19			
	IMPORTANT! Be su	are address and ship	oing instructions ar	e correct. Please	write plainly.		
SPECIAL N On items n Sales Tax"	OTICE: Remittances show not marked postpaid be s on non-exempt merchandi	old be made by po sure your remittance se.	stal money orders includes the neces	checks, or co sary amount for	ash. DO NOT SEN postage and also	ND STAMPS. the "State	
	that the seeds, bulbs gnized tolerances; but w s and plants. We give stutes Section 578.26 re ging damages sustained ten days after defect						
ree, wimin	ten days after defect	becomes apparent; (and copy of compi	uint be sent to .	Seller by registered	l Pric	
antity ITEMS WANTED					Dollars	e Ce	
		N.					
	P-2						
	100						
-31-5	ou included postage on i						+-
Have yo		te the next best or mo					

O
ear
-
-
9
-
*
Along
-
2
.3
Q
This
3
-
S
-
1
Lin

Quantity	ITEMS WANTED	_	Price Dollars	Cents
	To Gallery and A			
3 80 - 10 10 10				
			W	
			A to	
			0	
				-
				-
			7	
			11	
Have you included	posteria en itame not meste il and "Et e C. I. T. "			
	postage on items not postpaid and "State Sales Tax" o		١	
achines, garden tools, cans and o	hing you need in vegetable, flower and field crop sec canning equipment, poultry and bee supplies? Extra	eas, tertilizers, sprays, dusts, sp order blanks and envelopes sen	raying and t upon app	dusting lication
LEASE WRITE BELOW	The names of neighbors or friends who might be i			
NAMES				
INAMES	POST OFFICE	P.O. Box, Street, or R.F.D.	STAT	E
		0		

Important Directions to Customers

By carefully observing the following directions, mistakes and misunderstandings will be largely avoided.

How to order—Please be careful to sign your name, box number, street or rural route, post office and state on every order. Be sure to tell us your nearest express or freight office should you desire the seed or merchandise forwarded by express, freight or truck line. Please write plainly.

Cash with order or C. O. D.—Ours is a cash business and our prices are made on that basis. Customers are requested to remit when ordering, and by so doing save the c.o.d. fee charged by express companies, banks, and postal authorities for collecting and returning money.

Perishable items not sent C.O.D.—Perishable items, such as strawberry plants, seed potatoes, onion sets, etc., are not sent c.o.d. under any circumstances. Remittances must be with order.

How to send remittance—Remittances should be made by postal money orders, drafts, checks or cash. Please do not send stamps.

We pay postage to points in Florida — The postpaid prices indicated in this catalog include delivery to any point in Florida.

Prices subject to change—This catalog is printed during the early fall months. The market on many items of garden and field seeds, as well as supplies, has not yet been fully established. Consequently, prices on these items indicate value at time these pages are printed. In practically all cases orders will be filled at prices indicated in this catalog, but all prices are subject to change without notice. Write for special prices on larger quantities.

Export Shipments to Cuba and the West Indies—For the convenience of our customers in the West Indies, we have established the following agencies: E. G. Del Campo, Nueva Gerona, Isle of Pines, Cuba; F. C. Albury, Nassau, Bahamas.

For direct deliveries outside the continental United States, all prices quoted as "Postpaid" in this catalog are on an f.o.b. Plant City, Florida, basis, and the proper amount for postage must be added to the cost of the seeds.

Parcel Post rates to Cuba and the West Indies—The parcel post rate to Cuba, Bahamas and Jamaica is 45 cents, for Haiti 61 cents, and for the Dominican Republic 58 cents for the first pound or fraction thereof. If the net weight of an order is 1 pound or more, 1 pound must always be added to cover weight after being packed for shipment. In each case add 22 cents per pound for each additional pound or fraction thereof to the above indicated pound rates. Packages of merchandise going to Cuba, Haiti or Dominican Republic, weighing 8 ounces or less are subject to the postage rate of 1½ cents for each 2 ounces or fraction of 2 ounces. To the Bahamas and Jamaica, packages weighing 8 ounces or less are subject to the postage rate of 3 cents for each 2 ounces or fraction thereof, with a minimum charge of 15 cents.

WARRANTY AND LIMITATION OF LIABILITY—"We warrant that the seeds, bulbs and plants we sell are, at the time of delivery, as described on the container within recognized tolerances; but we limit our liability on this warranty to the amount of the purchase price of such seeds, bulbs and plants. We give no other or further warranty express or implied." The Kilgore Seed Company.

Florida Statutes Section 578.26 requires as prerequisite to suit for defective seeds that sworn complaint against Seller alleging damages sustained be filed with Florida Commissioner of Agriculture, accompanied by \$10.00 filing fee, within ten days after defect becomes apparent; and copy of complaint be sent to Seller by registered mail.

Air express shipments can now be made from Miami or Tampa, Florida, to any point in the West Indies.

Return of seeds for exchange—To safeguard our customers and ourselves, seeds may not be returned if seed package has been broken. In case of error on our part, unbroken packages may be returned within 10 days after shipment and we will gladly make correction at our expense.

Stock or lot number on all Kilgore's seed packages—Every lot of seed distributed by the Kilgore Seed Company will indicate a stock number or lot number. This stock number is a code reference to our records of the history, performance and production of that particular lot of seed, by means of which we can trace it back to the growers' field where it was produced and to our trial plantings. It is recommended that every commercial grower of vegetables retain the empty packages or tags with the stock numbers. Correspondence pertaining to the performance of Kilgore's seeds should include the specific lot number of the seed under consideration to aid investigation.

Postal, Express, and Sales Tax information—The parcel post weight limit for first and second zones is 40 pounds; 20 pounds in the third to eighth zones. Any weight up to 40 pounds in the first and second zones and 20 pounds in the third zone will usually go just as cheaply by parcel post as by freight or express.

In sections of Florida that are not easily accessible by freight or express, parcel post service will be found not only more satisfactory, but more economical. If on a rural route, delivery will be made to your door by mail.

Most Florida points are in the 1st (within 50 miles from Plant City, Fla.) or 2nd (within 50 to 150 miles from Plant City, Fla.) Postal Zone. Jacksonville north, Live Oak north and west, and points on the east coast south of West Palm Beach are in the 3rd Zone. (Within 150 to 300 miles from Plant City, Fla.) Puerto Rico and Virgin Islands are in the 6th Zone for surface mail; 8th Zone for Air Parcel Post. Surface mail 28c for 1st lb., 12c for each additional lb. Air Parcel Post 80c lb.

	1st an	d 2nd	d ZO	NES				3rd Z	ONE		
1	lb.	.23	11	lbs.	.63	1	lb.	.23	11	lbs.	.75
2	lbs.	.27	12	"	.67	2	lbs.	.29	12	"	.80
3	"	.31	13	"	.71	3	"	.34	13	"	.85
4	"	.35	14	"	.75	4	"	.39	14	"	.90
5	"	.39	15	"	.79	5	"	.44	15	"	.96
6	"	.43	16	"	.83	6	"	.49	16	"	1.01
7	"	.47	17	"	.87	7	"	.54	17	"	1.06
8	"	.51	18	"	.91	8	"	.60	18	"	1.11
9	"	.55	19	"	.95	9	"	.65	19	"	1.16
10	"	.59	20	"	.99	10	"	.70	20	"	1.21

If in doubt about your zone rate, obtain it from your postmaster, and be sure to send enough money to cover necessary postage, if the items ordered are not indicated "postpaid" in this catalog. Do not have your order delayed or reduced by failing to add necessary money for postage. If you send too much money, it will be returned to you. It will be found plainly stated above each item whether or not it is postpaid. Also be sure to include with your remittance the State Sales Tax on non-exempt merchandise. (See page 2).

When you order your seeds and supplies to come by express or freight we can send them collect, so you may pay the transportation charges on delivery. Give your express or freight station, if it is not the same as your post office.

SALES TAX INFORMATION

The Florida Revenue Act of 1949 (Sales and Use Tax) imposes a 3% tax on the retail sales of many commodities. Numerous agricultural supply items are exempt from the tax and for your guidance in determining tax exempt items shown in this catalog, we list below those items that are not subject to the tax.

EXEMPT ITEMS — Bee supplies Except Bodies & Frames)

Cans and Jars
Crate Materials
Feeds—For farm animals
Fencing—For farm use
Grafting Wax and
Rubber Strips
Hotkaps
Label Paste
Paper Pots—Plant Bands
and Flats
Peat Moss
Poultry and Stock
Remedies

Seedbed Cloth
Seed Inoculants
Seeds—All Vegetable,
Field Crop, Pasture
Grass Seeds, Onion
Sets, Seed Potatoes
Seed Treating
Compounds
Starter Solutions and
Root Inducing
Materials
Terra-Lite
Tractors
Weed Killers

EXEMPT ONLY IF USED FOR COMMERCIAL AGRICULTURAL PURPOSES

Dusters
Farm Tools
Fertilizers
Insecticides and
Fungicides
Soluble Plant Foods

Fumigants Garden Tools Raffia and Twine Rubber Bands Sprayers Tarpaulins All flower seed and other merchandise listed in this catalog is subject to 3% State Sales Tax except sales to Dealers, Government Agencies, Schools, Churches and Charitable organizations provided Tax Exemption Certificate has been obtained from the State Comptroller. All orders from institutions or organizations claiming tax exemption must clearly indicate on order sent us, the certificate serial number and date of certificate.

THE RATE OF TAX APPLICABLE TO NON-EXEMPT SALES IS AS FOLLOWS:

- (1) On single sales in amounts from 10c to 35c, both inclusive, 1c shall be added for taxes.
- (2) On sales in amounts from 35c to 65c, both inclusive, 2c shall be added for taxes.
- (3) On Sales in amounts from 65c to \$1.00, both inclusive, 3c shall be added for taxes.
- (4) On sales in amounts of more than \$1.00, 3% shall be charged upon each dollar of price, plus the above bracket charges upon any fractional part of a dollar in excess of even dollars.

SPECIAL NOTICE—In order to avoid delay in filling your order, on all non-exempt items, be sure to include with your remittance the proper amount of sales tax as indicated above. We are not permitted by law to assume the Sales Tax.

FLORIDA HOME GARDENS IN 1960

Florida is one of the most important commercial vegetable producing states of the Union. Florida home vegetable gardens offer almost no competition to commercial vegetables which are grown mainly for shipment to northern markets. Many Florida families who of not get fresh vegetables from their own gardens, seldom get them from the market, at least in sufficient quantity for good health. Florida people, city residents as well as rural folks, can help solve their food problems by producing vegetables in the home garden.

Florida climate and soils are favorable for producing the finest vegetables in the United States. In Florida fresh vegetables can be grown every month in the year, especially during the fall, winter and spring months when they cannot be grown in other sections of the United States. In addition to producing fresh vegetables in your Florida home garden, plans should also be made for preserving vegetables from the garden by canning or freezing for home use which will mean not only better health for the family, but also a material saving in cost. For best results, order your seeds from the Kilgore Seed Company — Florida's Leading Seedsmen. Mail Order Department, Plant City, Florida.





Location and plan of the garden

A level, well drained piece of ground, convenient to the house should be selected for the garden. It should be located where it can be irrigated if necessary and away from hedges, large trees or the shade of buildings. On the farm it should be well-fenced to protect it from rabbits, chickens and other animals.

After selection of a location, and determining the size of the garden, a list of vegetables and flowers should be made. In selecting varieties, primary consideration should be given to the likes of the family, also to the size of the plot available. Different vegetables give variety to the diet.

A map or plan of the garden should be made, showing the various varieties to be grown, and their position in the garden, indicating length of rows and the space between rows. In case of a small, backyard garden, entirely worked by hand, the rows should be close together, i.e., the smaller figure in column 5 of our chart on page 4, so that large quantities may be produced on a limited space.

The planting data given on pages 4, 5, 6 will help greatly in preparing a plan since seed requirements, planting depth, distances and time of maturity are given.

Making a seedbed

Small seeds such as lettuce, cabbage, collard, pepper, eggplant, tomato, and most flowers, should be sown in a seedbed and the young plants transplanted to the field or garden. A shallow box or flat may be used for this purpose, or a small plot in the garden may be used as a seedbed for starting plants for transplanting. The soil should be in perfect condition. Make seedbeds four to six inches high, to prevent overflow in case of heavy rains. In preparing seedbed soil it is desirable to mix in "Terra-Lite" (Vermiculite) in order to increase germination of seed and prevent damping off of seedlings. (page 87). Fertilize with 2 to 3 pounds per 100 sq. ft. of a commercial fertilizer such as Vigoro or Vertagreen (page 79), applied at least a week or ten days before sowing seeds. Sow the seed in rows about six inches apart. Very fine seed like celery, and petunias should not be covered with soil, but may be pressed into the soil with a light roller or with a wide board. Larger seeds like tomato, pepper, and cabbage may be covered, not over ½ to ½ inch deep.

We recommend the use of burlap or white muslin as a cover over the seedbed, stretching it on a three-or four-foot wide tent-shaped or slanting frame built over the seedbed, with the sides about 10 to 12 inches from the ground to provide ample circulation of air under the cover. If plants are being started in late

summer for fall planting, the seedbed cover will serve as a shade to protect seedlings from the hot sun, also to break the force of heavy rains. If plants are started in winter for a spring crop, this seedbed cover will protect seedlings from frost or freezes. When seeds begin to germinate these ground covers should be removed immediately to prevent spindly plants. After the seed is sown, it is desirable to sprinkle every evening or oftener if necessary with cool water by means of hand sprinkling pots.

Success in getting plants to grow satisfactorily depends very much upon how the plants are grown in the seedbed. For a week or ten days before transplanting, the plants in seedbed should be gradually hardened. This means leaving the cover off the plants for a week or so before transplanting, and water should be withheld from the plants during this period. When plants are about six inches high they are ready to transplant into the field or garden, preferably on a cloudy day or toward evening. Water the plants thoroughly several hours before removing from seedbeds, and again immediately after setting. A starter solution is helpful (page 79). To grow good vegetables and flowers, a large supply of moisture and fertilizer is essential until the crop is ready to harvest.

Preparation of the soil for the garden

The soil in the garden should be well prepared. Grass and weeds removed with all roots. A little extra time spent in preparing the soil for planting will save many hours of hoeing later. It is best to spade or plow, and apply commercial fertilizer to the soil ten days to two weeks before planting in order to avoid burning the seed or young plants. A sample of the soil should be tested, either by yourself with an inexpensive soil test kit (page 88) or by your county agent, and if found too acid or sour apply Hardwood Ashes or Hydrated Lime; if too alkaline or sweet apply Aluminum Sulphate or Manganese Sulphate (page 80).

Dairy Organic Compost is excellent for fertilizer and land conditioner, will not burn, is odorless and contains no weed seed. Use at the rate of 20 to 25 lbs. per 100 sq. ft. broadcast. (page 79). If available, leaf mold or black muck or hammock soil are ideal materials to work into soil. If none of these materials are available, a two or three inch layer of horticultural grade peat moss (page 87) may be worked into the garden soil. This material is clean and easy to handle, and is wonderfully effective for breaking up hard soil, making it mellow, or for adding humus to sandy soil, making it more retentive of moisture and fertilizer materi-

als. After the peat moss is worked into the soil, an inch or two inch layer of Pulverized Sheep Manure, or Dairy Organic Compost, along with a liberal application of a good garden fertilizer such as Vigoro or Vertagreen (page 79) should be used, applying one of these fertilizer materials broadcast at the rate of 4 to 6 lbs. per 100 square feet. If this material is raked into the surface soil a week or ten days before planting, and if well watered in case of no rain, in order to speed the incorporation and decomposition of the fertilizer or manure, you will have the foundation for an excellent garden. If the fertilizers are applied just before planting, they are very apt to burn the young seedlings as they emerge from the seed.

After the plants are well started in the garden, fertilizer should be applied lightly every two or three weeks on either side of the row, but not in contact with the leaves or stems to avoid burning.

Note:—All materials indicated above, such as Pulverized Sheep Manure, Dairy Organic Compost, Peat Moss, Terra-Lite, Super Phosphate, Vigoro, Vertagreen, Wood Ashes, Lime, Aluminum Sulphate, Manganese Sulphate, etc., are available at all Kilgore seed stores, and in our mail order department at Plant City, Florida.

For further information on Florida home gardening write the Florida Agricultural Experiment Station, Mailing Department, Gainesville,
Florida, for a free copy of Bulletin 131 entitled "The Florida Home Garden," and also for
Circular 104 entitled "Vegetable Garden Production Guide."



Reference Table for Florida Vegetable Gardens

Many mistakes may be avoided by following directions given in this table. The time requirements indicated for crops to reach edible size vary, of course, according to the season, varieties and soil conditions. Allowance also must be made for the relative estimate of the

yields, due to climatic and soil conditions. However, the averages indicated in the table will be found to be fairly accurate for Florida growing conditions on the basis of many years of observation and experience.

Kind of vegetable	Tender- Hardy	Number Running feet of row for a family of 3 or 4 people (Includes for canning)	Seeds or plants to provide for a family of 3 to 4 people	Depth to cover in inches	Space between rows in feet	Set plant from seedbed or thin out to inches in row	Approximate time required to produce crop, from seed planted in garden or from transplanting plants to garden (days)
BEANS (Bush)	т	100*	1/2 lb.†	11/2-2	2-21/2	2.4	40
BEANS (Pole)		50	1/4 lb.	11/2-2		3-4 9 or 3 ft. hills	48 to 56
BEANS (Bush Lima)		100*	1/2 to 1 lb.†	11/2-2	3-4 3		63 to 70
BEET		100*	1 oz.†	1/2-2	11/2-2	4-5	65 to 75
BROCCOLI		40	1 pkt. or 25 plts.	1/4-1/2		3-4	56 to 68
BRUSSELS SPROUTS	Н	25	15 plants	1/4-1/2	3	18-24	55 to 60 from pits.
CABBAGE		60	•		21/2-3	18-24	90 from plants
CHINESE CABBAGE		40*	50 plants	1/4-1/2	21/2-3	12-16	62 to 98 from plts.
CANTALOUPE		50	1 pkt. or 50 plts.† 1 pkt.†	1/4-1/2	21/2-3	8-12	70 from seed
CARROT	Н	100*	2 pkts.†	1/2-3/4	4-5	24 or 4 ft. hills	75 to 95
CAULIFLOWER		50			11/2-2	2-3	72 to 95
CELERY		50	30 plants	1/4-1/2	3-4	18-24	55 to 62 from plts.
COLLARD		50*	150 plants	1/8	3	4-5	105 to 130 from plts.
CORN (Sweet)	т Т	200*	1 pkt. or 35 plts.†	1/4-1/2	21/2-3	14-18	50 to 52 from plts.
CORN (Roasting Ear)		200*	1/4 lb.†	1-2	21/2-3	1215	65 to 80
COWPEAS (Table)		150	1/4 lb.†	1-2	3-31/2	15—18	68 to 87
CUCUMBER		50	1/2 to 3/4 lb.	1-11/2	3-4	56	68 to 85
EGGPLANT		50	1 pkt.	3/4	56	12 to 24	40 to 50
ENDIVE (ESCAROLLE)		40	17 plants	1/2	4-41/2	18 to 36	80 to 85 from plts.
GARLIC (bulbs or sets)		30	pkt. or 50 plts.	1/4-1/2	11/2-2	9	90 to 95 from seed
KALE		25	1 lb.	1 1	11/2	4	90
KOHL RABI		25*	1 pkt.	1/4-1/2	21/2	20-24	70
LEEK		50	1 pkt.†	1/4-1/2	11/2-2	68	50
LETTUCE (Head)		75*	1 pkt. or 120 plts.	1/2	11/2	5–6	140 from seed
LETTUCE (Leaf)		50*	1 pkt. or 90 plts.†	1/4-1/2	11/2-2	10-15	66 to 84 from seed
MUSTARD		40*	1 pkt. or 60 plts.†	1/4-1/2	11/2-2	10-12	46 from seed
OKRA			1 pkt.†	1/4-1/2	2	8-10	40 to 45
ONION SEED		75	l oz.	1-11/2	3-4	10-15	50 to 60
ONION SETS		80	1 pkt. or 400 plts.	1/2	11/2	2-4	110 to 150 from seed
PARSLEY		40 30	1 pt.	1	11/4-11/2	1–2	30 to 60
PARSNIP		50	1 pkt.	1/4-1/2	11/2-2	1012	90
PEAS (ENGLISH)		100*	1 pkt.	1/2	21/2-3	4	120
PEPPER	T		½ lb.†	11/2-2	3	12	50 to 53
POTATOES (IRISH)		40 200	24 plants	1/2	21/2-31/2	10-16	60 to 80 from plts.
PUMPKIN	_ †		12 lbs.	1-4	3-31/2	9–12	75 to 90
RADISH		50	1/2 oz.	1	8—10	5 ft. hills	100 to 118
RHUBARB	H	25*	l pkt.†	1/2	1-11/2	1-2	21 to 25
ROSELLE	T	25	1 pkt. or 15 plts.	1/2-3/4	$2\frac{1}{2}$	18-24	90 from plants
RUTABAGA	" .;	25	1 pkt.	1/2-3/4	5–6	24-36	175
SPINACH-SAVOY		50	1 pkt.	1/4-1/2	2	5–6	90
SPINACH-NEW ZEALAND	Н	50*	1 pkt.†	1/2-3/4	11/4-11/2	6	40
SQUASH (Bush)	1	25	1 pkt.	3/4—1	3	20	60
SQUASH (Running)	T	50	1 pkt.	1	3-4	2 ft. hills	42 to 52
SWISS CHARD		50	1 pkt.	1	6-8	3 or 4 ft. hills	60 to 115
OMATO-GROUND		25	1 pkt.	1/2-1	2	12-15	58 to 60
		150	50 plants	1/2	31/25	24-36	70 to 90 from plts.
OMATO-STAKED		75	50 plants	1/2	4	16-20	70 to 90 from plts.
URNIP	1	50	1 pkt.†	1/4-1/2	11/4-11/2	46	35 to 50
VATERMELON	T	75	1 pkt.	1/2-1	8-10	6 to 8 ft. hills	85 to 100

^{*} Make two or more plantings at different times during the season. Number of feet or row indicated is for each planting.

THE KILGORE SEED COMPANY—A Florida Institution

GENERAL OFFICES AND MAIL ORDER DEPARTMENT, PLANT CITY, FLORIDA

A Division of Associated Seed Growers, Inc.

Fourteen Kilgore stores serving Florida, located at:

Belle Glade Fort Myers Gainesville Homestead Miami— 600 N.W. Fifth St. Ocala Pahokee

Palmetto
Plant City
Pompano Beach
Sanford

Vero Beach Wauchula West Palm Beach 1310 Belvedere Road

[†] Number of feet and amount of seed or number of plants indicated are for each planting.

Reference Table for Florida Flower Gardens

Kind of Flower	Average Height (Inches)	or	When to Plant	Approxi- mate Time Sowing to Bloom (Months)	Space Between Rows (Feet)	Set Plants From Seedbed or Thin Out to Inches in Row (Inches)	to Sow	Use
AGERATUM Regular Dwarf	18 6	т	SeptMarch	2½	1	arf Blue Cap 6 le Perfection 12	Flats or Seedbed	Bouquets, Bedding, Borders, Rock Gardens
ALYSSUM	6 to 8	Н.	SeptJan.	11/2	1	6	Flats or	Borders, Rock Gardens
ARCTOTIS DAISY	18 to 22	Н	SeptJan.	3 to 4	3	18	Seedbed Garden	Bouquets, Bedding, Rock Gardens
ASTER	15 to 30	H	SeptApril	3 to 4	2	18 to 24	Flats or Seedbed	Bouquete Paddia - Pad C
BABY'S BREATH	18 to 24	Н	SeptJan.	2 to 21/2	3	18	Garden	
BACHELOR'S BUTTON	24 to 36	Н	SeptJan.	3	2	18	Garden	
BALSAM	18 to 24	Т	SeptMarch	11/2	2	18	Flats or Seedbed	Edwines or Boule
BEAN—SCARLET RUNNER	8 to 10 ft.	Т	FebMay	1½	Vine	6	Garden	Ornamental Climber for Shade and Attractive Blossoms
BELLS OF IRELAND	20 to 24	Н	OctMay	2	2	18	Garden	Bouquets, Borders
BLANKET FLOWER	20 to 24	Н	Any Month	2 to 3	2	18	Garden	Bouquets, Bedding
BLUE LACE FLOWER	24 to 36	Н	SeptJan.	2 to 3	3	36	Flats or Seedbed	Bouquets, Bedding, Borders,
CALENDULA	12 to 18	Н	SeptJan.	3 to 3½	2	18	Flats or Seedbed	Bouquets, Bedding
CALIFORNIA POPPY	12 to 14	Н	SeptJan.	3	2	6	Garden	Bedding, Borders, Rock Gardens
CALLIOPSIS	36 to 60	H	SeptFeb.	4	4 to 5	36	Garden	Bouquets, Bedding, Backgrounds
CARNATION	18 to 24	Н	SeptFeb.	2 to 3	2	12	Garden Flats or	Bouquets, Borders, Rock Gardens
CELOSIA	12 to 18	H T	SeptDec.	3 to 4	2	12 to 18	Seedbed	
COSMOS	48		JanJune	3	1½ to 2	12	Garden	Bouquets—Dried, Bedding
			March-May	2½ to 3	3 to 4	36	Garden	Bouquets, Backgrounds
CYNOGLOSSUM	18 to 24	Н	SeptMarch	2 to 3	2	18	Garden	Cuttings to Mix with Other Flowers in Bouquets. Bedding, Rock Gardens
DAHLIA 	12 to 14	T	March-April	3	2	12	Flats, Seedbed or in Garden	Bouquets, Bedding
DELPHINIUM	36	н	OctNov.	5	3	12 to 18	Flats, Seedbed or in Garden	Bouquets, Borders, Backgrounds
DIANTHUS	12 to 18	Н	SeptMarch	2½ to 3	2	6	Garden	Bouquets, Bedding, Borders, Rock Gardens
GOMPHRENA	12 to 18	Т	FebJuly	3	2	12	Flats or Seedbed	Borders
GOURD-ORNAMENTAL	8 to 12 ft.	T	FebMay	4 to Fruit	Vine	36	Garden	Shading and Ornamental Fruits
HELICHRYSUM	24 to 36	T	SeptMarch	2 to 21/2	3	12 to 18	Garden	Bouquets—Dried, Backgrounds
HOLLYHOCK	5 to 7 ft.	Н	AugJan.	5 to 6	3	24	Garden	Backgrounds
ARKSPUR	18 to 30	Н	NovJan.	2½ to 3	3	12 to 18	Garden	Bouquets, Bedding, Backgrounds
INARIA	16 to 18	н	SeptDec.	3	2 10" for bedding	8	Flats or Seedbed Garden	Bouquets, Bedding
UPINE	30 to 36	н	SeptDec.	2 to 21/2	3	12	Garden	Bouquets, Bedding, Backgrounds, Borders—Dwarf
MARIGOLD Standard Dwarf Sunset Giant	36 12 to 15 42 to 60	т	FebMay and again in Sept.	1½ to 2	2 Giganted 4	rf Varieties 15 to 18 a Sunset Giant 36 ner Varieties 15 to 18	Garden	Bouquets, Bedding—Dwarf types Borders—Dwarf
MORNING GLORY	8 to 10 ft.	Т	March-June	2	Vine	12 to 18	Garden	Shade and Showy, Large, Attractive Blossoms
NASTURTIUM	12 to 18	Т	SeptMarch	2 to 3	2	iem Mixture 12 ner Varieties 12	Garden	Bouquets, Bedding, Rock Gardens
PANSY	6 to 8	Н	SeptDec.	3	11/2	12	Flats or Seedbed	Bouquets, Bedding, Borders, Rock Gardens

Reference Table for Florida Flower Gardens (Continued)

Kind of Flower	Average Height (Inches)	Tender or Hardy	When to Plant	Approxi- mate Time Sowing to Bloom (Months)	Space Between Rows (Feet)	Set Plants From Seedbed or Thin Out to Inches in Row (Inches)	Where to Sow Seed	Use
PETUNIA Dwarf Standards	6 to 12 15 to 18	Н	SeptJune	2½ to 3	Hea 2	osy Morn Ivenly Blue 12 her Varieties 18	Flats or Seedbed	Bedding and Roadside Beautification. Borders, Rock Gardens
PHLOX Dwarf	12 to 16 6 to 8	н	SeptFeb.	3	2	6	Garden	Bedding för Gardens. Vacant Lots and Roadsides. Borders—Dwarf, Rock Gardens
PORTULACA	6	T	FebAug.	11/2	1	12	Garden	Borders, Rock Gardens
QUEEN ANNE'S LACE	36 to 60	Н	SeptJan.	2 to 3	4	24	Garden	Bouquets to Mix with Other Cut Flowers. Backgrounds
SALVIA	18 to 24	Н	SeptJune	3 to 4	3	18	Flats or Seedbed	Bouquets, Bedding, Borders
SCABIOSA	15 to 30	Н	SeptDec.	5	3	12	Garden	Bouquets, Bedding
SHASTA DAISY	18	Н	SeptJan.	4	3	12 to 15	Garden	Bouquets, Bedding
SNAPDRAGON	18 to 24	Н	SeptDec.	3 to 4	Giant 3 18 Half Tall 3 12		Flats or Seedbed	Bouquets, Bedding
STATICE	24 to 36	Н	SeptJan.	3	3	18	Garden	Bouquets, Bedding
SUNFLOWER	36 to 48	Т	FebAug.	1½ to 2	3	24	Garden	Bouquets, Backgrounds
SWEET PEA	48 to 60	Н	SeptDec.	2½ to 3	3	1 to 2	Garden	Bouquets
TITHONIA	48 to 60	T	FebJune	3	4	36	Garden	Bouquets, Backgrounds
VERBENA	6 to 8	Н	Any Month	3	2	12	Garden	Borders, Rock Gardens
VINCA	18	Н	Any Month	3	1	12	Garden	Bedding, Borders
ZINNIA Dwarf Standards	6 24 to 36	Т	FebAug.	11/2	2	Dwarf 12 her Varieties 18 to 24	Flats or Seedbed in Garden	Giants for Bouquets and Backgrounds. Dwarf Bedding, Borders, and Rock Gardens

What and When to Sow Grasses, Clovers, and Field Crop Seeds in Florida

JANUARY and FE	BRUARY		JULY		
Alfalfa	Bahia Grass	Rape	7327		
Clover, Crimson	Oats	Rye	Beans, Velvet	Cowpeas	Hegari
MARCH			Beggarweed	Grasses	Millet
Beans, Velvet	Italian Rye,	Peanuts	Chufas	Bermuda, Sudan	
Benne	Sudan, Bahia	Rape	Corn	and Bahia	
Corn. Field	Indigo	Rice			
Crotalaria	Hegari	Egyptian Wheat	AUGUST		
Grasses	Millet	Sorghum	Cowpeas	Bermuda, Sudan	Millet
Bermuda, Carpet,	Peas	Sunflower	Grasses	and Bahia	Sunflower
APRIL			0.0000		J
Beans, Velvet	Grasses	Peanuts	SEPTEMBER		
Beggarweed	Bermuda, Carpet,	Rice	SEPTEMBER		
Benne	Italian Rye, Sudan, Bahia	Egyptian Wheat	Clover, Sweet	Grasses	Lupines
Chufas	Indigo	Sorghum	Kenland Red	Bermuda, Bahia	Rape
Clover, Alyce	Hegari	Sunflower			Rye
Corn Field	Millet		ОСТОВИЯ		
Crotalaria	Peas		OCTOBER		
MAY			Clover, Sweet	Grasses	Lupines
Beans, Velvet	Grasses	Peas	Kenland Red	Bermuda, Italian	Oats
Beggarweed	Bermuda, Carpet,	Peanuts		Rye, Carpet, Bahia	Rape
Benne	Italian Rye, Sudan, Bahia	Sesbania			Rye
Chufas		Egyptian Wheat	NOVEMBER		•
Clover, Alyce	Indigo	Sorghum			
Corn, Field	Hegari	Sunflower	Clover	Grasses	Oats
Crotalaria	Millet		Crimson, Sweet,	Bermuda, Italian Rye, Carpet, Bahia	Rape
JUNE			White, Ladino, Kenland Red	Lupines	Rye
Beans, Velvet	Crotalaria	Peanuts			
Beggarweed	Grasses	Rice	DECEMBER		
Benne	Bermuda, Carpet, Sudan, Bahia	Sesbania			
Chufas	Indigo	Egyptian Wheat	Clover	Grasses	Oats
Clover, Alyce	Hagari	Sorghum	Crimson, Sweet,	Italian Rye,	Rape
Cowpeas	Millet	Sunflower	White, Ladino, Kenland Red	Carpet, Bahia Lupines	Rye
			,	-apines	

BEANS

Our bean seed is high germinating; grown in disease free areas, carefully milled, graded, fitted for the critical grower and is packed in 50 pound bags.

See page 36 for details of the Kilgore method of handling seeds.

Culture of Green and Wax Podded Bush Beans. Beans are grown on all types of soil in Florida, and are not very particular with reference to soil requirements. Beans are a very quick maturing crop.

Spring planting in the southern and central part of the State begins January 1 and continues until April. Fall planting starts in August. For a fall crop, plant from the middle of August until the middle of October; for a main crop on the Florida east coast plant from October to February. Sow seed in drills, 2 to 2½ feet apart, dropping a bean every 2 to 2½ inches, and cover 1½ to 2 inches deep. On muck soils apply 300 to 600 pounds of a 3-8-8 fertilizer mixture per acre. On sandy loam apply 1200 to 1600 pounds of a 4-7-5 fertilizer per acre, making one application. This application should be applied carefully, as there is always a risk of the fertilizer burning and killing the vitality of the seed when both are put in at the same time.

For the control of insects and diseases of beans see page 65. In order to help control soil diseases and insure a better stand, The Kilgore Seed Co. treats all their bean seed with Delsan.

The number of days in parentheses after each variety represents average time to maturity.

GREEN PODDED BUSH BEANS

Plant 50 lbs. per acre

In the garden make four plantings at different times.

Plant 1/2 lb. to 100 feet of row at each planting.

Write for special prices on larger quantities.



Contender

CONTENDER (45 days.) A consistently early, high yielding, attractive appearing, stringless bean, adapted for shipment to fresh markets. Plant is very vigorous and produces pods similar to Black Valentine, although the Contender pods will average three-fourths of an inch longer, are slightly heavier and thicker, are round to oval, and have a tendency to cure slightly.

½ 1b. 20c; 1 lb. 65c; 2 lbs. \$1.25, Postpaid
 Not prepaid, 10 lbs. \$3.10; 50 lbs. \$14.00

RESISTANT ASGROW VALENTINE. (48 days.) This new variety retains all the desirable characteristics that made its noted predecessor popular plus resistance to common bean mosaic. The plants are medium large, vigorous, and more uniformly productive. Pods are stringless, and thick oval in shape, holding up well under long distance shipping.

1/4 lb. 20c; 1 lb. 65c; 2 lbs. \$1.25, Postpaid
 Not prepaid, 10 lbs. \$3.10; 50 lbs. \$14.00



Tendergreen FS.

TENDERGREEN FS. (50 days.) This variety is popular due to its excellent shipping quality. Plants are large, erect, and thick stemmed. The pods average about five inches in length, are attractive in appearance, being nearly straight, round, fleshy, dark green, stringless, and of excellent quality.

1/4 lb. 20c; 1 lb. 65c; 2 lbs. \$1.25, Postpaid
 Not prepaid, 10 lbs. \$3.30; 50 lbs. \$15.00

SEMINOLE. (53 days.) A round podded bush bean released by the Everglades Experiment Station of Florida A.E.S. The plant is somewhat sprawling, with numerous leaves that protect the pods from wind-scars. Seminole has yielded well in widely scattered trials. The pods are similar to Tendergreen in length, slightly lighter in color. Pods are stringless, fiberless and of good flavor, texture, quality and appearance.

½ lb. 20c; 1 lb. 65c; 2 lbs. \$1.25, Postpaid
 Not prepaid, 10 lbs. \$3.30; 50 lbs. \$15.00

SLENDERGREEN. (54 days.) This variety is similar to Tendergreen, except the pods are longer, slimmer, and straighter. The bush holds the pods well off the ground, and the pod set is quite concentrated. It's very attractive appearance makes it outstanding on northern markets.

1/4 lb. 20c; 1 lb. 65c; 2 lbs. \$1.25, Postpaid
 Not prepaid, 10 lbs. \$3.40; 50 lbs. \$15.50

TENDERLONG 15. (50 days.) Developed for disease resistance; longer, straighter pods than Tendergreen. The plant is an erect, compact bush holding the pods well off the ground. Highly productive and resistant to common bean mosaic, several other virus diseases one form of powdery mildew, and with considerable tolerance to Halo blight. The round straight pods, averaging six inches, are stringless and fiberless and of excellent flavor and medium dark green color.

1/4 lb. 20c; 1 lb. 65c; 2 lbs. \$1.25, Postpaid Not prepaid, 10 lbs. \$3.20; 50 lbs. \$14.50

GREEN PODDED BUSH BEANS (Cont'd)

RESISTANT TENDERGREEN. (50 days.) Similar to regular Tendergreen in bush and pod type, but resistant to common bean mosaic, produces longer pods and higher yields than regular Tendergreen when the disease is present.

1/4 lb. 20c; 1 lb. 65c; 2 lbs. \$1.25, Postpaid Not prepaid, 10 lbs. \$3.20; 50 lbs. \$14.50


PLENTIFUL. (50 days). Plants are spreading in growth and should be given a little wider spacing in the row than most bush beans. The pods are long, flat, straight, and of medium green color. Heavy vielder.

1/4 lb. 20c; 1 lb. 65c; 2 lbs. \$1.25, Postpaid Not prepaid, 10 lbs. \$3.20; 50 lbs. \$14.50

WADE. (48 days). The vigorous plants are tall and hold the pods well off the ground. The heavy foliage helps to protect the pods from wind-scars. It is a fleshy, round podded bean of the Tendergreen type, but powdery mildew and mosaic resistant. It has a consistently higher yield and produces long, straight pods. The pods are of a dark green color and hold their green color exceptionally well in transit and on the market. Stringless and little fiber. Good for market as well as for processing. Because of its long bearing season, high quality and disease resistance, it is highly recommended for home gardens and is a good shipper.

1/4 lb. 20c; 1 lb. 65c; 2 lbs. \$1.25, Postpaid Not prepaid, 10 lbs. \$3.60; 50 lbs. \$16.50

WAX PODDED BUSH BEANS



Resistant Cherokee

RESISTANT CHEROKEE. (50 days.) This is a new mosaic resistant strain. Plant large, vigorous, erect and prolific. Early and an enormous yielder of beautiful, attractive appearing, golden-yellow, long, thick oval pods, nearly straight, entirely stringless and of excellent quality. Considered the best wax bean for Florida and the South.

1/4 lb. 20c; 1 lb. 65c; 2 lbs. \$1.25, Postpaid Not prepaid, 10 lbs. \$3.25; 50 lbs. \$14.75


GREEN PODDED POLE BEANS

Plant 30 lbs. per acre

In Garden 1/4 pound to 50 feet.

Culture of Green Podded Pole Beens. Pole beans are later in maturing than bush beans, bearing over a longer season and extremely productive, higher yielding than bush beans, and also produce larger pods. They are somewhat more tender than bush beans if harvested young.

Pote beans should be supported by wires or by good strong poles to keep them off the ground. Poles should be 8 to 9 feet above ground, and firmly in the ground, in rows 3 to 4 feet apart, and 2 to 3 feet apart in the row. In order to keep pole beans bearing over a long season, it is important to keep them closely picked. Pole beans are more susceptible to mildew, rust and other foliage diseases than are bush beans, and therefore should be sprayed or dusted quite often. (See page 65.)



Kentucky 191 (U.S. No. 4)

KENTUCKY 191 (U.S. No. 4). (63 days.) A high yielding type of white-seeded Kentucky Wonder, resistant to certain forms of rust. Plant is vigorous, with dark green leaves. The pods are dark green, oval, straight, long, averaging from 8 to 9 inches, and attractive in appearance. Stringless in the marketable stage. Productive and very desirable for market or home use. Grown most extensively on the west coast of Florida for shipment to southern and northern markets.

1/4 lb. 20c; 1 lb. 65c; 2 lbs. \$1.25, Postpaid Not prepaid, 10 lbs. \$3.60; 50 lbs. \$16.50

McCASLAN. (65 days.) A vigorous grower and heavy producer, unlike other varieties, producing from the ground up over a long season. The pods are medium green, eight inches long, large, flat, slightly curved, thick and meaty, stringless, brittle, fine grained, and of excellent quality if harvested young. Because of the high yield of large, attractive pods, this variety has become the leading pole bean on the lower east coast of Florida for shipment to northern markets.

1/4 lb. 20c; 1 lb. 65c; 2 lbs. \$1.25, Postpaid Not prepaid, 10 lbs. \$3.60; 50 lbs. \$16.50

FLORIGREEN. A pole bean developed by Dr. A. P. Lorz, horticulturist at the Florida Experiment Station, Gainesville, Fla., and Dr. J. M. Walters, Pathologist of the Gulf Coast Station, Bradenton, Fla.

Florigreen is highly resistant to rust. Also resistant to common and southern bean mosaics. The pods resemble U.S. No. 4 or 191, but a little darker. They are long, straight and retain a fresh dark green color, which is highly desirable for fresh market, stringless at the proper stage for harvesting.

1/4 lb. 20c; 1 lb. 65c; 2 lbs. \$1.25, Postpaid Not prepaid, 10 lbs. \$3.80; 50 lbs. \$17.50



LIMA BEANS

Fordhook—Plant 50 lbs. per acre. Henderson's—Plant 30 lbs. per acre.

In the garden make two plantings at different times. With Fordhook plant 1 pound to 100 feet or with Henderson's plant $\frac{1}{2}$ pound to 100 feet.

Culture of Lima Beans. In general, cultural requirements of Lima Beans are similar to those given for green and wax podded bush beans, but they need more room. Lima beans are not as hardy as string beans, and the ground should be warm when planting the seed. Fertilizer should be applied at least ten days before planting. In order to control soil diseases and get a better stand, it is always desirable to treat lima bean seed before planting (see pages 64 and 74). Plant in rows three feet apart, dropping a bean every 3 to 4 inches in the row, covering an inch deep.

Write for special prices on larger quantities.

FORDHOOK 242. (72 days.) It is an excellent bearer, producing a heavy, early yield, and is especially suitable to Florida because of its ability to make a heavy set and yield under high temperature conditions. The plants are as tall as regular Fordhook, grow more erect and have a more open appearance. Pods average 3 to 4 inches long by 1 and 1½ inches wide and about ½ inch thick, and contain 3 to 4 thick beans. Green shell beans are slightly smaller than regular Fordhook, have light green seed coats, and are of delicious, finely flavored edible quality.

½ 1b. 20c; 1 lb. 65c; 2 lbs. \$1.25, Postpaid
 Not prepaid, 10 lbs. \$3.35; 50 lbs. \$15.25

FORDHOOK-CONCENTRATED. (75 days.) Developed for concentration of pods on the plant, making it easy to harvest, and for concentration in time of reaching maturity. The plants are stocky, without tendrils and very prolific. Pods are dark green, very large, with thick fibrous walls giving them body for long distance shipments. Pods contain 3 to 4 large, very thick, plump beans of excellent quality.

½ 1b. 20c; 1 lb. 65c; 2 lbs. \$1.25, PostpaidNot prepaid, 10 lbs. \$3.30; 50 lbs. \$15.00

HENDERSON'S BUSH LIMA (Baby Lima). (65 days.) An early variety, hardy and a sure cropper. The plants are small and very productive. The pods are small in size, very short and narrow, thin, flat, straight to slightly curved. The seeds are smaller and not as plump as Fordhook. This variety matures about ten days earlier than Fordhook.

½ 1b. 20c; 1 lb. 65c; 2 lbs. \$1.25, Postpaid
 Not prepaid, 10 lbs. \$2.90; 50 lbs. \$13.00

BUTTER OR GREEN SHELL BEANS

Used principally as Green Shell Beans, like Lima Beans Jackson Wonder Bush—Plant 30 lbs. per acre. Horticultural—Plant 50 lbs. per acre. In the garden plant 3/4 pound to 100 feet,

Write for special prices on larger quantities.



Jackson Wonder

JACKSON WONDER (Old Florida Bush Speckled Butter Bean). (65 days.) Culture similar to Lima Beans. This variety is similar to Old Florida Pole, but produces a bush or dwarf plant. Plant is small to medium in size, spreading, with semi-runners resembling somewhat Henderson's Bush Lima. Pods are flat, containing 3 or 4 beans. The seeds are small flat and of light buff color. Early and very prolific, droughresistant, a sure cropper and of good quality.

½ lb. 20c; 1 lb. 65c; 2 lbs. \$1.25, Postpaid
 Not prepaid, 10 lbs. \$2.80; 50 lbs. \$12.50

FRENCH'S HORTICULTURAL. (68 days.) This is the most attractive bean of the speckled Cranberry type making long pods (6 to 8 inches), straight very heavily splashed with bright red or deep carmine, which gives them a very striking and attractive appearance. Strong, erect, sturdy bushy plants, which produce short runners. Very prolific. The shelled beans are pinkish buff color, mottled and streaked. Can be used as a dry shell or green shell bean. Of fine flavor.

½ lb. 25c; 1 lb. 80c; 2 lbs. \$1.50, Postpaid
 Not prepaid, 10 lbs. 5.70; 50 lbs. \$27.00

TAYLOR'S HORTICULTURAL. (64 days.) Used primarily for shelled beans, though immature pods are edible. An improved strain of Dwarf Horticultural type, with plants somewhat larger, more spreading than that variety and more productive. Pods (5 to 6 inches), green turning to cream-yellow, splashed with red.

½ 1b. 20c; 1 lb. 65c; 2 lbs. \$1.25, Postpaid
 Not prepaid, 10 lbs. \$3.90; 50 lbs. \$18.00



BEET

Plant 6 pounds of seed per acre. In the garden make three plantings at different times. Plant 1 oz. to 100 feet of row at each planting.

Culture. Successful culture demands good soil, to which has been applied 600 to 1,200 pounds per acre of good standard vegetable fertilizer. Plant seeds in rows 1½ to 2 feet apart and when 3 inches high thin to 3 or 4 inches apart in the row. In Florida, sow from latter part of September to March.

Prices quoted are postpaid.
Write for special prices on larger quantities.

ASGROW WONDER (Early Blood Red Market). (56 days.) One of the earliest and most uniform beets ever introduced. Roots very attractive in appearance, slightly flattened globe, smooth, with small tap root, of dark red color outside. Flesh of deep dark red, tender, and excellent quality.

Pkt. 10c; oz. 15c; ½ lb. 45c; 1 lb. \$1.40 5 to 25 lbs. \$1.30 per lb.

CROSBY'S EGYPTIAN (Improved). (60 days.) This standard variety produces beets of uniform flattened globe shape, very much flattened at the top, of deep red color outside, with a deep purple color with little zoning inside. Sweet and tender.

Pkt. 10c; oz. 15c; 1/4 lb. 45c; 1 lb. \$1.40 5 to 25 lbs. \$1.30 per lb.

DETROIT DARK RED. (68 days.) A globe-shaped beet of dark red color, outside as well as inside, without white zoning; sweet and of excellent quality.

Pkt. 10c; oz. 15c; ½ lb. 45c; 1 lb. \$1.50 5 to 25 lbs. \$1.40 per lb.

STOCK BEET

Culture. Plant 4 to 6 pounds per acre in rows 2½ to 3 feet apart, and thin to 6 inches apart in the row.

MAMMOTH LONG RED. (80 days.) An enormous yielder and one of the best for dairymen.

Pkt. 10c; oz. 15c; 1/4 lb. 35c; 1 lb. \$1.00 5 to 25 lbs. 90c lb.

SWISS CHARD

Plant 4 pounds per acre. In garden plant one packet to 25 feet of row.

Culture. Plant like beets in the fall, winter, or spring, in rows 2 feet apart, and thin to 12 or 15 inches in the row.

FORDHOOK GIANT. (68 days.) The thick, fleshy, dark green leaves are crinkled or savoyed, and the pearly white stems are very broad, thick, and fleshy. A vigorous, hardy grower.

Pkt. 10c; oz. 15c; ½ lb. 45c; 1 lb. \$1.50 5 to 25 lbs. \$1.40 per lb.

BRUSSELS SPROUTS

Sow 1/4 pound in seedbed to plant an acre. In the garden sow 1 packet in seedbed. Sow seed and cultivate same as cabbage.

LONG ISLAND IMPROVED. (90 days from setting out plants.) Plants grow to a height of two feet. Stems are thickly set with small firm, cabbage-like heads, 1½ inches in diameter, maturing small heads successively.

Pkt. 10c; ½ oz. 35c; oz. 60c; ¼ lb. \$1.90; 1 lb. \$7.00 5 to 25 lbs. \$6.90 per lb.

BROCCOLI

Sow 1/4 to 1/2 pound in seedbed to plant acre.

In the garden 1 packet to 40 feet of row, or 1 packet in seedbed for 25 plants to set 40 feet of row.

Culture. In Florida, seed may be sown in seedbeds from September to January. Plants should be set in rows 3 feet apart, and 1½ to 2 feet apart in the row. When the main head is cut, thick stems form at the base of leaves on the stalk below. At the end of each branch a medium-sized green head developes.

In general, varieties of broccoli fall into three groups which we term Early, Medium, and Late. Those varieties listed as Early must be planted early in the season to give satisfactory quality. Medium varieties are adapted to the widest range of planting dates, but even in this group planting too early in the summer may result in premature flowering. The late types are adapted to areas where broccoli can be grown throughout the winter.

GREEN SPROUTING (Medium). (90 days from plants.) A leading variety for shipping and freezing generally grown for late fall and winter harvest. Plants vigorous, tall, branching and very productive. Heads large, compact, followed by continuous crops of large size shoots suitable for bunching and for freezing.

Pkt. 10c; ½ oz. 20c; oz. 30c; ¼ lb. 85c; 1 lb. \$3.00 5 to 25 lbs. \$2.90 per lb.

WALTHAM 29 (Medium). (75 days from plants.) Low growing, heavy, short compact plant, which produces a large, central head. The plant is heavily branched with dark green, tight buds that are of medium size, evenly distributed. It has ability to survive periods of hot weather without the heads opening up. Because of its uniformity, a large portion of the crop matures at one time. Its excellent dark green color, fine head and bud texture and high yield make it a valuable addition for Florida. Excellent for fresh market and freezing.

Pkt. 10c; ½ oz. 25c; 1 oz. 40c; ¼ lb. \$1.15; 1 lb. \$4.00 5 to 25 lbs. \$3.90 per lb.



Waltham 29

CABBAGE

Sow 6 ounces in seedbed to plant an acre or 12 ounces per acre

For the garden, sow 1 packet in seedbed for 50 plants to set 60 feet of row.



A typical plant of Marion Market (Yellows Resistant) Cabbage

Culture. Cabbage is a heavy feeder and should be grown rapidly. For maximum yields on sandy loam soils 1500 to 2000 lbs. per acre of 4-7-5 fertilizer mixture should be applied in the row a few days prior to setting plants. In addition, two or three side dressings of Nitrate of Potash or Nitrate of Soda should be made at three to four week intervals. Where planted in seedbeds, which is the desirable and common practice, plant seed thin in drills 6 inches apart, cover 1/4 to 1/2 inch deep, using 8 ounces of seed for an acre of plants, and do not make the soil in seedbeds as rich as soil where plants are to be set. When seed is planted in the field, use 12 to 16 ounces of seed per acre. Make rows $2\frac{1}{2}$ to 3 feet apart and thin to 12 to 18 inches apart in row, depending on variety and size of head you wish to produce. For a winter crop of cabbage in Florida seed is sown in September and early October. For an early spring crop plant in November or early December. For the control of cabbage insects and diseases see page 65.

Our cabbage seed is grown in disease-free territory on the Pacific coast where Black Rot has never been found by plant pathologists who inspect our seed growing fields each year.

The number of days in parentheses after each variety indicates the relative number of days to marketable heads from setting out plants. From the time seed is sown until the plants are ready for setting, thirty-five days are usually required.

> Prices quoted are postpaid. Write for prices on larger quantities.

EARLY ROUND DUTCH. (71 days.) A hardy, and dependable variety of round headed cabbage. Heads are firm and of high quality. Plants vigorous, small, and compact, with erect leaves and short stems. Suitable for close planting, but must be fertilized heavily if planted close to produce size. Withstands cold weather. Especially suited for muck land production.

Pkt. 10c; oz. 35c 1/4 lb. \$1.00; 1 lb. \$3.50 5 to 25 lbs. \$3.40 per lb.

GREENBACK (Yellows Resistant). (72 days.) This cabbage is in the round Dutch class, about as early as Marion Market, but darker green and more compact. Plant is vigorous and short stemed.


> Pkt. 10c; oz. 40c; 1/4 lb. \$1.25; l lb. \$4.50 5 to 25 lbs. \$4.40 per lb.

MARION MARKET (Yellows Resistant). (75 days.) This is a late maturing disease-resistant strain of Copenhagen Market, highly resistant to Yellows. Larger and coarser in growth than Copenhagen Market but makes somewhat heavier yields, with large, firm, solid heavy heads.

> Pkt. 10c; oz. 35c; 1/4 lb. \$1.00; 1 lb. \$3.50 5 to 25 lbs. \$3.40 per lb.

MIDSEASON MARKET. (75 days.) The heads are of medium size, hard, globe-shaped on short stems. Leaves are dark green, thick and leathery, making this variety hardy and resistant to cold damage, and also making it a good shipper in bags, the heads not being bruised so readily as other varieties.

> Pkt. 10c; oz. 30c; 1/4 lb. 80c; 1 lb. \$2.75 5 to 25 lbs. \$2.65 per lb.



Golden Acre

GOLDEN ACRE. (64 days.) An extremely early, small to mediumsized round headed cabbage of the Copenhagen Market type, but con-siderably earlier than most stocks of Copenhagen Market. Our strain of this variety heads up very uniformly and is an exceptionally good strain.

Pkt. 10c; oz. 30c; 1/4 lb. 85c; 1 lb. \$3.00 5 to 25 lbs. \$2.90 per lb.

COPENHAGEN MARKET (No. 86). (68 days.) Our early strain of Copenhagen Market is very popular in all the large cabbage producing sections of Florida. It is short-stemmed, compact with very solid, round, medium-sized heads, having few outer leaves, thus facilitating close planting, very uniform in shape, and matures practically all at the same time. This uniform maturity is desirable from a marketing standpoint.

> Pkt. 10c; oz. 30c; 1/4 lb. 85c; 1 lb. \$3.00 5 to 25 lbs. \$2.90 per lb.



Copenhagen Market No. 86

CABBAGE (Continued)

BADGER MARKET (Yellows Resistant). (72 days.) This new yellows resistant cabbage is showing much



Badger Market

promise on Florida muck soils. On sandy soils it is inclined to be too small unless heavily fertilized. In season it is only a few days later than Early Copenhagen Market No. 86. It is extremely uniform as to type and time of maturity. It makes a uniform, small, compact, round head, of attractive dark bluegreen color, very firm and hard. even on loose muck, with a short to medium core. It is very resistant to splitting, and can be left in the field for ten days to

two weeks after it is mature and shows no signs of splitting. It possesses excellent edible quality. This new variety is strongly recommended for Florida muck and the better sandy loam soils where it makes heads two and one-half to three pounds in weight, just the right size for northern markets.

> Pkt. 10c; oz. 40c; 1/4 lb. \$1.15; 1 lb. \$4.25 5 to 25 lbs. \$4.15 per lb.



Early Glory

EARLY GLORY. (72 days.) A medium-early, round headed type. Somewhat similar to Copenhagen Market, but slightly later, larger, with a darker green color and a heavier yield. The heads are protected by large, thick, overlapping wrapper leaves, thus resisting cold damage. Heads are solid, of medium size, and of excellent shipping and market quality. An improved early maturing strain of the old Glory of Enkhuizen.

Pkt. 10c; oz. 30c; 1/4 lb. 85c; 1 lb. \$3.00 5 to 25 lbs. \$2.90 per lb.

RED ACRE. (72 days.) This variety fills a long felt need in being the only red cabbage that matures very early, only a few days later than Early Copenhagen Market. The plant is small, compact and makes goodsized round, hard, deep red colored heads. Very uniform as to type. Especially desirable for muck land production in Florida, but also does exceedingly well on fertile sandy loams.

> Pkt. 10c; oz. 35c; 1/4 lb. \$1.00; 1 lb. \$3.50 5 to 25 lbs. \$3.40 per lb.

STEIN'S FLAT DUTCH. (90 days.) This is an old standard, hardy, winter variety. It produces a very large-sized plant, spreading in growth, with few large, broad outer leaves. The head is of large size, flat, and very solid.

> Pkt. 10c; oz. 25c; 1/4 lb. 75c; 1 lb. \$2.50 5 to 25 lbs. \$2.40 per lb.

JERSEY WAKEFIELD. (62 days.) Very early and resistant to cold weather. The small, medium-sized conical-shaped, pointed heads are very firm, of finest quality, and especially desirable for home use.

> Pkt. 10c; oz. 30c; 1/4 lb. 85c; 1 lb. \$2.75 5 to 25 lbs. \$2.65 per lb.

CHARLESTON WAKEFIELD. (70 days.) The heads are larger and broader at the base, not quite so pointed and later in maturing than Jersey Wakefield. Of uniform size, very solid and of desirable market quality. Resistant to cold weather and premature seeding.

> Pkt. 10c; oz. 30c; 1/4 lb. 85c; 1 lb. \$3.00 5 to 25 lbs. \$2.90 per lb.

CHINESE CABBAGE


Also known as Celery Cabbage. Plant 1 pound per acre.

In the garden make two plantings at different times. 1 packet to 40 feet of row at each planting.

Culture. This is a cool-weather plant and should not be grown for a late spring or summer crop, because it bolts to seed in hot weather. Seeds should be planted in 2½ to 3 foot rows where the crop is to mature, thinning out as the plants develop to stand 8 to 12 inches in the row. Chinese Cabbage makes excellent cole slaw.

MICHIHLI. (70 days.) From seed planting to edible stage. Sure header, maturing uniformly into long, slender, very firm, well blanched, crisp, tender, and sweet heads. It is slightly more vigorous and darker green than other strains, the heads are more uniform. and slower to bolt to seed. The heads become 19 to 20 inches tall, 3½ to 4 inches thick, tapered near the tip.

Pkt. 10c; oz. 25c; 1/4 lb. 75c; 1 lb. \$2.50



CANTALOUPE


Plant 1 to 2 pounds per acre.

In the garden plant 1 packet to 50 feet row.

Culture. General cultural requirements, diseases and insects and their control, similar to those of cucumber. (See page 19.) Plant in hills 4 to 5 feet apart, 5 to 6 seeds to a hill. Thin out to two plants per hill when 4 inches high. Plant in 5 foot rows, thinning to 2 feet in the row. Apply 600 lbs. Fertilizer 10 days before planting, side dress with 400 to 600 lbs. at vining. For earlier melons, use **Hotkaps.** (See page 88.) For insects and disease control (See page 66). For protection from rats and birds use Kilgore's Melon Seed Treatment. (See page 76.)

Prices quoted are postpaid.

Write for special prices on larger quantities.



Edisto

EDISTO. (85 days.) This new variety developed by M. B. Hughs at the Edisto Branch of the South Carolina A.E.S., has good resistance to powdery mildew and alternaria and considerable tolerance to downy mildew. Melons average about a third larger than Jumbo Hale's Best. The rind is hard and the flesh firm, withstanding long distance shipping. The flesh is very thick, with small seed cavity, ripening to an attractive deep salmon color. Excellent flavor and texture.

> Pkt. 10c; oz. 35c; 1/4 lb. 75c; 1 lb. \$2.50 5 to 25 lbs. \$2.40 per lb.

SMITH'S PERFECT. (90 days.) A strong, vigorous vine growth, withstanding wind and cold damage. Very resistant to downy mildew disease commonly called rust. Also resists aphids. Fruits average 4 lbs. Round in shape, slightly flattened at stem and blossom ends, free of ribbing, a tough rind; makes a good shipper and market melon. For market, it is best to harvest them a little on the green side just as they begin to turn yellow. The flesh is firm, very thick, with a small seed cavity, of attractive deep orange or salmon color, and has a delicious flavor.

> Pkt. 10c; oz. 25c; 1/4 lb. 75c; 1 lb. \$2.50 5 to 25 lbs. \$2.40 per lb.

JUMBO HALE'S BEST. (78 days.) A large fruited strain of Hale's Best, grown very extensively in Georgia and other southern states. Fruits are oval in shape, showing some trace of ribbing, with heavy coarse netting. Average about 4 to 5 lbs. Flesh thick, relatively small seed cavity, deep orange color.

Pkt. 10c; oz. 20c; 1/4 lb. 60c; 1 lb. \$2.00 5 to 25 lbs. \$1.90 per lb.

HALE'S BEST, No. 36. (75 days.) An improved strain of Hale's Best, developed for uniformity in shape and complete netting. Fruits are round in shape, with very heavy, coarse netting, and average about three pounds. Flesh thick, small seed cavity, deep salmon color, very sweet, fine quality. Excellent shipper.

Pkt. 10c; oz. 20c; ½ lb. 60c; 1 lb. \$2.00

2 to 5 lbs. \$1.90 per lb.

NO. 45. (80 days.) Attracting much favorable attention in Florida because of its high yielding, good shipping and holding quality. Vines moderately large and produce a very heavy crop. Highly resistant to powdery mildew, but not to downy mildew. Medium sized fruits (about 3½ lbs.) are short-oval, indistinct ribbing and well netted. Flesh thick, firm, sweet and of good quality, light orange color. Can be picked almost mature for local markets or distant shipping.

Pkt. 10c; oz. 20c; 1/4 lb. 60c; 1 lb. \$2.00 5 to 25 lbs. \$1.90 per lb.

RIO GOLD. Resistant to downy and powdery mildew and to aphids. Comparable in size to Imperial 45, has a greenish-yellow rind with well raised net. Flesh is deep and a bright orange color. Of excellent flavor and sweetness. A very productive melon and valuable for its resistance.

Pkt. 10c; oz. 25c; 1/4 lb. 65c; 1 lb. \$2.25 5 to 25 lbs. \$2.15 per lb.

CAULIFLOWER

Sow 6 ounces in seedbed to plant an acre. In the garden sow one packet in seedbed for thirty plants. Culture. Plant in seedbeds from Sept. to Dec. and cultivate the same as cabbage, spacing 3 to 4 feet between rows and 18 to 24 inches in row. Cauliflower needs rich soil and plenty of moisture. When head develops to the size of an apple, draw outer leaves over head and tie with soft cord to produce attractive, white, solid heads. Our cauliflower seed is grown in black rot free territory.

Prices quoted are postpaid. Write for special prices on larger quantities.

SUPER SNOWBALL. (55 days from plants.) An early maturing strain. Snowball type, in which curled formation of leaves over head protects it from sun and from freezing. Dwarf, compact plants are strong, vigorous. Heads are large, smooth, solid, pure white and very uniform; recommended as an early winter crop to mature in cold weather. Susceptible to "whiptail" and should be grown in alkaline soils with high pH to avoid trouble.

Pkt. 25c; ½ oz. \$1.00; oz. \$1.50; ¼ lb. \$4.00; 1 lb. \$15.00

EARLY SNOWBALL (Snowdrift). (62 days from plants.) Vigorous plants grow larger and taller than Snowball, are more compact and upright in growth with long leaves which protect head or curd in its early growth. Large, snow-white attractive heads are very deep. Heavy and firm. An ideal variety for early spring production in Florida. Free of leafy and ricey heads. Very resistant to "whiptail" disease.

Pkt. 25c; ½ oz. \$1.00; oz. \$1.50; ¼ lb. \$4.00; 1 lb. \$15.00



Early Snowball

CARROT

Plant 4 pounds of seed per acre. In the garden make three plantings at different times.

Plant 2 packets to 100 feet of row at each planting.

Culture. Sow in good, deeply worked, fine soil, enriched with 800 to 1200 pounds per acre of a good vegetable fertilizer, in drills 1½ to 2 feet apart, and cover seed ¼ to ½ inch deep. When 2 inches high, thin out 2 to 3 inches apart. Carrots can be planted in Florida from August to March.

Prices quoted are postpaid.

Write for special prices on larger quantities.

LONG IMPERATOR. (77 days.) The roots average about 10 inches in length, with a shoulder diameter of 1½ to 1¾ inches, smooth, uniformly tapered to a semiblunt end, deep rich orange color, both exterior and interior with comparatively inconspicuous core. The flesh is of rich orange color, extending to the center of the root, with an indistinct core; fine grained, tender and of excellent quality. The tops are of medium size, but strong enough for good bunching. For shipping quality, attractiveness and popularity in the market. Long Imperator is superior to other long-rooted varieties.

Pkt. 10c; oz. 30c; 1/4 lb. 85c; 1 lb. \$2.75 5 to 25 lbs. \$2.65 per lb.

RED CORE CHANTENAY. (72 days.) The roots are tapered, less stump rooted, and slightly longer than the old Chantenay, averaging about 6 inches. The interior is solid orange-colored without the usual lemon color in the core. The flesh is very tender and sweet, which with the excellent outside color and shape combines to make this variety valuable for the market.

Pkt. 10c; oz. 20c; 1/4 lb. 50c; 1 lb. \$1.75 5 to 25 lbs. \$1.65 per lb.



Developed to meet the need for a better market type.



Comparison of root characteristics of carrot varieties. From left to right: Imperator, Nantes (Coreless), Danvers Half Long,
Red Core Chantenay.

DANVERS HALF LONG. (75 days.) Our stock of this variety is half stump or semi-pointed, from 6 to 7 inches in length. The shoulder at the crown is from 1½ to 1¾ inches thick, and tapers uniformly to a thickness of about ¾-inch just where the root begins to taper to the tip. The tops are a little larger than those of Chantenay, and the crown is flat without a depression. The roots are smooth and attractive in appearance.

Pkt. 10c; oz. 20c; 1/4 lb. 50c; 1 lb. \$1.75 5 to 25 lbs. \$1.65 per lb.

NANTES IMP. CORELESS. (70 days.) Cylindrical, roots about 5½ to 6½ inches long and 1 to 1½ inches thick, very stump rooted with a small rat tail. The color is clear orange-red and the core is rather undefined, being of the same color as the flesh so that it is called "Coreless." The tops are small and short, but heavier than ordinary stocks of Nantes, and the root is very fine grained, tender, delicate, sweet and of highest edible quality. An excellent high quality variety for the home garden and local market.

Pki. 10c; oz. 20c; 1/4 lb. 50c; 1 lb. \$1.75 5 to 25 lbs. \$1.65 per lb.

LONG ORANGE. (86 days.) Our stock is the true thick-shouldered strain of Long Orange, 1¾ to 2 inches thick at the shoulders. 10 to 20 inches long, and tapering in a straight line to the tip. The tops are rather large. Grown in sections of Florida where a smooth, reddish-colored, long-rooted carrot is desired.

Pki. 10c; oz. 20c; 1/4 lb. 50c; 1 lb. \$1.75 5 to 25 lbs. \$1.65 per lb.



CELERY

Sow 1/2 pound in seedbed to plant an acre.

For the garden sow 1 packet in a well protected seedbed for 150 plants to set 50 feet of row.

Prices quoted on celery seed are postpaid. Write for special prices on larger quantities.

Culture of Celery. First sowings in Florida are made the latter part of June and continue to December. The main sowings for a winter crop are usually made in July or early August, and for the spring crop, grown principally on muck, they are usually made in October and early November. Celery is always started in protected seedbeds. The soil must be put in perfectly fine condition and made into beds 4 to 6 or more inches high, to prevent overflow in case of heavy rains. Sow the seed broadcast or in rows 6 inches apart, using 1 pound of seed to 125-150 yards of bed 3 feet wide. The seed should be barely covered with soil.

We recommend use of burlap or white muslin as a cover over the seedbed stretching it on a tent-shaped or slanting frame built with the sides about 12 inches from the ground to provide ample circulation of air. This cover will serve as shade and break the force of heavy rains. After the seed is sown, sprinkle with a very fine spray in order to keep the soil cool and moist, which will increase the germination of the seed during the hot summer months. With surface irrigation, water should be circulated continuously around the beds.

When plants in seedbed are 6 inches high, transplant to the field in double rows 3 to 3½ feet apart, with 7 inches between the double rows, and set the plants 4 to 5 inches apart in the row. The plants may be set in single rows 3 feet apart, allowing 4 to 5 inches between plants in the row. The Pascal varieties should always be planted at least 6 inches in the row. To grow fine celery, a large supply of moisture and fertilizer is essential from the time the seed is sown in the seedbeds until the crop is ready for harvest.

For the control of insects and diseases of celery see page 65.

Good control of black heart in celery has been obtained by spraying weekly with 10 lbs. Calcium Nitrate dissolved in 100 gal. water per acre.

The number of days in parentheses after each variety indicates the relative number of days to marketable stage or harvest from setting out plants. It usually requires about seven or eight weeks to grow plants large enough to be set in the field.



Summer Pascal



Florida Green Pascal

FLORIDA GREEN PASCAL. (125 days.) A very vigorous, compact-growing variety, with broad, semi-round, medium-long, thick, very meaty, smooth, attractive ribs. The ribs or petioles will average from 7½ to 8 inches to the first joint and plants will range from 20 to 23 inches overall. The ribs or stalks are of ideal edible quality, being tender, brittle, stringless and crisp.

This stock is not subject to premature seeding, and can be grown at any time during the fall, winter or spring in Florida.

Pkt. 15c; ½ oz. 40c; 1 oz. 75c; ¼ lb. \$2.35; 1 lb. \$8.50

TALL GOLDEN SELF BLANCHING. (115 days.) One of the best varieties for a mid-winter crop in Florida, because it withstands cold weather better than most varieties of the Special or Golden Plume type. Selected for uniformly long, thick, round ribs, large full hearts of rich golden yellow color, and for heavy yields of good sizes. The ribs average eight to ten inches in the first joint, rounded and exceptionally thick. The plants range from 21 to 24 inches over all, making an attractive pack. It is intermediate between the Old Golden and Special or Golden Plume types, possessing the desirable characteristics of both types. Recommended especially for a Florida mid-winter crop.

Pkt. 15c; ½ oz. 40c; 1 oz. 75c; ¼ lb. \$2.35; 1 lb. \$8.50

SUMMER PASCAL (Waltham Strain). (115 days.) This is an early maturing Green Pascal variety. A rapidly growing type of celery with long, upright compact ribs or petioles, making a very attractive appearing celery for the markets. The ribs are thick and semi-rounded, but not quite as smooth as our strain of Florida Green Pascal, although the Waltham strain will mature about a week earlier, and makes a high yield of large sizes. This strain must be cut promptly when ready or it will tend to go down in the field after maturity.

Pkt. 15c; ½ oz. 40c; 1 oz. 75c; ¼ lb. \$2.35; 1 lb. \$8.50



COLLARD


Sow ½ pound in seedbed to plant an acre, or plant 2 pounds per acre in field.

For the garden make two sowings at different times. Sow one packet in seedbed for 35 plants for each planting to get 50 feet of row.

Culture. Collard is an old standby for winter greens all over the South. May be started in seedbeds or direct in the field, in rows 2½ to 3 feet apart, with plants 14 to 18 inches apart in the row. Use about 1200 pounds of fertilizer per acre before setting plants. Collard can be grown every month of the year in Florida, but it grows best and produces the highest edible quality during cool weather. This vegetable should be included in every Florida family garden, as it is very hardy, easy to grow and will furnish an abundance of greens, high in health-producing vitamins.

The number of days in parentheses after each variety indicates the time from setting out plants to harvest. It requires about a month in the seedbed to get plants from seed sowing to transplanting stage.

Prices quoted are postpaid.
Write for special prices on larger quantities.



Georgia 912

GEORGIA 912. (50 days.) A non heading type of the cabbage family, very extensively grown for greens in home and commercial gardens in Florida. This is a distinct strain of Georgia type collard, developed at the Associated Seed Growers' Southeastern Breeding Station at Bradenton, Fla. through inbreeding and progeny testing. Compared to the standard Georgia type it is slower to bolt or to seed; has more compact plant habit, with shorter petioles and broader leaf blades, and has more savoyed leaves. Very uniform plant type and a heavy yielder.

Because of the more compact plant habit, broader and more savoyed leaves, good uniformity of stock, cold hardiness and long standing habit. Georgia 912 should find a place for commercial shipping and processing, as well as for general use in collard growing areas.

Pkt. 10c; oz. 15c; 1/4 lb. 40c; 1 lb. \$1.20 5 to 25 lbs. \$1.10 per lb.

GEORGIA OR SOUTHERN. (50 days.) Plant grows from two to three feet tall, is erect and spreading, with many large, undulated leaves on a white stem. Withstands cold as well as heat, and adverse soil conditions. A high yielding hardy type, of excellent quality when cooked. This is the old standard variety for Florida and the South.

Pkt. 10c; oz. 15c; 1/4 lb. 40c; 1 lb. \$1.20 5 to 25 lbs. \$1.10 per lb.



Vates

VATES. (55 days.) A very large, vigorous, rather low growing plant, not subject to wind damage, with very large, broad, heavy, thick leaf blades, the leaf blade developing from the base of the short stem or petiole.

The leaves are smooth with an attractive deep, dark green color, the stems and leaves never turning yellow or purplish even in the coldest weather or on poor soils. It retains this deep green color in shipment and on the markets. A heavy yielder, slow to bolt to seed in cold or hot weather. Uniform as to plant type.

Pkt. 10c; oz. 15c; ¼ lb. 45c; 1 lb. \$1.40 5 to 25 lbs. \$1.30 per lb.

ASGROW VIGORPAK SEED

Seed of superior vigor • Scientifically pre-conditioned
• Packed in airtight cans

For many years the ASGROW RESEARCH DEPARTMENT has studied the factors affecting the life of vegetable seed. It was found that, within limits, the drier the seed the longer it will keep its vitality. The higher the moisture content of the seed and of the surrounding atmosphere, the more rapidly will the seed lose its capacity to germinate and grow into healthy plants. From these studies ASGROW has perfected a way to process and package seeds so that they will keep their initial vitality for years under almost any storage conditions, short of killing temperatures.

The practical application of these investigations is in ASGROW VIGORPAK SEED. The internal moisture of the seed is brought to a safe level for that particular kind of seed, as determined by the ASGROW RESEARCH DEPART-MENT. Then, after treatment with an adequate seed protectant, the seeds are packed in moisture-proof cans and hermetically sealed. So long as it is in the unopened can, ASGROW VIGORPAK SEED is protected not only from the adverse effects of high humidity but also from damage by insects and vermin and from loss by spilling or mixing. With these hazards eliminated, seed can safely be carried over to the following season, provided the can has not been opened.

YOUR INSURANCE AGAINST

Loss in germination and vigor ● Loss from pests ● Waste from mixture

FIELD CORN

Plant 7 pounds ($\frac{1}{2}$ peck) per acre or 56 pounds (1 bushel) to 8 acres.


To increase yields with field corn in Florida two things are essential, namely closer spacing and ample fertilizer. On good land 20 to 26 inch spacing in 38-inch rows is best with 500 lbs. per acre of fertilizer at planting time followed by side dressings of nitrate of soda up to 300 lbs. per acre about 5 weeks after planting.

The number of days indicated after each item represents the time from seed planting to harvest under average spring conditions in Florida.

Write for special prices on larger quantities.
All prices quoted are subject to change without notice

FUNK'S G-740 HYBRID. (120 days.) This new yellow corn has given excellent results in numerous commercial trials in the Everglades muck and offers fine possibilities for other sections of the state. It consistently makes high yields (two ears per stalk) of attractive, medium large ears, well covered to the tips with hard, semi-flint yellow kernels, with extra heavy shuck, very weevil resistant and of excellent keeping quality. Makes strong, sturdy stalks which stand drouth better than many others.


1/4 lb. 25c; 1 lb. 60c; 2 lbs. \$1.00, postpaid. Not prepaid, 14 lbs. (pk.) \$3.25; 56 lbs. (bu.) \$12.20



Funk's G-740 Hybrid

IMPROVED GOLDEN DENT. (86 days.) A wonderful yielder, and does splendidly here in Florida. It is planted extensively for early use, but is too soft for a general crop. Has large, deep, broad grains, and is desirable for early feed.

1/4 lb. 15c; 1 lb. 45c; 2 lbs. 75c; postpaid Not prepaid, 14 lbs. (pk.) \$1.75; 56 lbs. (bu.) \$6.50



Dixie 18 Hybrid

DIXIE 18 HYBRID (Certified). (68 days.) Dixie 18 is the best yellow hybrid corn for central and north Florida and south Georgia; a heavy yielder, averaging 2 ears per plant. Yields of 80 to 100 bushels per acre have been reported. The plants are relatively short, sturdy and very vigorous, with a deep root system, resisting wind damage, lodging and drouth. The ears are uniform and borne low down on the plant. It makes a semi-hard corn and is very desirable for "hogging off." The grain is yellow and intermediate in texture being harder than Whatley. Has good resistance to weevil.

½ lb. 15c; lb. 45c; 2 lbs. 75c; postpaid
 Not prepaid, 10 lbs. (pk.) \$2.25; 56 lbs. (bu.) \$8.50

Roasting Ear Corn

Plant 7 pounds ($\frac{1}{2}$ peck) per acre. In the garden two plantings at different times. Plant $\frac{1}{4}$ pound to 200 feet of row.

Write for special prices on larger quantities.

OKLAHOMA SILVERMINE. (82 days.) An exceedingly heavy yielder of very large, heavy ears. The cobs are covered with 14 to 18 rows, and run from 11 to 12 inches in length. Very resistant to worm damage. An excellent variety for shipping, with a heavy dark green husk that holds its color in shipment and on the market. This is the standard roasting ear variety grown in Central Florida for shipment to northern markets.

1/4 lb. 15c; 1 lb. 50c; 2 lbs. 85c; postpaid Not prepaid, 14 lbs. (pk.) \$2.00; 56 lbs. (bu.) \$7.00

SOUTHERN SNOWFLAKE. (80 days.) The ears are large, 10 to 12 inches long, well filled with 16 rows of large, broad kernels, and of very attractive appearance. The long tight husks protect the ears from worms.

1/4 lb. 15c; 1 lb. 50c; 2 lbs. 85c; postpaid
 Not prepaid, 15 lbs. (pk.) \$1.75; 56 lbs. (bu.) \$6.50

NORFOLK MARKET (Improved Trucker's Favorite). (68 days.) An extremely early variety. Ears are medium large, averaging about 10 inches long, with 12 to 14 rows of large, white kernels, very tender, and of splendid eating quality. Especially valuable for planting after winter truck crops.


½ lb. 15c; l lb. 50c; 2 lbs. 85c; postpaid Not prepaid, 14 lbs. (pk.) \$2.00; (bu.) \$7.00

SWEET CORN

15 pounds seed required per acre. In the garden make two plantings. Plant 1/4 lb. to 200 feet of row at each planting.

Our hybrid Sweet Corn seed is now graded and sized for better, more accurate stands and more uniform maturity.

Culture. Rows should be 2½ to 3 feet apart with the plants 8 to 12 inches depending on type and season. Sweet corn requires much more fertilizer than field corn. Use about 1000 lbs. per acre of a good commercial fertilizer mixture high in potash. When corn is about knee high apply 200 lbs. per acre of sodium nitrate or its equivalent. This fertilizer practice will greatly increase the yield. Early plantings are made in Florida from January 1st until March. For control of corn ear worms and leaf blight see page 66.



Golden Security

GOLDEN SECURITY. (76 days.) This variety is one of the most popular with Florida farmers. A high yielding hybrid, suited for fall, winter, or spring planting. The stalks are very vigorous, averaging 6½ ft. tall with dark green foliage. The ears are 7½ in. to 8½ in. long with 14-16 rows of light yellow kernels of excellent quality. No grower should overlook this variety.


½ lb. 25c; l lb. 80c; 2 lbs. \$1.35, postpaid
 Not prepaid, 10 lbs. \$5.20; 25 lbs. \$12.50; 50 lbs. \$24.50

FLORIDA 104 (Iobelle). (76 days.) This variety was developed by Dr. E. S. Huber of Iowa State College, Ames, Iowa and has been thoroughly tested under Florida conditions. The ears are about 7½ in. to 8½ in. long, with 14 to 16 rows of high quality, light yellow kernels. Particularly adapted to cool weather production on the muck soils.

1/4 lb. 25c; 1 lb. 80c; 2 lbs. \$1.35, postpaid Not prepaid, 10 lbs. \$5.10; 25 lbs. \$12.25; 50 lbs. \$24.00

GOLDEN CROSS BANTAM. (73 days.) Golden Cross Bantam is a hybrid sweet corn that has done well as a high quality, yellow-kerneled, early variety in Florida. The small, short stalk ($5\frac{1}{2}$ to 6 feet) is very sturdy and vigorous, suckers quite badly. Ears 6 to 7 inches long, 10 to 14 rows.

 $^{1}\!\!/_{4}$ lb. 25c; 1 lb. 80c; 2 lbs. \$1.35, postpaid Not prepaid, 10 lbs. \$4.10; 25 lbs. \$9.75; 50 lbs. \$19.00



Sixtypak


SIXTYPAK. (74 days.) This hybrid developed by Associated Seed Growers, Inc., has been thoroughly tested with other leading varieties of hybrid sweet corns in all important corn growing sections of Florida. In every test "Sixtypak" produced a higher yield than any other variety, and showed a marked resistance to Helminthosporium Leaf Blights.

The plants are uniform, about six and one-half feet tall with heavy dark green foliage, a high percentage producing two fancy ears per stalk. The ears are 7½ to 8 inches long, well filled to the tip, with an average of 14 rows of golden yellow kernels. The husk and broad flags are dark green, making a very attractive five dozen pack.

½ lb. 25c; l lb. 80c; 2 lbs. \$1.35, postpaid
 Not prepaid, 10 lbs. \$5.60; 25 lbs. \$13.50; 50 lbs. \$26.50

IOANA. (74 days.) A very productive, yellow-kerneled, hybrid. Vigorous plants are highly resistant to drouth. Medium tall, 6 feet, sturdy, with large, broad and numerous dark green leaves. Ears 7½ to 8 inches long, 12 to 14 rowed, well-filled with medium wide attractive light yellow kernels of fine quality and flavor, making it highly resistant to worms. Husks retain green color for a long time, making for fresh, attractive market appearance.

1/4 lb. 25c; 1 lb. 80c; 2 lbs. \$1.35, postpaid
 Not prepaid, 10 lbs. \$4.00; 25 lbs. \$9.50; 50 lbs. \$18.50



Ioana



A field of Marketer (Early Green Market)

CUCUMBER

In the garden plant 1 packet to 50 feet of row. Plant 2 pounds per acre.

Culture. Rich, moist, sandy loam soil is the best for growing cucumbers. The soil should be well fertilized with a good vegetable fertilizer applied at the rate of 1200 lbs. per acre a week or ten days before planting; 600 to 1000 lbs. is generally used as a side-dresser just before the plants start to run. Fall plantings are made in late August and in September. Spring planting in Florida begins early in January and continues through February. Plant in rows 5 to 6 feet apart, 6 inches apart in the row, and thin to 1 to 2 feet between every two plants in the row. Sometimes it is difficult to get a stand early if the ground is wet and cold, as the seed will rot before it germinates. Therefore seed should be treated. (See page 64).

If young plants are very badly damaged with frost or cold, it is best to plant over. When young plants are stunted they never make a normal crop and usually not much earlier than if they were planted over. A row of oats planted between every two rows of cucumbers makes a good windbreaker.

It is very necessary to have hives of bees in or near cucumber fields to insure pollination and a good set of fruits. The bees should not be placed in the field until the cucumbers begin to blossom, in order to provide food for the bees so they will remain in the cucumber fields. Without proper pollination the cucumbers are deformed, and a large percentage of bottlenecks and nubbins are produced.

For the control of insects and diseases of cucumbers (see page 66). To control angular leaf spot treat seed

with a mercuric chloride. (See page 64.) The best control of downy mildew, sometimes called "rust," is Nabam spray or Zineb Dust. (See page 66.) The most effective and safest control for aphids and some other cucumber insects is Lindane spray of Kilgam dust. (See page 66.)

For protection from rodents and birds use Kilgore's Melon Seed Treatment. See page 76.)

The number of days after each variety represents the time from seed planting to first picking under average spring or fall conditions in Florida.


Prices quoted are postpaid.

Write for special prices on larger quantities.

MARKETER (Early Green Market). (60 days.) The leading variety for yield and quality. Holds its dark green color much better than most varieties, when grown in late spring or early fall. It averages about 8 inches in length, and is especially desired by the eastern and southern markets where a medium size cucumber is preferred. The fruits are exceptionally uniform in size, shape and color, with practically no jumbos or oversize even when not harvested for several days. Stands heat of late spring better than other varieties. Exceedingly prolific, throwing an exceptionally large number of female blossoms per plant. It will average over 90 per cent fancies. Buyers pay a premium for this cucumber, when other varieties go begging at lower prices.

Pkt. 10c; oz. 25c; 1/4 lb. 65c; 1 lb. \$2.25 5 to 25 lbs. \$2.15 per lb.

CUCUMBER (Continued)




Ashley

ASHLEY. (55 days.) A new downy mildew resistant cucumber developed at Clemson Truck Experiment Station, South Carolina. Vigorous vines and a heavy producer. Fruit dark green color, about 8 inches long, tapering at the stem end.

> Pkt. 10c; oz. 25c; 1/4 lb. 75c; 1 lb. \$2.50 5 to 25 lbs. \$2.40 per lb.

STRAIGHT-8. (47 days.) This plant is vigorous in growth and very prolific. Fruits cylindrical, blunt at both ends. Exterior medium green. Recommended for late fall planting in Florida.

> Pkt. 10c; oz. 25c; 1/4 lb. 65c; 1 lb. \$2.25 5 to 25 lbs. \$2.15 per lb.



Puerto Rico No. 39

PUERTO RICO NO. 39 (Special Long Strain). (50 days.) This variety is highly resistant to downy mildew. The vines are rather compact in growth with fruits averaging about 7½ inches in length, dark green

> Pkt. 10c; oz. 25c; 1/4 lb. 65c; 1 lb. \$2.25 5 to 25 lbs. \$2.15 per lb.

PALMETTO . (60 days.) This variety is highly resistant to downy mildew, the most serious cucumber disease of the South. Palmetto will survive and produce a good crop, when other varieties may go down with mildew under cool, wet weather conditions.

> Pkt. 10c; oz. 25c; 1/4 lb. 75c; 1 lb. \$2.50 2 to 25 lbs. \$2.40 per lb.

OHIO MR 17. (40 days.) Highly tolerant of cucumber mosiac, replacing National Association and other susceptible varieties in areas where mosaic is prevalent. Vines vigorous and productive. Fruits longer and darker colored than National Association Pickling but comes well within acceptable range for type.

> Pkt. 10c; oz. 20c; 1/4 lb. 60c; 1 lb. \$2.00 5 to 25 lbs. \$1.90 per lb.




Stays Green (Pickling Type)

STAYS GREEN. (Pickling Type). (40 days.) This variety should be grown only for pickles, not for slicing or as a fresh market cucumber. It is very early and exceedingly productive of short, dark green warty cucumbers most desirable for dill pickles.

> Pkt. 10c; oz. 20c; 1/4 lb. 50c; 1 lb. \$1.75 5 to 25 lbs. \$1.65 per lb.

PALOMAR. (55 days.) A downy mildew resistant cucumber developed by crossing Palmetto and Marketer. Vines are heavy and vigorous. Extra early and a heavy yielder. Fruit about 8½" long, dark green color. Some fruits are slightly curved.

> Pkt. 10c; oz. 30c; 1/4 lb. 85c; 1 lb. \$2.75 5 to 25 lbs. \$2.65 per lb.



Palomar

EGGPLANT

Sow 1/4 pound in seedbed to plant an acre.

For the garden sow one packet in seedbed for 17 plants to set 50 feet of row.

Culture. For a spring crop, plant seed in November, December, and January; for fall crop, plant in June, July, and August, in muslin or burlap-covered seedbeds. Set in rows 4 to 5 feet apart and 2½ to 3 feet between the plants. Eggplant being a long season crop requires several fertilizer applications. Best results are obtained by applying half or more of a complete fertilizer at planting time and the remainder when the crop is one-third to one-half grown. The initial application should be made in two bands each located 2 or 3 inches below and 3 inches to the side of the plant row. Subsequent application should be drilled close to the plant row.


Top-dressing applications of nitrogen or a combination of nitrogen and potash throughout the season is necessary to hold up size and color of fruit. For the control of insects and diseases of eggplant see page 66.

The number of days in parentheses after each variety indicates the time to first marketable fruits, from setting out plants in the field or garden. It usually requires six to eight weeks to produce plants for field setting.

Prices quoted are postpaid.
Write for special prices on larger quantities.

FORT MYERS MARKET. (83 days.) We introduced this variety, which is today one of the leading varieties grown in the South. Our breeding department has made special selections to improve the variety for uniformity in type of plant, size, shape and color of fruit. Produces immense yields of uniform, large-sized fancy fruits over a long period. The plants are of the high-bush type, with very tall, vigorous growth. The fruits are slightly elongated, cylindrical in shape, and they possess the intense deep black color so much in demand among buyers and in the market.

Pkt. 10c; oz. 75c; ½ lb. \$2.40; 1 lb. \$9.00 5 to 25 lbs. \$8.90 per lb.



Florida Market (Cook's Strain)

FLORIDA MARKET (Cook's Strain). (75 days.) Resistant to tip over, leaf blight and fruit rot, all diseases caused by phomopsis fungus which have become so severe in the state as to make eggplant production unprofitable in some sections. Makes vigorous growth, sturdy bush, holding the fruits well off the ground. It is extremely early and a heavy yielder, bearing over a long season. This makes it a very desirable variety for south Florida where it can be planted in early fall for production throughout the winter and spring months. The attractive fruits are cylindrical in shape and develop to a large size. They are of an attractive, dark color. After several years of selection on the original Florida Market variety, we have in this Cook's Strain a great improvement in color, shape and uniformity.

Pkt. 10c; oz. 75c; 1/4 lb. \$2.50; l lb. \$9.50 5 to 25 lbs. \$9.40 per lb.



A field of Fort Myers Market Eggplant at harvest time.



ENDIVE



Improved Florida Deep Heart

Sow 1/2 to 1 pound in seedbed to plant an acre, or 1 to $1\frac{1}{2}$ pounds per acre in the field.

For the garden sow one packet in seedbed for 50 plants to set 40 feet of row, or plant one packet to 40 feet of row.

Culture. Plant from August to February in seedbeds or in the fields or garden the same as lettuce. Set in rows 1½ to 2 feet apart and 9 inches apart in the row. Soil and fertilizer requirements are the same as for lettuce. (See page 23.)

The number of days indicated after each variety is the number of days from seed planting in the field or garden to edible or harvest stage. For insect control (see page 67).

Prices quoted are postpaid. Write for special prices on larger quantities.

ESCAROLLE—FLORIDA DEEP HEART. (70 days.) One of the best and most prolific types to grow commercially. In great demand on northern markets, and well adapted to Florida soil and climatic conditions. The leaves are large, broad, thick, slightly crumpled, of medium green color, with thick, nearly white midribs. Our improved stock of this variety is of the Full-Heart type, developing an upright growth, producing a very heavy, thick matted plant with large, deep, well balanced creamy white hearts.

Pkt. 10c; oz. 20c; ¼ lb. 60c; 1 lb. \$2.00

5 to 25 lbs. \$1.90 per lb.

PANCALIER (Pink Ribbed). (75 days.) A distinctive variety of curly-leaved type. Plants spreading, 12-15 inches, producing dense growth of deeply cut, curled dark green leaves with pink midribs, blanching well when planted closely.

Pkt. 10c; oz. 20c; 1/4 lb. 60c; 1 lb. \$2.00

5 to 25 lbs. \$1.90 per lb.

RUFFEC GREEN CURLED. (Green Ribbed). (75 days.) Sometimes called "Chicory." A hardy vigorous sort, with deep green, curled, finely cut leaves, giving a mossy appearance. The midrib is thick, broad, green, tender and fleshy. A dense mass of deeply divided leaves forms the head which blanches readily to a beautiful creamy white.

Pkt. 10c; oz. 20c; 1/4 lb. 60c; 1 lb. \$2.00 5 to 25 lbs. \$1.90 per lb.

DILL

A packet will sow 10 ft. of row; an ounce 50 ft. of row. Plant from November thru December

LONG ISLAND MAMMOTH. (65 days.) Partially matured seed clusters and flowerheads are useful for flavoring dill pickles. A tall growing plant, 2 to 2½ feet high, vigorous, with very finely divided leaves. Seed clusters are borne in umbels or clusters, have a strong aromatic odor, and mildly pungent flavor.

Pkt. 10c; oz. 35c; 1/4 lb. \$1.00; 1 lb. \$3.50

GARLIC

Plant 1 pound to 30 feet of row.

Garlic sets come in the form of bulbs which are divided into so-called sets or cloves, which are planted the same as onion sets, from August to November, in rows 15 to 20 inches apart, setting the cloves 3 to 4 inches apart in the row, being sure not to cover the tips of the cloves. Garlic is a southern crop and does well in

1/2 lb. 45c; 1 lb. 85c; 5 to 25 lbs. 75c per lb., postpaid

KALE

Plant 1 ounce to 100 feet of row, or 1 packet to 25 feet of row.

Culture. Kale is grown for greens. Plant in Florida from August to April, in rows 2½ feet apart, and for large plants thin from 20 to 30 inches apart in the row. The plants are extremely hardy, withstanding cold and heat.

VATES. (55 days.) A strain of the older Dwarf Blue Scotch, which it has largely superseded as a commercial crop for fresh market and processing. Plants low, spreading, hardy, very slow in bolting when grown over winter for spring harvest. Leaves tightly curled, dark bluish green. Holds color well in cold weather.

> Pkt. 10c; oz. 20 c; 1/4 lb. 50c; 1 lb. \$1.75 5 to 25 lbs. \$1.65 per lb.

KOHL RABI

For the garden make short row plantings at different times. Plant 1 packet to 25 feet of row.

Culture. Grow in row 11/2 to 2 feet apart, thinning from 6 to 8 inches apart in the rows. Planting every ten days will give a succession of tender, fleshy bulbs until hot weather. Start fall planting September, and successive plantings can be made until early spring.

WHITE VIENNA. (50 days from seed sowing to edible stage.) The fleshy stems or so-called bulbs, are formed above ground and are very light green, tender and delicate, and of best quality if used when about 2 to 3 inches in diameter, before fully grown.

> Pkt. 10c; oz. 25c; 1/4 lb. 75c; 1 lb. \$2.50 5 to 25 lbs. \$2.40 per lb.

LEEK

Sow 11/2 lbs. in seedbed to plant an acre, or plant 3 lbs. per acre in the field.

For the garden sow 1 pkt. in seedbed for 120 plants to set 50 feet of row, or plant 1 pkt. to 50 feet of row.

Culture. Leek seed may be planted any time during the fall or winter directly in the field, or preferably in seedbeds, in good rich soil. When plants in seedbeds are six to eight inches high, transplant to rows 18 inches apart, and 5 to 6 inches apart in the row, or thin out to this spacing in the field.

LARGE AMERICAN FLAG. (140 days from seed planting in field or garden to full edible stage.) The heaviest yielding and the best variety for Florida growing conditions. It makes larger and thicker, more fleshy stems averaging eight to ten inches long and 1½ inches thick.

Pkt. 10c; ½ oz. 25c; oz. 45c; ¼ lb. \$1.40; 1 lb. \$5.00 5 to 25 lbs. \$4.90 per lb.

LETTUCE

Sow ½ to 1 pound in seedbed to plant an acre, or plant 2 to 3 pounds per acre in field. For the garden make two plantings at different times. Sow 1 packet in seedbed for 60 to 80 plants to set 50 to 75 ft. of row at each planting.

Culture. First sowings are made in Florida the beginning of September and continued until March. Low, damp soil is best suited for this crop. Lettuce must be grown rapidly in order to give it size, tenderness, crispness and high edible quality. Therefore, the land must be highly fertilized and moist. Broadcast the seed on good moist seedbeds that have been thoroughly prepared, or sow in rows six inches apart and use a light roller to press seed into soil. Seedbeds should be kept thoroughly moist at all times after the seed is planted. In hot weather germination may be hastened and improved by placing lettuce seed in moist cloth in the ice box for several days before sowing. In level culture, lettuce plants are usually set 15 by 15 inches in the field, or in rows 1½ to 2 feet apart, and 10 to 15 inches in the row, or in double rows 16 inches apart, with 3½ to 4 feet from center to center of double rows, and 14 inches in the rows. Plants should be watered as set in field. For insect control see page 67.

Rusting of head lettuce and Romaine is usually due to boron deficiency. To prevent or stop this trouble, it is recommended to spray with 1 lb. borax to 50 gallons of water applied at the rate of not more than a total of 4 pounds borax per acre in two applications. Too much borax is toxic.

The number of days in parentheses after each variety represents the time required to develop marketable heads from the planting of seed.

Write for special prices on larger quantities.

Prices quoted are postpaid.

EIFFEL TOWER COS. (73 days.) Tall, vigorous and makes large heads with correspondingly heavy yields. It can be left in the fields for 10 days to two weeks after it is ready to harvest without danger of bolting to seed. It forms firm, solid heads with the leaves well capped over the top, and is very uniform.


Pkt. 10c; oz. 25c; 1/4 lb. 75c; 1 lb. \$2.50 5 to 25 lbs. \$2.40 per lb.

PARRIS ISLAND. (68 days.) Resistant to lettuce mosaic, and highly resistant to premature seeding. Similar to Dark Green Cos, but a shade lighter green, with the leaves of somewhat heavier texture. Of value wherever mosaic is a limiting factor. Uniform and attractive.

Pkt. 10c; oz. 25c; 1/4 lb. 85c; 1 lb. \$2.75 5 to 25 lbs. \$2.65 per lb.



Bibb



Great Lakes 118

GREAT LAKES 118. (83 days.) This strain has been developed for uniformity of size and type, and is the very best strain for Florida. The plants are large, vigorous and have a distinctive, deep grassy-green, attractive appearance. The heads are relatively large, compact, firm and solid, with abundant wrapper leaves which cover the heads, protecting them from cold and heat injury. Great Lakes 118 will stand high temperatures, is much less subject to bottom rot, tip burn and slime rot, and does not bolt so readily as other varieties of this type. It offers a possibility of prolonging the Florida lettuce growing season in late spring, and in early fall.

Pkt. 10c; oz. 40c; 1/4 lb. \$1.15; 1 lb. \$4.00 5 to 25 lbs. \$3.90 per lb.

WHITE BOSTON (Ocoee Strain). (75 days.) This strain has been selected under Florida conditions for the development of a large framed lettuce similar to Big Boston. Produces large heads and heavy yields.

Pkt. 10c; oz. 25c; 1/4 lb. 65c; 1 lb. \$2.25 5 to 25 lbs. \$2.15 per lb.

BIG BOSTON. (75 days.) The heads are of cabbage or butterhead type, large sized. The outer leaves are broad, almost smooth, but wavy at the edge, medium light green with a tinge of brown on outer leaf margins.

Pkt. 10c; oz. 25c; 1/4 lb. 65c; 1 lb. \$2.25 5 to 25 lbs. \$2.15 per lb.

MIGNONETTE. (67 days.) Of excellent table quality, used mostly for home gardens. Plants small, compact. Heads small, round, firm, with well blanched heart. Leaves crumpled, reddish brown with dark green tinge. Bolts quickly.

Pkt. 10c; oz. 35c; 1/4 lb. \$1.00; 1 lb. \$3.50 5 to 25 lbs. \$2.15 per lb.

BIBB. (54 days.) A distinctive variety of high table quality, used for commercial production, and also suitable for home garden. Plants form a small rosette head; very early maturing and rather quick to bolt. Leaves thick, tender, quite smooth; deep green outside, blanched to creamy yellow within.

Pkt. 10c; oz. 25c; 1/4 lb. 75c; 1 lb. \$2.50 5 to 25 lbs. \$2.40 per lb.

SIMPSON, BLACK-SEEDED. (46 days.) The best large, very early, nonheading or leaf variety. The leaves are broad, much blistered and crumpled, and excessively frilled on the borders, light yellowishgreen in color, very tender, of fine delicate texture, and of excellent quality.

Pkt. 10c; oz. 20c; 1/4 lb. 60c; 1 lb. \$2.00 5 to 25 lbs. \$1.90 per lb.

MUSTARD

Plant 4 pounds of seed per acre.

In the garden make two plantings at different times.

1 packet to 40 feet of row at each planting.


Culture. General cultural directions similar to those for cabbage (see page 11), but is always planted direct in the field or garden. Mustard is one of the quickest growing greens. Sow seed August to March in good rich garden soil in rows 24 inches apart and thin from 8 to 10 inches in the row. Heavy fertilization with a fertilizer mixture high in nitrogen or ammonia, and moist soil are conducive for quick growth, high yield and good edible quality. Does best when grown in cool weather of late fall, winter and early spring.

The number of days after each variety indicates the

time from seed to harvest.

Prices quoted are postpaid.

Write for special prices on larger quantities.




Fordhook Fancy

FORDHOOK FANCY. (42 days.) A vigorous, upright grower, producing large, bright green, very deeply fringed leaves, attractive in appearance. A very heavy yielder. The plants stand relatively long without bolting to seed. Of good edible quality, crisp and tender in texture, and mild in flavor. Our stock of this variety is uniformly true to type.

Pkt. 10c; oz. 15c; 1/4 lb. 35c; 1 lb. \$1.00 5 to 25 lbs. 90c per lb.

TENDERGREEN (MUSTARD SPINACH). (35 days.) Although not a true mustard, it is like a quick growing mild mustard, with a spinach flavor, and remains tender even in dry, hot weather. An all-season plant, it withstands heat and may be grown throughout the spring and summer. The smooth, oblong leaves are dark green in color, with small white midribs. Relatively slow in bolt to seed.

Pkt. 10c; oz. 15c; ½ lb. 35c; 1 lb. \$1.00 5 to 25 lbs. 90c per lb.



Florida Broad Leaf

FLORIDA BROAD LEAF. (40 days.) Plants tall and upright, with large, thick, smooth, round to oval, medium green leaves, 8 to 10 inches long and 8 inches wide. The leaves have a large, distinct, flattened, whitish stem and midrib. It produces more leaf growth than other varieties. Plants stand a long time without bolting to seed, and do well in hot weather. Of excellent edible quality, the flavor being very mild. Fine for greens and salads.

Pkt. 10c; oz. 15c; 1/4 lb. 35c; 1 lb. \$1.00 5 to 25 lbs. 90c er lb.

CHINESE BROAD LEAF. (40 days.) A hardy and vigorous grower, producing large, broad, elongated or oblong leaves (9 to 10 inches long and 6 inches wide) which are slightly crumpled and of medium dark green color. The leaves have a small, rounded midrib and differ in this respect as well as in shape of leaf from the Florida Broad Leaf. Tender and of excellent quality, with mildly pungent flavor, when boiled for greens.

Pkt. 10c; oz. 15c; 1/4 lb. 35c; 1 lb. \$1.00 5 to 25 lbs. 90c per lb.



Chinese Broad Leaf

OKRA

Plant 8 pounds per acre.

In the garden plant 1 ounce to 75 feet of row.

Okra does well all over Florida where the young seed pods are used in different ways and are highly esteemed. In addition to being served alone, pods are used in soups and stews to which they give body. It is a healthful vegetable, and should be planted in every Florida garden. We have done a great deal of work on our seed stocks of this valuable Florida vegetable.

Each year, we make special selections for stock seed from only the best, earliest maturing plants. Our seed growing fields are carefully rogued so as to eliminate any slightly off-type plants. We are confident we have the best strains of okra seed obtainable anywhere.


Culture. Okra can be grown in Florida any time from February until September, but it is a warm-weather plant and will not do much until the ground becomes warm. For early planting when the ground is cold always treat okra with Spergon or Arasan. (See pages 74 and 75.) Plant in rows 3 to 4 feet apart and sow thinly in the drill. Okra should be cut every second day. If this is not done some of the pods become too old for market and the young plants will stop bearing. Plants should bear for several months after harvesting starts. Okra does best on sandy loam soil with a fair amount of fertility. On sandy loam soils, 500 to 700 lbs. per acre of fertilizer at time of seeding, and two or three side dressings of nitrate of potash during the bearing period are desirable. Okra is very susceptible to nematodes causing root knot disease, and therefore it is best to plant on land not previously used for growing crops that are subject to attack by nematodes. For the control of root knot nematodes use (See page 76.) For insect control see page 67.

Prices quoted are postpaid.

Write for special prices on larger quantities.



Clemson Spineless



Perkins' Spineless

PERKINS' SPINELESS. (53 days.) Pods similar in size, shape and appearance to Perkins' Long Green, except they are practically spineless. Early, productive, semidwarf plant 4½ to 5 feet tall. Selected for uniformly higher yields of dark green, long slender pods desired in the markets. This Perkins' Spineless should not be confused with Clemson Spineless, being an entirely different and distinct variety, the pods having the long, slender shape and dark green appearance of Perkins' Long Green. Perkins' Spineless can be sold in any market for Perkins' Long Green.

Pkt. 10c; oz. 15c; 1/4 lb. 35c; 1 lb. \$1.00 5 to 25 lbs. 90c per lb.

CLEMSON SPINELESS. (55 days.) Plants are semi-dwarf, growing to a height of 4½ feet on fairly good soil and produce uniform, straight, large, thick, ridged pods of the Perkins' Long Green type, of good length, averaging 3 to 3½ inches when ready for market, green in color, and entirely free from spines, thus making it easy to pick without discomfort. Pods are thicker than Perkins' and fill the crates quickly. Plants are close jointed, and produce an enormous yield. Our strain of this variety is highly bred, uniform, and one of the best, the fields showing practically no variation in plant type. Very well adapted for home gardens and local markets, but does not stand long distance shipments as well as Perkins'. In greatest demand on southern markets.

Pkt. 10c; oz. 15c; ½ lb. 35c; 1 lb. \$1.00 5 to 25 lbs. 90c per lb.

ONION


For dry bulbs sow 1½ lbs. in seedbed to plant an acre, or plant 3 lbs. per acre in the field. For the garden sow 1 packet in seedbed for 350 to 450 plants to set 80 feet of row or plant 1 packet to 80 feet of row.

Culture. For large mature bulbs in Florida, first plantings are made the end of August and continued until November. September and October are the best months to plant. The seed is generally planted in the open field, where the onions are to grow, in rows 18 inches apart, and when 4 to 5 inches high they are thinned out from 3 to 4 inches apart, depending on variety and size of bulbs desired. Those taken out can be used to transplant any possible skips. Onion seed can also be planted in seedbeds and transplanted to the field later. This is a safe and sure method, and is becoming more popular in Florida. Transplant when plants are small. The larger the plants when transplanting, the more splits will develop. It is also important to avoid checking larger the plants when transplanting, the more splits Do not work soil to onions after they start bulbing. Onions do best in sandy loam or muck soil, and require at least a ton of fertilizer mixture per acre for sandy soils or 1,000 to 1,200 on muck.

The number of days after each variety indicates the time from planting the seed to large mature bulbs under average Florida conditions.

Prices quoted are postpaid.

Write for special prices on larger quantities.



Texas Grano

Granex Hybrid


Excel.

EXCEL (BERMUDA 986). (100 days.) Produces more onions of high marketable quality ten to fourteen days earlier than regular Yellow Bermuda. It is practically free of bolters, doubles, splits, and color defects. Uniform in maturity, size, color and shape of bulbs, the shape being a deep or thick-flat, the color a light yellow. Crisp and mild in flavor, and keeps better in storage than other Yellow Bermudas.

Pkt. 10c; $\frac{1}{2}$ oz. 20c; oz. 30c; $\frac{1}{4}$ lb. 85c; 1 lb. \$3.00 5 to 25 lbs. \$2.90 per lb.

GRANEX. (120 days.) The first short-day F¹ hybrid onion adapted to the South, filling the requirements for an onion suitable for growing in Florida. Very early, extremely uniform in type, color and maturity. Practically no splits or bolting to seed. In many trials, it has out-yielded both its Grano type and Excel type parents. Yields of 600 to 900—50-lb. bags of U.S. No. 1 onions per acre have been reported. The bulb of Granex is short top-shaped or slightly flat, mild in flavor, yet keeps unusually well. On the basis of earliness, total yield, percentage of marketable bulbs, attractive appearance and general all-around good quality, Granex is strongly recommended.

Pkt. 10c; ½ oz. 40c; oz. 75c; ¼ lb. \$2.25; 1 lb. \$8.00 5 to 25 lbs. \$7.90 per lb.



Texas Grano 502

TEXAS GRANO 502. (130 days.) This strain is fully ten days earlier, more uniform in shape, and slightly smaller in size than regular Grano. Bulbs have light straw-colored skin, with pure white flesh, are globeor top-shaped, slightly tapering to the root. Mild in flavor with a sweet taste.

Pkt. 10c; ½ oz. 25c; 1 oz. 40c; ¼ lb. \$1.25; 1 lb. \$4.50 5 to 25 lbs. \$4.40 per lb.

CRYSTAL WAX BERMUDA. (110 days.) Early maturing, flat in shape, of waxy white color, attractive in appearance, and of mild flavor.

Pki. 10c; ½ oz. 20c; oz. 30c; ¼ lb. 85c; 1 lb. \$3.00 5 to 25 lbs. \$2.90 per lb.

EVERGREEN (Nebuka). (60 days to green onions.) Makes a long stem, much longer than ordinary bunching onions, pure white in color, and without a bulb. Resistant to pinkroot, thrips and smut.

Pkt. 10c; $\frac{1}{2}$ oz. 20c; oz. 35c; $\frac{1}{4}$ lb. \$1.00; 1 lb. \$3.50 5 to 25 lbs. \$3.40 per lb.

ONION SETS

Plant 8 to 12 bushels of sets per acre. In the garden 1 pint of sets to 40 feet of row.

A bushel weighs about 32 lbs.

Set out in rows 1½ feet apart and 1 to 2 inches apart in rows of green onions. Plant from September to March.



Sets for green onions or early maturing bulbs.

Prices quoted are f.o.b. Plant City except pints and quarts which are postpaid. 1 bushel weighs about 32 lbs.

White SetsYellow Sets	Pint 35	Quart .60		Bushel 4.50 4.50
Available from S	Septembe	r to Marci	h	




PARSLEY

Plant 1 packet to 150 feet of row, or 2 pounds per acre.

In the garden plant 1 packet to 30 feet of row.

Culture. Parsley can be planted in the fall or spring, in drills 18 inches aart, and thinned to 6 inches in the row. Seed is slow to germinate and it usually takes from two to three weeks to produce a satisfactory stand.



Paramount (Improved Moss Curled)

PARAMOUNT (Improved Moss Curled.) (90 days from seed plantings to first harvest.) Plant very vigorous in growth, usually 12 inches tall and spreading 20 inches when given sufficient room, with very long, stout stems for bunching. This variety has been bred and selected for uniform, attractive leaves of very dark green color.

Pkt. 10c; oz. 20c; 1/4 lb. 60c; 1 lb. \$2.00 5 to 25 lbs. \$1.90 per lb.

PARSNIP

Plant 1/2 ounce to 150 feet of row.

In the garden plant 1 packet to 50 feet of row.

Culture. Parsnips do best in deep, rich, sandy soil. Sow seed in Florida from September to December, in rows 1½ to 2 feet apart; cover with one-half inch of soil. When plants are well up, thin to 4 inches apart in row. Parsnip seed germinates very slowly. The Parsnip is a winter vegetable with more starch and sugar than other root crops. To prepare, cook roots one hour in water, scrape off the skin. Slice or chop coarsely or mash, and season with milk, pepper and salt, or slices may be fried after seasoning. A delicious and nutritious dish.

HOLLOW CROWN. (120 days from seeding to harvest.) Roots long, 10"-12", 2½"-3" at shoulder; hollow crowned; tapering gradually. Flesh white, with small core.

Pkt. 10c; oz. 20c; ¼ lb. 50c; 1 lb. \$1.75 5 to 25 lbs. \$1.65 per lb.

PEAS

Plant 1 bushel (56 lbs.) per acre in single rows; 1½ bushels (84 lbs.) per acre in double rows.

In the garden make three plantings at different times.

Plant 1/2 pound to 100 feet of row at each planting.

Culiure. Peas are very hardy and are not commonly attacked by insects. Peas can be grown on almost any kind of soil and with about the same cultivation as beans. Planting in double rows, six to eight inches apart, is a good plan, because this enables the plants to support each other. Single rows should be spaced two and a half to three feet apart, and double rows three to three and a half feet apart from center of double rows. Plant seed 1 to 2 inches apart in the row, and 1½ to 2 inches deep. By working the soil up against the stems of the plants they will bear longer. Use a high grade fertilizer, at the rate of 1,000 to 1,200 pounds per acre. To control soil diseases, and to get better stands, more vigorous growth and increased yields, seed should be treated (see pages 74 and 75). Being a legume, the seed should also be inoculated with Nitragin. (See page 82).

The number of days after each variety indicates the relative time from planting to first picking.


Write for special prices on larger quantities.

HUNDREDFOLD. (53 days.) This variety is one of the finest large early peas for Florida sandy loam soils, and in cool seasons is also adapted to Florida muck soils. It is a sure cropper and a better yielder than other early large podded varieties. The plants are vigorous in growth, eighteen to twenty inches in height, with heavy, very dark green foliage, producing a great abundance of large (4 to 4½ inches long) broad, deep green, pointed pods containing usually eight large peas of fine quality.

½ 1b. 20c; 1 lb. .65; 2 lbs. \$1.25 Postpaid
 Not prepaid, 10 lbs. \$2.90; 50 lbs. \$13.00

LITTLE MARVEL. (52 days.) The most popular pea for Florida, because of its good quality, earliness and high yield of well filled pods. Especially adapted to Florida muck soils where it is a sure cropper with well filled pods, even in warm seasons. The vines are vigorous, stock, dark green in color, growing about 1½ feet tall, and loaded with straight, round, blunt, plump pods about 3 inches long, well filled with 7 to 8 tender peas.

½ lb. 20c; 1 lb. 65c; 2 lbs. \$1.25 Postpaid
 Not prepaid, 10 lbs. \$2.80; 50 lbs. \$12.50



Little Marvel



PEPPER

Sow 1/2 to 1 pound of seed in seedbed to plant an acre. Sow 2 to 3 lbs. per acre when drilled in field. For the garden sow 1 packet in a protected seedbed for 24 plants to set 40 feet of row.

Culture. For a crop in central Florida sow the seed in June, July and August; for a long season winter crop on the east coast plant in August, September and October. For an early spring crop in Florida west coast sections sow seed in November and in north Florida in December or January.

High temperature is needed to germinate pepper seed satisfactorily, and it is difficult to get a good stand when the ground is wet and cold. Hammock land, or rather moist soil, is best suited to this crop, although any medium good sandy loam soil, properly fertilized, will grow good peppers. Make rows three feet apart and set plants 10 to 16 inches in the row. As this crop is in bearing for a long time, it is most profitable to make several applications of fertilizer, using 400 pounds per acre about every three weeks until at least a ton has been used.

For the control of insects and diseases see page 68. The number of days in parentheses after each variety indicates the relative time from setting out plants to picking of marketable green peppers. It requires six to eight weeks to produce plants ready for setting.

Prices quoted are postpaid.
Write for special prices on larger quantities.

WONDER GIANT. (74 days.) This variety of the thick meat Florida Giant type is popular, especially on the west coast of Florida where it has been grown the past few seasons. The vigorous growing plant is more spreading and does not grow as tall as Florida Giant. Developed and selected for uniformity in plant and fruit type and productivity. The blocky fruits borne upright on the plant are mostly four-lobed, with sweet, thick flesh, of deep glossy green color, making an attractive pack.

Pkt. 10c; oz. 65c; ¼ lb. \$2.15; 1 lb. \$7.50 5 to 25 lbs. \$7.40 per lb.

YOLO WONDER A. (70 days.) A superior Asgrow strain, replacing the original Yolo Wonder. Yolo Wonder A is a tobacco-mosaic-resistant variety, resembling California Wonder, of general use for shipping, local market, home garden, and suitable for processing where that type is desired. Plants upright, vigorous, with a good foliage cover, but compact, permitting close planting; productive, with heavy crown set. Fruits pendent, dark green, more uniformly blocky than original with considerably higher percentage of four-lobed fruits; walls thick.

Pkt. 10c; oz. 75c; 1/4 lb. \$2.25; 1 lb. \$8.00 5 to 25 lbs. \$7.90 per lb.



Yolo Wonder A



Florida Giant

by us, produces an abundant yield of large, heavy, thick walled, fleshy, dark green fruits, over a long period, but requires heavy and frequent fertilization, and a rich, moist soil. The tall plants are compact but very vigorous and upright in growth. Slightly later than World Beater, they are heavy producers, often bearing six to eight large fruits at one time. The fruits are longer than most thick-meated peppers, about 4 to 4½ inches long, and 3 to 3½ inches in diameter, usually with four lobes, are smooth and well filled at the blossom end. Makes a very attractive pack and carries well to distant markets.


Pkt. 10c; oz 60c; 1/4 lb. \$1.90; l lb. \$7.00 5 to 25 lbs. \$6.90 per lb.



California Wonder

CALIFORNIA WONDER. (75 days.) An outstanding medium-late variety. Plant is vigorous, upright, prolific. The thick meated, smooth, large, heavy fruits average about four inches in length and about the same in diameter. This is an excellent shipping type, holding up well in transit and on the market, and is in great demand by truckers. Solid thick walls and very sweet and mild. Blocky in shape. Color an attractive, glossy dark green. Requires frequent fertilizing and continuous moisture to hold the size of the fruit up over a long harvesting period.

Pki. 10c; oz. 60c; 1/4 lb. \$1.75; 1 lb. \$6.50 5 to 25 lbs. \$6.40 per lb.



Early Calwonder

EARLY CALWONDER. (65 days.) An early maturing strain of California Wonder, maturing 7 to 10 days ahead of other strains, and producing fruits over a long season. The plant is not as tall and erect in growth as regular California Wonder, but is very vigorous and prolific. The fruits usually are four-lobed and attractive in appearance, being smooth, dark green, uniform and thick meated. This early strain is rapidly growing in popularity.

Pkt. 10c; oz. 75c; ½ lb. \$2.25; 1 lb. \$8.00 5 to 25 lbs. \$7.90 per lb.

WORLD BEATER (Thick Wall Strain). (70 days.) This strain of World Beater Pepper was developed in our breeding station at Bradenton, Florida, for a more uniform, thicker walled, heavy yielding strain. In numerous trial plantings in different sections of Florida we have found this to be a heavy yielder of uniform, true to type, smoother, thicker walled fruits, which are larger in size and carry better to distant markets than regular World Beater which usually are very thin-walled. In some sections it has brought a premium on the market. The plants are erect in growth and hold the large heavy fruits well off the ground.

Pkt. 10c; oz. 60c; 1/4 lb. \$1.90; 1 lb. \$7.00 \$5 to 25 lbs. \$6.90 per lb.




World Beater (Thick Wall Strain)

ITALIAN "EL". (65 days.) This green-fruited strain of Neapolitan is in increasing demand on northern Italian markets where it is often preferred to the regular yellow-fruited Neapolitan because of its peculiar sharp taste. Fruits are slightly larger and thicker fleshed than the regular yellow Neapolitan. The vigorous growing plants produce a heavy yield of light green fruits over a long season.

Pkt. 10c; oz. 80c; 1/4 lb. \$2.65; l lb. \$9.50 5 to 25 lbs. \$9.40 per lb.

NEAPOLITAN. (50 days.) This is an early, medium large, prolific variety, growing in popularity on some northern markets where it is in demand among Italian people. This plant is wonderfully productive, carrying as many as forty fruits on a single plant. The fruits are upright in growth, averaging 4 inches long and 2 inches in diameter, three-lobed, yellowish green, becoming bright red when ripe. The flesh is mild and sweet.

Pkt. 10c; oz. 75c; 1/4 lb. \$2.35; l lb. \$8.50 5 to 25 lbs. \$8.40 per lb.



Cuban

CUBAN. (62 days.) A distinctly longer fruited variety of Neapolitan type, particularly desirable for those areas where peppers are widely used for frying. Plants medium large, somewhat spreading, very productive. Fruits pendent, greenish yellow, tapering; two and three-lobed, medium thick walls averaging 6 inches in length and 2 inches in diameter at stem end.

Pkt. 10c; oz. 80c; ½ lb. \$2.60; 1 lb. \$9.50 5 to 25 lbs. \$9.40 per lb.

ANAHEIM. (80 days.) A popular, rather long mildly pungent variety, not as hot as Long Thick Cayenne. The fruits are meaty, 6 to 7 inches in length and 1 to $1\frac{1}{2}$ inches in diameter at stem end, green ripening to a bright red. Often dried for winter use.

Pkt. 10c; oz. 60c; 1/4 lb. \$1.75; 1 lb. \$6.25 5 to 25 lbs. \$6.15 per lb.

HUNGARIAN YELLOW WAX. (60 days.) A very early, hot sort, valuable for canners. Plants are rather dwarf in growth, and exceedingly prolific. Fruits pendent, medium slender, long, tapering, smooth and very hot or pungent. Color waxy yellow, changing to bright crimson at maturity.

Pkt. 10c; oz. 75c; 1/4 lb. \$2.35; l lb. \$8.50 5 to 25 lbs. \$8.40 per lb.

LONG THICK CAYENNE (Hot). (70 days.) This favorite hot variety produces large and productive plants. Fruit 5 to 6 inches long, 1 inch thick, slender, tapering, pendent, very hot or pungent, deep green in color, ripening to a brilliant red.

Pkt. 10c; oz. 65; 1/4 lb. \$2.15; 1 lb. \$7.75 5 to 25 lbs. \$7.65 per lb.

POTATOES


Plant 600 to 900 pounds (10 to 15 bu.) per acre in 3-ft. rows.

In the garden plant 12 pounds to 200 feet of row.

Culture. Plant in rows 36 to 42 inches apart, dropping a seed piece every nine to ten inches in the row. The seed piece should be at least one to one and one-half ounces in size. It is best to cut seed within one day of planting. Fertilizer should be put in seven or eight days before planting. Use from 1,500 to 2,000 pounds per acre, depending on type of soil. A 6-8-8 fertilizer mixture is a good standard for potatoes. On rich Everglades muck from 200 to 500 pounds per acre of 0-10-12 or 0-8-16 mixture applied in one application at planting time is a common practice. To control blight and increase the yield, spray with Nabam using 2 quarts Nabam, 34 lb. zinc sulphate, to 100 gallons water. For control of insects and diseases see page 68.

RED PONTIAC (Dakota Chief) (State Certified). (80 days.) Packed in 100-lb. bags. This variety is the result of a cross, Red Bliss and Katahdin, made by the Michigan Experiment Station. In tests at the Sub-tropical Experiment Station in Homestead and elsewhere in Florida, it has greatly outyielded Red Bliss. The tubers of Red Pontiac are slightly longer, and have shallower eyes, but the color is a bright red like Red Bliss. Pontiac tubes are tougher, stand shipment better and are better keepers than Bliss.

Write Plant City or your nearest Kilgore store for price on seed potatoes stating quantity wanted.



Red Pontiac

PUMPKIN

Plant two pounds seed per acre. In the garden plant $\frac{1}{2}$ ounce to 50 feet.

Culture. Plant 4 to 5 seeds in hills 8 to 10 feet apart one way and 5 feet apart the other way, when the weather gets warm. Can be planted in corn. Thin to two or three plants after well started. Give shallow and frequent cultivation until vines begin to run. Cut pumpkins when fully ripe.

Prices quoted are postpaid.

Write for special prices on larger quantities.

king of the mammoths. (100 days.) Commonly known as a pumpkin, but is really the very large fruited Mammoth Chili squash, known as the giant among squashes. Fruits are globular, slightly ribbed, light yellow color mottled with orange. Flesh is solid, light yellow-orange, coarse, of fair quality. Grown mainly for stock feed. Fruits measures 20 to 22 inches across and weigh 60 to 80 pounds.

Pkt. 10c; oz. 20c; 1/4 lb. 50c; 1 lb. \$1.75 5 to 25 lbs. \$1.65 per lb.



Small Sugar

SMALL SUGAR. (110 days.) The best variety for general use, and particularly desirable for pies. Fruits are round, flattened at ends. Skin is hard, smooth, somewhat ribbed, or deep orange color. Flesh is thick, sweet and dry, of bright orange color, and very high quality. Makes delicious pumpkin pies. Fruits measure eight to ten inches and weigh six to eight pounds.

Pkt. 10c; oz. 20c; ½ lb. 50c; l lb. \$1.75 5 to 25 lbs. \$1.65 per lb.

CONNECTICUT FIELD (Cornfield). (118 days.) Prolific and vigorous, usually planted in cornfields. Extensively grown for making pies, for canning and for stock feed. Fruits are large round to slightly flattened, with a hard, smooth surface, slightly ribbed, deep yellow or orange color. Fruits measure 12 to 14 inches and weigh 15 to 20 pounds.

Pkt. 10c; oz. 20c; 1/4 lb. 50c; 1 lb. \$1.75 5 to 25 lbs. \$1.65 per lb.

ASGROW VIGORPAK SEED

Seed of superior vigor • Scientifically pre-conditioned • Packed in airtight cans

For many years the ASGROW RESEARCH DEPARTMENT has studied the factors affecting the life of vegetable seed. It was found that, within limits, the drier the seed the longer it will keep its vitality. The higher the moisture content of the seed and of the surrounding atmosphere, the more rapidly will the seed lose its capacity to germinate and grow into healthy plants. From these studies ASGROW has perfected a way to process and package seeds so that they will keep their initial vitality for years under almost any storage conditions, short of killing temperatures.

will keep their initial vitality for years under almost any storage conditions, short of killing temperatures.

The practical application of these investigations is in ASGROW VIGORPAK SEED. The internal moisture of the seed is brought to a safe level for that particular kind of seed, as determined by the ASGROW RESEARCH DEPART-MENT. Then, after treatment with an adequate seed protectant, the seeds are packed in moisture-proof cans and hermetically sealed. So long as it is in the unopened can, ASGROW VIGORPAK SEED is protected not only from the adverse effects of high humidity but also from damage by insects and vermin and from loss by spilling or mixing. With these hazards eliminated, seed can safely be carried over to the following season, provided the can has not been opened.

YOUR INSURANCE AGAINST
Loss in germination and vigor • Loss from pests • Waste from mixture

RADISH

Plant 6 to 8 pounds of seed per acre. In the garden make four plantings at different times. 1 packet to 25 feet of row at each planting.

Culture. Plant in rows 12 to 18 inches apart and thin to 1 to 2 inches in the row. Successive plantings should be made every few weeks for a continuous supply. Our radish seed stocks are produced each year from carefully selected transplanted roots. For insect and disease control see page 68.

Prices quoted are postpaid. Write for special prices on larger quantities.

BUCCANEER. (21 days.) Developed for uniformity and attractive market appearance. The root is globeshaped, with a lustrous scarlet exterior, and with short top growth.

> Pki. 10c; oz. 15c; 1/4 lb. 35c; 1 lb. \$1.00 5 to 25 lbs. 90c per lb.

CAVALIER. (23 days.) An attractive, short-topped Scarlet Globe type, with olive-shaped roots. Used both for bunching and for pre-packaging. Tops very short; roots bright scarlet, very firm.

Pkt. 10c; oz. 15c; 1/4 lb. 35c; 1 lb. \$1.00 5 to 25 lbs. 90c per lb.

CHERRY BELLE. (23 days.) A widely used variety for market, suitable for forcing and equally desirable for summer crops; both for bunching and for pre-packaging. Tops very short, even in warm weather. Roots bright cherry red, round, firm, free from pithiness.

Pkt. 10c; oz. 15c; 1/4 lb. 35c; 1 lb. \$1.00 5 to 25 lbs. 90c per lb.

GLOBEMASTER (Early Scarlet Globe, Special Forcing). (23 days). A scarlet globe type for late summer and fall harvest. Tops medium short. Roots globe-shaped, of crisp texture, attractive, with lustrous scarlet exterior.

> Pkt. 10c; oz. 15c; 1/4 lb. 35c; 1 lb. \$1.00 5 to 25 lbs. 90c per lb.

WHITE ICICLE. (23 days.) The straight roots are from 5 to 6 inches long, cylindrical, and very smooth and waxy white throughout.

> Pkt. 10c; oz. 15c; 1/4 lb. 35c; 1 lb. \$1.00 5 to 25 lbs. 90c per lb.




Cavalier

SPINACH

Virginia Savoy—Plant 12 to 15 pounds per acre in rows, or 20 to 25 pounds per acre broadcast. In the garden make two plantings at different times. Plant one packet to 50 feet of row at each planting.

> Prices quoted are postpaid. Write for special prices on larger quantities.

Culture. Successful for fall and winter planting. Does not grow well in hot weather. Requires a moist rich soil. Plant seed in rows 16 to 20 inches apart, and thin to 6 inches apart in the row.



DIXIE MARKET. (37 days.) New. A savoy-leafed shipping variety, resistant to both downy mildew (blue mold) and cucumber mosaic (blight) for winter production in the South. Plants upright, compact. Leaves dark green, rounded, thick, and well savoyed; more productive than standard Bloomsdale types.

Pkt. 10c; oz. 15c; 1/4 lb. 35c; 1 lb. \$1.00 5 to 25 lbs. 90c per lb.

EARLY HYBRID 7. (37 days.) The first hybrid spinach to be introduced, the result of a breeding program conducted by the U.S.D.A. with a view to incorporating disease resistance with the increased yield expected from F: hybrids.

Fast growing and heavy yielding, intended primarily for fall and winter harvest in the South and West. Bolts rapidly, and therefore not well suited for late spring and summer use. Plants large, vigorous, somewhat spreading, resistant to blue mold (downy mildew) and to blight (cucumber mosaic). Leaves large, somewhat savoyed, medium dark green.

> Pkt. 10c; oz. 20c; 1/4 lb. 55c; 1 lb. \$1.50 5 to 25 lbs. \$1.40 per lb.

VIRGINIA SAVOY. (40 days.) Better for market than the smooth leaved varieties, because it does not pack as closely and thus is not subject to heating. Developed for resistance to blight or mosaic disease. The plant is upright and compact in growth, with very thick, curly or savoyed dark green leaves.

Pkt. 10c; oz. 15c; 1/4 lb. 30c; 1 lb. 80c 5 to 25 lbs. 70c per lb.

SAGE

Culture. Plant in fall or spring. Sage does well on sandy soil. Plant seed in rows 15 to 18 inches apart. Thin to six inches in the row. The plant is a hardy perennial, and of easy culture in Florida.

Leaves should be cut as plant blooms, and should be

dried quickly in the shade. Grows 18 inches tall.

Pkt. 10c; 1/4 oz. 35c; 1 oz. \$1.00

SQUASH

Culture. This is one of the quickest and easiest crops that can be grown, and usually proves very profitable. The earliest varieties begin bearing in six or seven weeks from seed. Squash can be grown on almost any kind of soil. Use from 800 to 1,000 pounds of fertilizer per acre.


For bush varieties, drill seed in rows at least 3 to 4 feet apart in the row, thinning to 1 foot when 3 inches high. Bush squash in Florida is a good paying crop for fall or spring. Plant for spring, during January, February and March; for fall, from August until October. Mosaic has become a most serious disease of squash in many sections of Florida. See Sanitation on page 60. Insect and Disease control on page 66.

The number of days after each variety represents the average time required from seed planting to edible or marketable fruits.

For protection from rats and birds use Kilgore's Melon Seed Treatment. (See page 76.)

All prices quoted are postpaid.

Write for special prices on larger quantities.



Long Cocozelle

BUSH VARIETIES

Plant 2 to 3 pounds of seed per acre. In the garden one packet to 50 feet.


LONG COCOZELLE. (50 days.) This is a true Italian Vegetable Marrow. When ready for market, the fruit is 10 to 12 inches long, slender, 1 to $1\frac{1}{2}$ inches in diameter, cylindrical in shape, straight, smooth, dark green with lighter green stripes.

Our stock of this variety produces a bushy, open type of plant, with deeply cleft leaves, and a heavy yield of uniformly true-to-type fruits. We have devoted many years to the development of this superior strain, and we can say with full confidence that our Cocozelle is the earliest, most prolific, and uniformly true-to-type strain you can secure anywhere.

Pkt. 10c; oz. 20c; 1/4 lb. 50c; 1 lb. \$1.75 5 to 25 lbs. \$1.65 per lb.

EARLY PROLIFIC STRAIGHTNECK. (42 days.) This variety is ten days earlier and more productive than other strains of Straightneck Squash. The plants are vigorous, and the fruits are of brighter lemon-yellow color. Valuable for both home and market use. Because of its uniformity, earliness, high yield and attractive appearance this variety has grown greatly in popularity, especially in northern markets. Makes a very attractive pack.

Pki. 10c; oz. 15c; 1/4 lb. 45c; 1 lb. \$1.50 5 to 25 lbs. \$1.40 per lb.




Early Summer Crookneck

EARLY YELLOW SUMMER CROOKNECK. (45 days.) A very early and prolific, light lemon-yellow color strain with an open type of plant growth. The most popular and most extensively grown of any variety of squash in Florida. Fruits are small with curved neck, and uniform. Its rich, bright, lemon-yellow color gives it a very good market appearance, and it is in demand on all southern markets. Our strain of this variety is very early and exceedingly prolific, and has been selected for uniformly shaped fruits with a pronounced crook. No better strain of this most popular variety can be secured anywhere for Florida planting.

Pkt. 10c; oz. 15c; ½ lb. 45c; 1 lb. \$1.50 5 to 25 lbs. \$1.40 per lb.

BLACK ZUCCHINI. (45 days.) The plant of this improved selection, instead of developing a large, vigorous bush like the regular Black Zucchini, forms a short, single semi-runner, and produces a heavy yield of fruit over a long period. The plant is an open type of growth, with deeply cleft leaves, allowing the air and sunlight to circulate and prevent fruit rot. The fruits are dark green in color, six to eight inches long at market stage. A very early, prolific, improved strain of Black Zucchini.


Pki. 10c; oz. 20c; 1/4 lb. 60c; 1 lb. \$2.00 5 to 25 lbs. \$1.90 per lb.



Early Prolific Straightneck



BUSH VARIETIES (Continued)



White Bush Scallop

WHITE BUSH SCALLOP (EARLY WHITE). (48 days.) This is a popular type of squash for shipment to northern markets. The smooth fruits are of pale green color when very young, becoming white as they mature, are round and deep, with ridged or scalloped edges, measuring up to five or six inches across by threee inches thick when ready to market.

Pkt. 10c; oz. 15c; ½ lb. 45c; 1 lb. \$1.50 5 to 25 lbs. \$1.40 per lb.

BLACK BEAUTY. (40 days.) An attractive dark green Zucchini type, newly introduced from the University of Connecticut, maturing somewhat earlier than older varieties. Plants relatively compact, productive. Fruits blocky shaped, not prominently ridged, glossy dark green at prime picking stage, becoming black at full maturity.

Pkt. 10c; oz. 25c; 1/4 lb. 75c; 1 lb. \$2.50 5 to 25 lbs. \$2.40 per lb.

WOOD'S EARLY PROLIFIC. (42 days.) The earliest, meatiest, and most prolific of the patty pan type. It will average seven days earlier than White Bush Scallop, matures its fruit quickly. The plants are prolific, producing round, smooth, deep, thick fruits, slightly scalloped, and of creamy-white color, usually with a light greenish cast. A very good variety for home use or for shipment.

Pkt. 10c; oz. 20c; ½ lb. 50c; 1 lb. \$1.75 5 to 25 lbs. \$1.65 per lb.




Table Queen


RUNNING VARIETIES

Plant two pounds per acre of running varieties.

In the garden one packet to 50 feet.

Culture. Cultural requirements are similar to bush squash (see page 32), but running varieties require much more room, planting in rows 6 to 8 feet apart and hills 3 to 4 feet apart. Plant 3 to 5 seeds in each hill and thin to two plants when three inches high. Generally grown as a spring crop in Florida.

In these types of squash the vines run 10 to 24 feet. The fruits are left on the vines until fully grown with a hard rind, and are then harvested by cutting the stem. Running squash may be stored in a cool, dry place until desired for use, but must be handled carefully to avoid bruising. Unlike bush squash, fruits of the running varieties may be stored and used boiled, baked or for making squash pies.



Butternut 23

BUTTERNUT 23. (80 days.) Vines produce pear-shaped yellowish brown or deep buff-colored fruits. Fruits weigh 2½ to 4 lbs., 8 to 10 inches long, and 3 to 4 inches in diameter at blossom end. The outer shell is smooth and hard, making this squash a wonderful shipper and keeper. The dry flesh is yellow, fine-textured, sweet and of excellent flavor. Also excellent for making pies.

Pkt. 10c; oz. 25c; 1/4 lb. 75c; 1 lb. \$2.50 5 to 25 lbs. \$2.40 per lb.

ROYAL ACORN. (65 days.) To meet an increasing demand in Florida for a larger fruited strain of the Table Queen or Acorn Squash, we are offering this large-fruited strain. It is slightly later than regular Table Queen, not quite so prolific or as uniform in shape and color but the fruits are consistently larger. In other respects this strain is similar to the regular Table Queen.

Pkt. 10c; oz. 20c; ½ lb. 50c; 1 lb. \$1.75 5 to 25 lbs. \$1.65 per lb.

TABLE QUEEN. (60 days.) Extremely early for a running squash. Makes an enormous yield over a long season. Fruits dark green, weighing about 2 lbs., with sides deeply ribbed, and with a point at the blossom end. Yellow-colored flesh is very fine-grained and sweet when baked like Hubbard squash. It is a very good keeper and desirable for home and market use, also for shipping.

Pkt. 10c; oz. 20c; ¹/₄ lb. 50c; 1 lb. \$1.75 5 to 25 lbs. \$1.65 per lb.



TOMATO

Sow $\frac{1}{2}$ pound in seedbed to plant an acre, or 1 pound per acre in field.

For the garden sow one packet in a seedbed for 50 plants to set 150 feet of row.

If plants are staked, sow one packet in seedbed for 50 plants to set 75 feet of row.

Culture. Tomatoes are grown on all kinds of soil, from a light sand to a muck. The land should be put in good condition. Lay off rows 3½ to 5 feet apart, open a furrow, and drill in 1200 pounds per acre of high grade fertilizer such as 4-7-5, and mix thoroughly with the soil. It is good policy to let this stand for seven to ten days and then plant seed or set the plants. Seed may be planted direct in the field or in a seedbed. Plants started in a seedbed should be 6 to 8 inches high when ready to transplant to the field in 4 to 7 foot rows and 2½ to 3 feet apart in a row, or if staked 16 to 20 inches in the row, rows 4 feet apart.


On the east coast of Florida, first plantings are made during August and continue until January. North and central Florida begin planting for a spring crop the middle of February. For a fall crop in Central Florida, plantings are made in July and August.

For control of insects and diseases of Tomatoes see page 70. Spraying or dusting tomato plants with Nabam plus Manganese Sulphate or Maneb before the disease appears will help prevent the destructive late blight disease.

The number of days indicated after each variety represents the time required from setting of plants in the field to produce marketable fruits. It usually takes four or five weeks to produce plants for field setting.

All prices quoted are postpaid.

Write for special prices on larger quantities.



Jefferson (Wilt Resistant)

JEFFERSON (Wilt Resistant). (76 days.) The plant is very vigorous and makes densely-leaved tall vines. It is more open and less subject to bunching than Rutgers. Produces well in the fall but is best suited as a spring tomato on wilt infested soils. The fruits are globe-shaped ripening to a uniform, attractive dark red color, with a good flavor, thick walled with a small core, and relatively free from cracking and catfacing. This variety holds up under adverse conditions better than many other varieties. A good shipper as a green wrap tomato. Jefferson also has become a favorite for pink harvest, has good carrying qualities and will stand rougher handling when shipped as a vine ripe than most other varieties.

Pkt. 10c; ½ oz. 35c; oz. 60c; ¼ lb. \$1.90; 1 lb. \$7.00 5 to 25 lbs. \$6.90 per lb.




Homestead 24 (Wilt Resistant)

HOMESTEAD 24 (Wilt Resistant). (70 days.) A selected strain of Homestead tomato. Our aim in developing this strain has been elimination of off types, leading to uniformity of plants and fruits and a better crop. The plants of Homestead 24 are semi-determinate, open, early in maturing and resistant to Fusarium Wilt. Fruits free from puffiness and cracking, meaty and firm, excellent for green wrap. Homestead 24 is a heavy yielder and will make good crops even on wilt-infested soils. It has done well in the Homestead area, both on heavy marl soil and on the rockland, and seems well adapted to all Florida soils.

Pki. 10c; ½ oz. 40c; oz. 75c; ¼ lb. \$2.35; 1 lb. \$8.50 5 to 25 lbs. \$8.40 per lb.

RUTGERS. (82 days.) Our improved strain is uniform in plant growth and produces uniform smooth, thick wall, deep globe-shaped fruits which ripen to a bright red. The plant is vigorous and a heavy producer. Very firm and free of puffs. Fruits size up well out to the end of the vine. Recommended for a spring crop in Florida and is well suited for light sandy soils and also for shallow, dry pine land on the lower east coast of Florida. Not suitable for deep, wet glade lands where it makes too much vegetable growth.

Pkt. 10c; ½ oz. 25c; oz. 45c; ¼ lb. \$1.40; 1 lb. \$5.00 5 to 25 lbs. \$4.90 per lb.



Rutgers Tomato

WILT RESISTANT GROTHEN'S GLOBE. (70 days.) This variety was developed from a cross of Pan American and regular Grothen's Globe. Wilt Resistant Globe possesses all the good qualities of the regular Grothen's Globe, and in addition it is very resistant to fusarium wilt, although slightly earlier and more vigorous than regular Grothen. In numerous comparative trials between Wilt Resistant and regular Grothen's Globe, even where wilt was not a factor, the Wilt Resistant strain yielded just as high as the regular strain. This fusarium wilt resistant strain is especially recommended for old, wilt infested soils, where the regular Grothen's Globe and other varieties cannot be grown successfully. Is not resistant to all wilts, but is highly resistant to the most serious wilt disease of tomatoes caused by the fusarium soil fungus.


Pkt. 10c; ½ oz. 35c; oz. 60c; ¼ lb. \$1.75; 1 lb. \$6.50 5 to 25 lbs. \$6.40 per lb.

INDIAN RIVER (70 days.) This new variety was developed by the Indian River Field Laboratory at Fort Pierce and the Gulf Coast Experiment Station at Bradenton. It is resistant to Fusarium Wilt, Gray Leaf Spot, and Gray Wall. The plants are indeterminate, with enough foliage to protect the fruits from sunburn, yet open enough to permit thorough spray coverage. They are well adapted for ground culture as well as stake or trellis. The glove-shaped fruits have small blossom-end scars, and are of excellent quality for both "green-wrap" and "pink" shipment. They have good firmness, with thick outer walls and heavy, fleshy inner walls. This variety is adaptable to either fall, winter, or spring production and resists bunching.

Pkt. 10c; ½ oz. 60c; oz. \$1.00; ¼ lb. \$3.00; lb. \$10.50 4 to 25 lbs. \$10.40 per lb.



Indian River



Manalucie

MANALUCIE. (82 days.) This vigorous growing, disease resistant variety was developed by the Gulf Coast Experiment Station at Bradenton, Fla. It possesses multiple resistance, being very resistant to two serious Florida tomato diseases, namely, fusarium wilt, and gray leaf spot. It is also mildly resistant to black spot stem canker caused by the Phoma fungus. Makes a very strong, vigorous growing plant with large leaves and large, strong stems. Fruits are late, but are large size throughout the harvest period. Smooth and relatively resistant to cracking. Firm and fleshy, and good edible quality. Very slow ripening and must be allowed to fully mature before harvesting. Recommended primarily for pink and vine ripened harvest.

Pkt. 15c; ½ oz. 40c; oz. 75c; ¼ lb. \$2.35; 1 lb. \$8.50 5 to 25 lbs. \$8.40 per lb.

KOKOMO. (80 days.) As a result of numerous trials in different sections of Florida during the past few years we are fully convinced that this new, fusarium wilt resistant variety of the Rutgers type is exceedingly well adapted to Florida growing conditions. It will do well wherever Rutgers does well. It is of the general Rutgers type, not as vigorous or bunchy in growth as Rutgers. Vines open up more, making it easier to spray or dust and to harvest. Kokomo sets a heavy crop of good size, smooth, deep globe-shaped, meaty, thick-walled fruit, is a good shipper. It makes a very high percentage of smooth, fancy fruits and it is claimed to set well in hot weather. Is particularly valuable where fusarium wilt is a serious crop hazard, under which conditions Rutgers cannot be successfully grown.

Pkt. 10c; ½ oz. 35c; oz. 60c; ¼ lb. \$1.90; 1 lb. \$7.25 5 to 25 lbs. \$7.15 per lb.

MARGLOBE. (77 days.) This is an excellent strain of the old standard type Marglobe. It produces a vigorous plant resistant to nailhead rust. Fruit deep globeshaped, smooth and thick walls, free of navel and blossom-end scars. Ripens to a beautiful deep scarlet-red color. Well adapted to and recommended for north Florida.

Pkt. 10c; 1/4 oz. 35c; oz. 60c; 1/4 lb. \$1.75; 1 lb. \$6.75 5 to 25 lbs. \$6.65 per lb.

OXHEART. (90 days.) A smooth, thick, tender and almost seedless pink-fruited tomato of delicious flavor. An ideal variety for the home garden and for the local

Pkt. 15c; ½ oz. 60c; oz. \$1.10; ¼ lb. \$3.50; l lb. \$13.00 5 to 25 lbs. \$12.90 per lb.

TURNIP

Plant 2 to 3 pounds of seed per acre.

In the garden make three plantings at different times.

1 packet of turnip seed to 50 feet of row.

Culture. Plant in rows 15 to 20 inches apart. Thin early to one inch, because any over crowding will result in misshapen, slow growing, poor quality roots. Final thinning should be 4 to 6 inches apart. Thinnings may be used for turnip greens. For best quality, turnips should be grown rapidly in rich, moist soil. When turnips are grown for commercial canning and freezing, where only the tops are used and harvested with machinery, plant in double rows $4\frac{1}{2}$ to 5 feet apart, 14 inches apart in the double row. 6 to 8 lbs. of seed is required, no thinning. Turnips can be planted every month in the year.

Prices quoted are postpaid.

Write for special prices on larger quantities.

WHITE EGG. (45 days.) An early, oval or egg-shaped variety with clear white roots of medium size. The flesh is white, firm, fine grained and sweet.

Pki. 10c; oz. 15c; 1/4 lb. 30c; 1 lb. 90c 5 to 25 lbs. 80c per lb.


WHITE GLOBE. (44 days.) A large, round turnip of fine quality. It has small tops, grows rapidly, and gets very large, producing a heavy yield per acre. Resembles Purple Top White Globe except for color.

Pkt. 10c; oz. 15c; 1/4 lb. 30c; 1 lb. 90c 5 to 25 lbs. 80c per lb.

SHOGOIN. (50 days.) An aphid-resistant foliage turnip which resists the hot, dry weather. It produces an immense yield of upright growing foliage, excellent for freezing and canning. The roots are white, semiglobe shaped, but more largely planted for the tops which are large, thick, succulent, tender, and of finest mild flavor.

Pkt. 10c; oz. 15c; 1/4 lb. 30c; 1 lb. 90c 5 to 25 lbs. 80c per lb. **PURPLE TOP WHITE GLOBE.** (50 days.) Ours is a very uniform and well colored strain of this popular variety. Roots globe-shaped, flesh and skin are white, with purple or red at the stem end above ground. Fine grained and sweet, of superior quality. Grown extensively in Florida for shipment to northern markets.

Pkt. 10c; oz. 15c; 1/4 lb. 30c; 1 lb. 90c 5 to 25 lbs. 80c per lb.



Purple Top White Globe

RUTABAGA

Plant 1 pound per acre.

In the garden plant one packet to 100 feet of row.

Culture. Rutabaga is best grown as a late fall and winter crop. Planted in rows 2 feet apart, and the plants thinned to about 5 to 6 inches in the row.

AMERICAN IMPROVED. (90 days.) The smooth roots attain a much larger size than turnips, slightly oblong, pale yellow with purple top, and with a very small neck. Flesh is yellow, solid, tender.

Pki. 10c; oz. 15c; 1/4 lb. 30c; 1 lb. 90c 5 to 25 lbs. 80c per lb.

HIGH VITALITY SEED MUST BE PLANTED TO SECURE A GOOD CROP

The Kilgore Seed Company, has in its central warehouse in Plant City, Florida, the most up-to-date dehumidifying machinery in order to maintain high vitality in all Asgrow seed. Surplus seed supplies in our stores are also kept in specially constructed dry rooms where the humidity is maintained at a uniformly low degree.

Many of our vegetable seeds, such as cabbage, celery, eggplant, onions, pepper, tomato, and watermelon, are now Vigorpacked by Associated Seed Growers. Asgrow "Vigorpak" seed must first have a high germination and vitality standard. The seed is scientifically preconditioned, treated and sealed in air-tight cans. The cans should be opened only as needed for planting. In unopened cans the seed will retain its vigor and germination even under Florida conditions of high humidity and temperature.


In addition to keeping our seeds in artificially dried storerooms, we have a seed testing laboratory where samples of all our seed stocks are tested at regular intervals to further insure our customers against planting low vitality seed. In our seed testing laboratory, tests are made every six weeks of all stocks in our warehouse, mail order department and fourteen Florida stores. We know at all times the correct germination of all seed we have on hand. If any seed stocks are found to be dropping in germination they are immediately removed from our warehouse and stores and are destroyed.

We are taking all these precautions and giving our customers this extra service without any additional cost, in order to provide in Asgrow Seeds the best that money can buy. Play safe and get your seed from The Kilgore Seed Company, Florida's leading seedsmen.

WATERMELON

Plant 1 pound of seed per acre. In the garden plant 1 packet to 75 feet of row for a family of 3 to 4 people.

Culture. Watermelons may be grown on nearly all kinds of Florida soils that have good drainage, but rolling, sandy pine land is preferred. In general, new land is best for this crop in order to avoid diseases. The field should be checked off, planting in hills 8 feet apart each way or 7 by 9 feet. Plant ½ to 1 inch deep, and plants thinned to 2 or 3 plants per hill when they are still small and when all danger of frost is past, later thinning to 1 plant per hill. The most successful growers apply 600 to 800 lbs. of fertilizer mixture about a week before planting the seed. A second application of 400 to 600 lbs. of fertilizer or top dressing may be made if necessary when the vines begin to run. The first plant-



Congo

starts early in December. In central Florida starts early in December. In central Florida planting begins January 1st and extends to about February 15th. Plantings in north Florida are made 2 to 3 weeks later. For earlier melons use **Hotkaps.** (See page 88.)

For pollination of watermelons, see third paragraph under cucumber culture—page 19.

All our melon seed is now treated to control seedborne diseases and to give better stands especially if planted early in cold ground. For disease and insect control (see page 70.)

The number of days after each variety indicates the time from seed planting to ripe fruit.

For protection from rats and birds use Kilgore's Melon Seed Treatment. (See page 76.)

All prices quoted are postpaid.

Write for special prices on larger quantities.

GARRISONIAN. (90 days.) Resembles Garrison in size, shape, color and quality of flesh; but the very important difference is its resistance to anthracnose. Growers who specialize in extra large and attractive melons should try this. Fruits: long, distinctly striped with light and dark green; weight about 40 pounds. Flesh bright red and sweet. Rind harder than Garrison making it a much better shipper. Developed by Southeastern Vegetable Breeding Laboratory from the same parents that produced Charleston Gray.

Pkt. 10c; oz. 25c; 1/4 lb. 65c

VERIFIED*

1 lb. \$2.25; 5 to 25 lbs. \$2.15 per lb.

HOPE DIAMOND. (92 days.) Developed at the Arkansas A. E. S. as a fusarium and anthracnoseresistant variety of Black Diamond, or Florida Giant type. The fruits average slightly smaller, with brighter red flesh and sweeter flavor, free from white heart and hollow heart.

Pkt. 10c; oz. 25c; 1/4 lb. 75c

VERIFIED*


1 lb. \$2.50; 5 to 25 lbs. \$2.40 per lb.

CONGO. (90 days.) This outstanding variety is resistant to both fusarium wilt and anthracnose diseases. The vines remain green longer than other varieties to protect the melons from sun burning. Congo has an exceedingly hard, tough rind which resists breakage and bruising in transit. In addition to these good qualities it possesses many other desirable features. It is a heavy yielder of large size melons, under good growing conditions ranging from 32 to 40 pounds. Similar in size and shape to Garrison but having a darker green rind with darker stripes. It is free of the tendency to produce gourd necks. Flesh of attractive bright red color, of excellent quality and flavor, ripening well out to the rind, with small white seeds having light tan smears, and very attractive interior appearance. This variety does not need as heavy fertilization as other varieties, and little, if any, side dressing with nitrogen fertilizer should be used because heavy nitrogen may cause hollow heart or separation of the flesh. Melons of this variety should be harvested just before full maturity to avoid hollow hearts.

Pkt. 10c; oz. 20c; 1/4 lb. 55c

VERIFIED*

1 lb. \$1.80; 5 to 25 lbs. \$1.70 per lb.



Garrisonian

WATERMELON (Continued)

FLORIDA GIANT (BLACK DIAMOND or CANNON BALL). (92 days.) The vines are of vigorous growth and produce large, nearly round, solid dark greenskinned melons, with thick rind and firm, rather coarse grained and fibrous, sweet, deep red, attractive flesh with small black seeds. Very hardy, and withstands poor growing conditions better than most other varieties. Because of the dark green rind, it is in great demand, always having a fresh appearance. Has grown in favor among commercial melon growers in Florida. Also a desirable home garden variety.

Pkt. 10c; oz. 15c; 1/4 lb. 45c


VERIFIED*

1 lb. \$1.80; 5 to 25 lbs. \$1.70 per lb.

BLACKSTONE. (90 days.) A new anthracnose resistant variety of Black Diamond type for shipping developed at the U.S. Regional Vegetable Breeding Laboratory, Charleston, S. C. Earlier maturing than Black Diamond, with better flesh quality and higher sugar content; intended for commercial production and home garden use throughout the South. Not resistant to fusarium wilt. Fruits large, 30 to 35 lbs., with dark green rind and yellowish ground spot. Flesh bright red, crisp, and sweet. Seeds large, stippled black.

Pkt. 10c; oz. 20c; 1/4 lb. 60c

1 lb. \$2.00; 5 to 25 lbs. \$1.90 per lb.



Blackstone

SUGAR BABY — A small icebox type watermelon weighing from 6 to 10 lbs. Early and prolific, perfectly round, with a thin hard tough rind, which is desirable for shipping. The melons are distinctly striped when immature, but become almost black when ripe. Inside appearance and flavor good. This melon is considered superior to other types of the icebox melons presently grown.

Pkt. 10c; oz. 30c; 1/4 lb. 90c


VERIFIED*

1 lb. \$2.30; 5 to 25 lbs. \$2.20 per lb.

*VERIFIED WATERMELON SEED

Associated Seed Growers, Inc. verifies that these seeds were: (1) grown in California, in an area free from anthracnose; (2) grown from seed stock tested and approved by the Asgrow Breeding and Research Department; and (3) are the product of a crop that was inspected by trained Asgrow fieldmen and found to meet the standards for freedom from disease and for trueness to variety type established by the Asgrow Quality Control Department.

> ASSOCIATED SEED GROWERS, Inc. New Haven, Connecticut



CHARLESTON GRAY. (92 days.) Developed by the U.S.D.A., Southeastern Breeding Station at Charleston, S. C. It is highly resistant to both fusarium wilt and anthracnose. It makes a long, gray melon very slow to "sunburn," with a tough but not particularly thick rind, very striking in appearance when cut with red meat and black seed. It is of delicious flavor and quality. A good yielder and very uniform in size, averaging under good growing conditions about 28 to 35 lbs. Will stay in the field in good condition for several days after it is ripe. This, along with its high resistance to the two most serious diseases of watermelons, makes it especially valuable for home garden use and to hold for a strong market. This new variety offers much promise as a high quality, disease-resistant, good shipping type melon for Florida.

Pkt. 10c; oz. 20c; 1/4 lb. 60c

VERIFIED*

1 lb. \$2.00; 5 to 25 lbs. \$1.90 per lb.

NEW HAMPSHIRE MIDGET. (65 days.) This small icebox type melon, weighing 3 to 5 lbs. is attracting attention in Florida for home use, roadside stands,



New Hampshire Midget

and local markets. It is the earliest of all watermelons making ripe fruits in from sixty to seventy days from seed. The vines lack vigor, but are prolific, setting many small melons, round to slightly elongated in shape with a light gray mottled rind, which is very thin and tender. Flesh is a beautiful deep red with small black seeds.

Pkt. 10c; oz. 20c; 1/4 lb. 55c

VERIFIED*

1 lb. \$1.85; 5 to 25 lbs. \$1.75 per lb.



FLOWER SEED

All Flower Seeds are subject to the Florida State Sales Tax. Rate of sales tax—10c to 35c—1c tax; 36c to 65c—2c tax; 66c to \$1.00—3c tax.

All flower seeds priced at 15c per packet All prices quoted are postpaid Write for prices on larger quantities What to plant, when and how is listed on Calendar, Pages 5 and 6

If interested in any straight colors, not listed, of any of the following kinds of lowerers, write The Kilgore Company, Plant City, Florida, stating your desires and ask us to quote you prices.

AGERTATUM-See Floss Flower-Page 42

ALYSSUM

Alyssum makes an attractive winter border or edging plant, covered with tiny blooms which form a compact showy carpet lasting for months through the winter season if kept cut back. Set plants in garden or thin to 6 inches apart in 1 foot rows.

Little Gem (White)—Small white flowers completely cover the dwarf, compact plants. Wonderful for edgings, of easy culture and extremely hardy

ings, of easy culture and extremely hardy.

Pkt. 15c 1/8 oz. 30c; 1/4 oz. 55c; 11/2 oz. 80c

1 oz. \$1.40; 1/4 lb. \$4.75

Royal Carpet—This Alyssum is very dwarf, growing only 2 to 3 inches tall and 10 inches across. Makes a sheet of rich, deep blue-purple color and the flowers make a solid mass so thick it's hard to see any green leaves. Ideal for edging paths and flower beds or for mass planting. For continuous bloom shear the full grown plants lightly.

Pkt. 15c; 1/16 oz. 50c; ½ oz. 80c; ¼ oz. \$1.40 ½ oz. \$2.25; 1 oz. \$4.00



Wilt Resistant Crego Aster (Mixed)

ASTERS

All Wilt Resistant Varieties

Used for cut flowers, bedding and in rock gardens. Sow seed in flats or seedbeds. Set plants in rather rich, well drained soil, protected from too much midday and afternoon sun. All mixtures are a blending of blue, crimson, pink, purple and white.

Crego Mixture—This variety, also called "Ostrich Feather," blooms a few days after the Imperial Mixture. The double blooms have long, shaggy, twisted petals and are borne abundantly on branching plants of robust growth. Plant 18 inches apart in row.

of robust growth. Plant 18 inches apart in row.

Pkt. 15c; /16 oz. 30c; ½ oz. 55c; ¼ oz. 90c

½ oz. \$1.50; 1 oz. \$2.50; ¼ lb. \$8.00

Imperial Mixture—A blending of the best colors of Early Bauty Asters. Plant 18 inches apart in row. Pkt. 15c; 1/16 oz. 45c; ½8 oz. 75c; ½ oz. \$1.25
½ oz. \$2.00; 1 oz. \$3.50; ½ lb. \$10.50

Powderpuffs (Mixed)—A mixture of bouquet type plants. Grows upright, bearing quilled flowers on long stems. Very early.

Pkt. 15c; 1/16 oz. 60c; ½ oz. \$1.10; ¼ oz. \$1.85 ½ oz. \$3.00; 1 oz. \$5.00

Princess (Mixed)—Large blooms having full crested center of small quilled petals with a narrow collar of guard petals.

Pkt. 15c; 1/16 oz. 50c; ½8 oz. 80c; ¼ oz. \$1.40 ½ oz. \$2.25; 1 oz. \$4.00

BABY'S BREATH

Baby's Breath is one of the necessities in every Florida garden where flowers are raised for cutting. Does best in rich soil in semi-shade. Sow seed in garden, thinning plants to 18 inches apart in 3 foot rows.

Pure White Pkt. 15c; 1 oz. 35c; 1/4 lb. \$1.00

BACHELOR BUTTON

The Bachelor Button or Cornflower is very easily grown. We offer only the double varieties. They are free blooming, and must have plenty of sun; excellent for bouquets, borders and backgrounds.

Sky Blue—One of the most beautiful and deepest blues in the flower kingdom.

Dainty Mixture—Composed of white, delicate pink, lavender and blue flowers.

Each of the above: Pkt. 15c; ½ oz. 35c; ½ oz. 60c; 1 oz. \$1.00; ½ lb. \$3.00

BALSAM (Double Camellia-Flowered)

Well-known old garden favorite with large, double flowers borne in axils of the leaves. Sow seeds in flats or seedbeds. Plant 12 inches apart in semi-shady spot.

Selected Mixture—Finest double blooms, white, light ping, rose, crimson and scarlet.

Pkt. 15c; 1/4 oz. 50c; 1/2 oz. 75c; 1 oz. \$1.25; 1/4 lb. \$4.00

BEAN-SCARLET RUNNER

Plants grow quickly, making strong climbing plants for shade, with showy sprays of attractive bright scarlet flowers. Grown principally as an ornamental vine.

Pkt. 15c; 1/4 lb. 35c; 1 lb. \$1.25

BELLS OF IRELAND

An old favorite, for flower arrangements. Odd green bells on 2 foot spikes. Plant seeds in sandy loam about one-quarter inch deep. An easy to grow annual useful for bedding also.

> Pkt. 15c; ½ oz. 35c; ¼ oz. 60c; ½ oz. \$1.00 1 oz. \$1.50; ¼ lb. \$5.00



Bells of Ireland




BLANKET FLOWER OR GAILLARDIA

These are brilliant daisy-like flowers of yellow, orange, and dark crimson hues, very desirable for cutting and for garden display. Easily grown, require a sandy, moist, well drained soil and a sunny location. Sow seed in garden, spacing plants 18 to 24 inches apart each way.

Double Gaiety Mixture.

Pkt. 15c; $\frac{1}{8}$ oz. 35c; $\frac{1}{4}$ oz. 60c; $\frac{1}{2}$ oz. \$1.00; 1 oz. \$1.50



Blue-Eyed African Daisy-Arctotis Grandis

BLUE-EYED AFRICAN DAISY OR ARCTOTIS

A charming flower for cutting as well as for garden display. Plants produce many branches with long-stemmed, pure white daisy-shaped flowers having a blue center and pale lilac-blue shading on the reverse of the petals.


Pki. 15c; ½ oz. 35c; ½ oz. 60c; 1 oz. \$1.00; ¼ lb. \$3.50

BLUE LACE FLOWER

A delightful blue annual, growing quickly from seed, producing long-stemmed and clustered heads. Sow seed in flats or seedbeds, setting plants in garden about 3 feet apart each way.

Heavenly Blue

Pkt. 15c; 1/16 oz. 30c; ½ oz. 55c; ¼ oz. 90c ½ oz. \$1.50; 1 oz. \$2.50; ¼ lb. \$8.00



Blue Lace Flower or Didiscus

CALENDULA

The Calendula is one of the finest, and showiest of the free-flowering winter annuals for Florida. It grows quickly from seed, transplants easily, and is not affected by the coldest weather in Florida. Sow seed in flats or in garden.

Campfire Improved—Dark orange, scarlet sheen. Pkt. 15c; 1/4 oz. 35c; 1/2 oz. 60c; 1 oz. \$1.00; 1/4 lb. \$3.00

Lemon Queen—Light golden yellow, light center. Pkt. 15c; 1/4 oz. 35c; 1/2 oz. 60c; 1 oz. \$1.00; 1/4 lb. \$3.00

Excellent Mixture—Made up of shades of orange and yellow.

Pkt. 15c; 1/4 oz. 35c; 1/2 oz. 60c; 1 oz. \$1.00; 1/4 lb. \$3.00

Art Shades Mixture—Contains delicate shades. Pkt. 15c; 1/4 oz. 55c; 1/2 oz. 90c; 1 oz. \$1.50; 1/4 lb. \$4.50



Calendula—Campfire Improved

CALIFORNIA POPPY

Beautiful, low-growing, spreading plants of easy culture in sunny locations. Gray-green, fern-like foliage with bright colored cup-shaped blooms. Sow seed thinly in well prepared beds where they are to grow.

Gorgeous Mixture. (Single Flowering)—A fine mixture containing carmine, orange-scarlet, as well as white, yellow and orange shades.


Pkt. 15c; 1/4 oz. \$35c; 1/2 oz. 60c; 1 oz. \$1.00; 1/4 lb. \$3.00

CALLIOPSIS

Bushy, wiry plants, producing a profusion of bicolor, semi-double blooms. Requires very little care. Plant in groups, borders or backgrounds, giving plenty of room, and a sunny location.

Choice Mixture—Yellow, gold, brown and maroon bi-colors.

Pkt. 15c; 1/4 oz. 35c; 1/2 oz. 60c; 1 oz. \$1.00; 1/4 lb. \$3.00



Carnation—Charm Mixture—Chabaud's Giant Improved Strain

CARNATION

Improved Chabaud strain, superior to other outdoor types for cut flowers. Each plant produces many stalks of very large fragrant flowers. Sow seed in flats or seedbeds and transplant, allowing about 12 inches for each plant.

Charm Mixture.

Pkt. 15c; 1/16 oz. 35c; ½8 oz. 60c; ¼ oz. \$1.00 ½ oz. \$1.75; 1 oz. \$3.00

CANDYTUFT

Charming clustered heads of daintily colored flowers. Blooms from autumn to spring, preferring the cooler weather. It can be transplanted, but is better if sown where it is to bloom and thinned out to about 12 inches apart in rows.

White Hyacinth-Flowered. This is the Coronaria strain which holds its blooms in spikes like hyacinths, with several spikes per plant.

Pkt. 15c; 1/4 oz. 35c; 1/2 oz. 60c; oz. \$1.00; 1/4 lb. \$3.00

Borden Group (Umbellata)—This type grows about 16 inches tall, the plants being globe shaped with flowers covering the plant in a mass. The mixture contains shades of crimson, pink, lavender, rose and white.

Pkt. 15c; 1/4 oz. 35c; 1/2 oz. 60c; 1 oz. \$1.00; 1/4 lb. \$3.00

CELOSIA (Cockscomb)

Easily grown in any soil. Flowers may be dried for lasting bouquets. Sow seed in $1\frac{1}{2}$ to 2 foot rows during the winter and early spring, and thin to about 12 inches in the row.

Tall Royal Velvet — Comb Type — Rich carmine-crimson, foliage bronze green, reddish margins. 18 inches tall.

Pkt. 15c; 1/16 oz. 60c; ½ oz. \$1.00; ¼ oz. \$1.50 ½ oz. \$2.50; 1 oz. \$4.50

Pampas Plume—Plume Type. Mixture of long stemmed plumes in bronze, gold, henna, scarlet, copper and crimson shades.

Pkt. 15c; ½ oz. 50c; ¼ oz. 70c; ½ oz. \$1.20 1 oz. \$2.00; ¼ lb. \$7.00

CHINESE FORGET-ME-NOT OR CYNOGLOSSUM

A beautiful bushy plant, covered with sprays of lovely sky-blue flowers. Used with other flowers for bouquets. Fine for bedding and rock gardens. Sky Blue

Pkt. 15c; ½ oz. 30c; 1 oz. 50c; ¼ lb. \$1.75

CLEOME (Spider Plant)

Large trusses of salmon-pink blooms all summer. Excellent for bedding, borders and backgrounds.

Pkt. 15c; ½ oz. \$30c; ¼ oz. 50c; ½ oz. 75c l oz. \$1.25; ¼ lb. \$4.00

COSMOS

Plants covered with finely divided, bright green, fern-like foliage. The blooms are large and double, or single with a crested center.

Mandarin—A brilliant orange flower, often having 40 or more petals. Very heavy, dense foliage.

Pkt. 15c; 1/16 oz. 45c; ½ oz. 75c; ¼ oz. \$1.25 ½ oz. \$2.00; 1 oz. \$3.50

Fiesta Semi-Double—A brilliant, semi-double bloom of golden orange, striped with red, borne freely on rather bushy, sparsely foliaged plants.

Pki. 15c; 1/16 oz. 30c; 1/8 oz. 55c; 1/4 oz. 90c 1/2 oz. \$1.50; 1 oz. \$2.50

Choice Mixture—Plants 2½ to 3 feet tall, free blooming with attractive tufted centers. An excellent mixture in colors of crimson, pink, white and rose.

Pkt. 15c; 1/16 oz. 70c; ½ oz. \$1.00; ¼ oz. \$1.75 ½ oz. \$2.75; 1 oz. \$4.75



Cosmos - Mandarin

DAHLIA

Useful for bouquets and bedding. Sow seed in early spring in flats or seedbed and transplant to 12 inches apart in 2 foot rows. Easy to grow from seed.

Unwin's Bedding Mixture—Exquisite, semi-double dwarf dahlias in a varied assortment of attractive colors. Early flowering, bushy plants.

Pkt. 15c; 1/16 oz. 35c; ½ oz. 60c; ¼ oz. \$1.00 ½ oz. \$1.75; 1 oz. \$3.00; ¼ lb. \$10.00


DELPHINIUM (Perennial Larkspur)

Useful for bouquets and borders. Sow seed in flats or seedbeds or direct in garden in 3 foot rows and thin or transplant to from 12 to 18 inches in the row. To help germination it is desirable to place seed in moist cloth in the ice compartment of a refrigerator for about forty-eight hours before sowing. A constant moisture supply must be provided. Are used for display and cutting. They do best in cool weather.

Belladonna Hybrids Mixed—Single flowers of large size in solid erect spikes, shades of blue. Highly prized for cutting. This is the best type for Florida.

Pkt. 15c; 1/16 oz. 35c; ½ oz. 60c; ¼ oz. \$1.00

½ oz. \$1.75; 1 oz. \$3.00



Delphinium—Belladonna Hybirds Mixed

FLOSS FLOWER OR AGERATUM

Ageratum is popularly known as the Floss Flower. It is a favorite garden cut flower, and is used for edgings, borders or bedding in Florida. Plant grows rapidly, has light green foliage, and many clusters of small, tassellike flowers. Blooms all summer. Sow seed in flats or seedbed.

Dwarf Blue Cap—Dwarf, growing to a height of about six inches. Compact habit, small foliage, and symmetrical dome-like form. The color is a deep, rich, bright blue-lavender.

Pkt. 15c; 1/16 oz. 60c; 1/8 oz. \$1.00; 1/4 oz. \$1.50; 1/2 oz. \$2.50; 1 oz. \$4.50

Blue Perfection—Compact plant, growing to a height of about 18 to 24 inches, covered with true blue flowers. Desirable for cutting.

Pkt. 15c; 1/16 oz. 35c; 1/8 oz. 60c; 1/4 oz. \$1.00 1/2 oz. \$1.75; 1 oz. \$3.00; 1/4 lb. \$10.50

GLOBE AMARANTH (Gomphrena or Florida Bachelor Button)

This is a popular old-time favorite having clover-like blooms in shades of white, purple and rose. Can be used for dried bouquets if cut and hung in a dark place to dry. Can be depended upon to succeed under almost any conditions in Florida during the summer. Sow seeds in flats or seedbed and transplant, spacing 12 inches in row.

Assorted Colors

Pkt. 15c; ½ oz. 30c; 1 oz. 50c; ¼ lb. \$1.75

GOURD-ORNAMENTAL

Gourds have become esteemed for many uses. The small fancy fruits are of multitudinous shapes, sizes and colors. The fruits are oddly shaped and highly colored. Vines should be supported on a trellis or fence, spacing plants about 3 feet apart. The seed should be sown in a well drained, sunny location when danger of frost is passed.

Small Varieties Mixed

Pkt. 15c; 1 oz. 40c; 1/4 lb. \$1.50

HOLLYHOCK (Annual)

These types of hollyhock are well adapted to the climatic conditions of Florida. Thrives best on rich, well drained soil, but needs plenty of water. Seed should be planted in the garden, spacing 24 inches in the row. Hollyhock is difficult to transplant. Makes an ideal background.

Indian Spring—Produces a colorful display of single and semi-double blooms, three to four inches across, borne on a central spike. Shades of pink, rose, crimson, lavender, yellow and white.

Pkt. 15c; $\frac{1}{8}$ oz. 30c; $\frac{1}{4}$ oz. 50c; $\frac{1}{2}$ oz. 75c; 1 oz. \$1.25

Double Triumph Mixed—Frilled, graceful rosette blooms in many beautiful shades, including maroon, crimson, scarlet, rose, pink, white and lavender.


Pkt. 15c; ½ oz. 50c; ¼ oz. 80c; ½ oz. \$1.40; 1 oz. \$2.25

LARKSPUR

Larkspur is a cool weather plant, so it is best to wait until November to plant. Seeds will fail to germinate if planted in hot weather. Sow seed 1/8 inch deep and water with fine spray. Thin to stand 12 to 18 inches apart in 3 foot rows. The young plants are hardy and like full sun, plenty of fertilizer and water.

Regal Finest Mixed-This is the branching type of Larkspur which has been greatly improved by selecting for doubleness and size of flowers, new colors and better habit of growth. The plants are well branched and produce many long, thick spikes closely set with a large array of Delphinium-like florets of mixed colors, including blues, carmine, rose, white, etc. Showy in beds and lovely for cutting. Vigorous and early.

Pkt. 15c; 1/8 oz. 30c; 1/4 oz. 50c; 1/2 oz. 75c; 1 oz. \$1.25



Larkspur-Regal Finest Mixed

LINARIA (Baby Snapdragon)

Linaria blooms heavily, but for a limited period, making successive plantings desirable. The plants produce spikes of dainty snapdragon-like flowers. Resists frost. Very bright mixtures of desirable colors, scarlet, chartreuse, white, cerise, yellow, purple, rose, scarletorange, cream and lilac.

Northern Lights—An improved, taller growing Linaria. Very spectacular with wide range of color including darker shades.


Pkt. 15c; 1/8 oz. 30c; 1/4 oz. 55c; 1/2 oz. 80c; 1 oz. \$1.40

LUPINE (Annual)

Plants make decorative, palm-like foliage and stems covered with bristly hairs. Blooms are pea-shaped, borne in spires of varying length. Desirable for cutting and bedding. Also makes good backgrounds. Thin the plants to 12 inches apart in three foot rows. They will grow in comparatively poor soil, but they like full sun. Will also flourish in partial shade, and require an abundance of water during the blooming season. Assorted shades of dark blue, light blue, rose and white.

Assorted Colors

Pkt. 15c; 1/8 oz. 30c; 1/4 oz. 55c; 1/2 oz. 80c; 1 oz. \$1.40



Linaria - Northern Lights

MARIGOLD


The attractive plants, with fern-like foliage, produce gigantic, double, frilled flowers. The dwarf types for bedding and edging, the tall types for backgrounds. All types are magnificent when cut for bouquets. Marigolds withstand heat and drouth, thriving where many plants would perish.

Crown O' Gold (collarette) — Flowers are bright golden-orange color, having a collar of big, wide, drooping petals. The only Marigold with odorless foliage and fragrant flowers.

Pkt. 15c; 1/16 oz. 30c; ½ oz. 50c; ¼ oz. 75c
½ oz. \$1.25; 1 oz. \$2.00; ¼ lb. \$6.75

Crackerjack—A giant flowered Marigold, similar to Sunset Giants, but much earlier. The plants have a superior habit of growth, are compact and sturdy, approximately 2½ feet tall. The blooms are uniformly large and double, measuring three to four inches across. They have the full range of marigold colors, primrose, yellow, gold and orange.

Pkt. 15c; 1/16 oz. 35c; ½8 oz. 60c; ¼ oz. \$1.00 ½ oz. \$1.75; 1 oz. \$2.75; ¼ lb. \$8.00



Marigold—Crackerjack

Real Gold—Carnation Type—Very large blooms; pure gold color, plants very uniform, 20 to 24 inches high.


Pkt. 15c; 1/16 oz. 35c; ½8 oz. 60c; ¼ oz. \$1.00 ½ oz. \$1.50; ¼ lb. \$5.00

Yellow Supreme—Carnation Type—free flowering compact, and somewhat pyramidal in habit. The color is a cream lemon-yellow which fairly glows and sparkles.

Pkt. 15c; ½ oz. 50c; ¼ oz. 75c; 1 oz. \$1.25 1 oz. \$2.00; ¼ lb. \$6.75

Gigantea Sunset Giants Mixture—Tall, free flowering. The most remarkable feature is the delicate sweet fragrance of the blossoms. The stems are long and the flowers of startling bold colors of varied shades of lemon-yellow to gold-orange, are excellent keepers after cutting.

Pkt. 15c; ½ oz. 35c; ¼ oz. 60c; ½ oz. \$1.00 1 oz. \$1.50; ¼ lb. \$5.00



Marigold — Gigantea Sunset
Giants Mixture
(Continued on next page)

MARIGOLD—(Continued)

Dwarf Gigantea Pot O' Gold—A dwarf, sweet scented counterpart of Gigantea type. Plants are dwarf and compact, 12 to 15 inches in height. Flowers are double, deep golden-orange. Early and free blooming.

Pkt. 15c; ½ oz. 30c; ¼ oz. 50c; ½ oz. 75c 1 oz. \$1.25; ¼ lb. \$4.00

Dwarf French Double Harmony—An excellent dwarf French Marigold, growing to a height of about 12 inches. Free-flowering with a bright golden-yellow center, bordered with a dark maroon-brown color. Plants are of bushy habit and form excellent material for bedding or borders.

Pkt. 15c; ½ oz. 30c; ¼ oz. 50c; ½ oz. 75c 1 oz. \$1.25; ¼ 1b. \$4.00



Marigold — Glitters

Glitters—Chrysanthemum Flowered—Large, fluffy, bright canary yellow chrysanthemum-like flowers. Makes a bushy, compact, erect and free flowing plant with long strong stems.

Pkt. 15c; ½ oz. 50c; ¼ oz. 75c; ½ oz. \$1.25 1 oz. \$2.00; ¼ lb. \$6.75

Extra-Dwarf Double Mixed—An excellent mixture of free flowering dwarf double varieties. Large blooms in orange and yellow, as well as combinations of these colors. Plants grow 10 to 15 inches tall.

Pkt. 15c; ½ oz. 30c; ¼ oz. 55c; ½ oz. 80c 1 oz. \$1.40; ¼ lb. \$4.75

Lemondrop—Canary yellow, full double, free blooming. Plants grow eight to nine inches high.

Pki. 15c; ½ oz. 30c; ¼ oz. 50c; ½ oz. 75c 1 oz. \$1.25; ¼ lb. \$4.00

MORNING GLORY Improved Ipomea

This is the favorite quick growing vine with large showy attractive blossoms for shading porches, covering trellises and fences. Sow seeds after frost, when soil is warm. Germination may be hastened by cutting or filing through the hard outer shell and soaking in warm water for twenty-four hours before planting.

Scarlett O'Hara (Deep Wine Red)—Flowers 3½-4 inches across, borne profusely on vigorous climbing vines.

Pkt. 15c; ¹/₄ oz. 35c; ¹/₂ oz. 60c; 1 oz. \$1.00 ¹/₄ 1b. \$3.00 **Pearly Gates** (White)—The large flowers are a lovely lustrous white, and literally cover the vigorous, fast growing vines.

Pkt. 15c; ½ oz. 30c; ¼ oz. 50c; ½ oz. 75c 1 oz. \$1.25; ¼ lb. \$4.00

Heavenly Blue Improved—The large sky blue flowers stay open late in the day.

Pkt. 15c; 1/4 oz. 30c; 1/2 oz. 50c; 1 oz. 75c
1/4 lb. \$2.50


NASTURTIUM

The nasturtium is very easily grown, and succeeds best in rather dry, sandy or gravelly, well-drained but well fertilized soil. The large, double and semi-double, sweet-scented flowers are valuable for cutting. The plants have a long blossoming period (3-4 months) provided the flowers are kept picked off before seed is formed. Flowers are borne on long stems, well above the foliage, making them very attractive for bedding plants and as cut flowers for bouquets. Soaking seed in water over night before planting in moist soil helps germination. Plant seed in 3 foot rows and thin to 12 inches in the row. If covered on cold nights, plants will grow and bloom all winter.

NASTURTIUM VARIETIES

Semi-Double Mahogany Gleam—Large blooms, rich mahogany-red color. Very fine for cutting.

Pkt. 15c; 1 oz. 35c; 1/4 lb. \$1.00; 1 lb. \$3.00



Nasturtium — Semi-Double Golden Gleam

Semi-Double Golden Gleam—Large, deep goldenyellow, sweet-scented blooms, borne on long stems well above the foliage. A blaze of intense, gleaming, golden-yellow color when in full bloom.

Pkt. 15c; 1 oz. 35c; 1/4 lb. \$1.00; 1 lb. \$3.00

Semi-Double Scarlet Gleam—The flowers are a fiery orange-scarlet. The plants bear the numerous flowers well above the foliage.

Pkt. 15c; 1 oz. 35c; 1/4 lb. \$1.00; 1 lb. \$3.00

(Continued on next page)



NASTURTIUM—(Continued)



Nasturtium - Semi-Double Gleam Hybrids

Semi-Double Gleam Hybrids—The hybrids are a collection of gorgeously colored, large, well formed flowers of dazzling beauty. A very brilliant mixture of an amazing range of showy colors. There are a few singles among the blossoms, although mostly double and semi-double.

Pkt. 15c; 1 oz. 35c; 1/4 lb. \$1.00; 1 lb. \$3.00



Nasturtium — Semi-Double Dwarf Gem Mixture

Semi-Double Dwarf Gem Mixture—Dwarf, compact, bushy or globular plants about one foot high. Blooms are sweet-scented of excellent color range.

Pkt. 15c; 1 oz. 35c; 1/4 lb. \$1.00; 1 lb. \$3.00

PANSY

Pansies are the most ideal of all flowering plants for edging or for bedding in late winter and early spring. The pansy is distinctly a cool weather plant, and for a long blooming period it should be grown in cool weather and in a cool location. Seeds will not germinate well in hot weather. Sow seeds very shallow in flats or in well prepared seedbed soil in a shaded place in late fall. Transplant seedlings 12 to 18 inches apart, giving some shade and plenty of water.



Engelmann Giant Pansy

Engelmann Giant Mixture—This is a large flowered pansy. Flowers have thick, velvety petals, borne on long stems, and are of most striking brilliance and endless variety of color and design.

Pki. 15c; 1/16 oz. \$1.75; ½ oz. \$2.75; ¼ oz. \$5.00

PETUNIA

Petunias are the most colorful of annual garden flowers. They will grow in practically any soil and require very little attention after they are established.


The culture of petunias is simple and quite easy although many amateurs find them somewhat difficult to start, due to the fact that they cover the seed too deeply. Whether sown directly in the garden or as is generally recommended in flats or seedbed, the seeds must not be buried. The very small seeds should be broadcast thinly on a well-prepared seedbed, pressed into the soil, and then covered very lightly to a depth of not more than 1/16 to 1/8 of an inch, using sand or better, sand mixed with peat moss or humus. The moisture content must be kept uniform until the seeds have sprouted, and the beds should never be allowed to dry out after seeds are sown. It is best to start seed-lings in a flat or seedbed in all cases where moisture conditions can be controlled. The seedbeds should be covered with moist burlap or muslin until sprouts appear, when this shading material should be removed immediately.

When the seedlings are well grown in flats or seedbeds, transplant to the garden or border, setting plants in rows 2 to 3 feet apart and 12 to 18 inches in the row. Petunias do best in moderately rich, well drained, moist soil. Since they grow and bloom over long periods they require a lasting fertilizer like pulverized sheep manure (see page 79). In transplanting, set plants the same depth they were before. If they have sent up a center shoot, it should be pinched to induce branching and make a more compact plant form.

(Continued on next page)



PETUNIA-(Continued)



Petunia-Carnival (F2 Hybrids)


Carnival (F2 Hybrids). This new hybrid petunia has attracted great attention wherever grown. The vigorous plants are bedding type. Blooms are large, with ringed and ruffled petal edges, also some smaller plain edged flowers. Colors include blue, white, crimson, rose, scarlet, salmon and cream-pink in solid colors; also some bi-colored. This variety exceeds anything of its type yet shown in vigor or growth, and range of color. Very free blooming.

Pkt. 15c; 1/16 oz. \$3.00; 1/8 oz. \$5.50; 1/4 oz. \$10.00

Fluffy Ruffles

These are among the most beautiful petunias and well worth the ambition of everyone to produce. The huge blossoms are wonderfully fringed and ruffled and range bewilderingly through shades of pink, blue, scarlet, lavender, crimson, purple, white, and intermediate hues, markings, and curious blotches. The flowers measure 2 to 3 inches across when properly grown. Suited for bedding purposes and for cutting, as well as for porch and window boxes. Sow seeds carefully.

Pkt. 15c; 1/16 oz. \$3.00; 1/8 oz. \$5.50; 1/4 oz. \$10.00



Petunia—Fire Chief

DWARF, ERECT PLAIN PETALED PETUNIAS

Grow 6 to 12 inches high.

Rose Morn—Dwarf, compact, constantly blooming plants with small, brilliant, rose colored flowers with white centers.

Pkt. 15c; 1/16 oz. 75c; ½ oz. \$1.25; ¼ oz. \$2.00 ½ oz. \$3.50; 1 oz. \$6.00

Heavenly Blue—Plant is dwarf in growth. Flowers are a beautiful light blue.

Pki. 15c; 1/16 oz. \$1.00; ½ oz. \$1.75; ¼ oz. \$3.00 ½ oz. \$5.00; 1 oz. \$9.50

Fire Chief—A free-flowering Petunia with a brilliant scarlet-red color, turning to a deep salmon-rose with age. Plants vigorous and upright, flowering over a long season of bloom.

Pkt. 15c; 1/16 oz. \$1.50; 1/8 oz. \$2.50; 1/4 oz. \$4.50

PLAIN PETALED PETUNIAS

Grow 15 to 18 inches high

Snow Queen—Large, pure white flowers borne on plants of compact growth.

Pkt. 15c; 1/16 oz. 60c; ½ oz. \$1.00; ¼ oz. \$1.50 ½ oz. \$2.50; 1 oz. \$4.50



Petunia-Superb Mixture

Superb Mixture—A well selected blend of the finest types and deepest, richest colors of the single plain petaled, tall growing petunias.

Pkt. 15c; 1/16 oz. 45c; ½ oz. 75c; ¼ oz. \$1.25 ½ oz. \$2.00; 1 oz. \$3.50; ¼ 1b. \$10.00

Howard's Star Improved—Crimson maroon flowers with a clearly defined star of pure white.

Pkt. 15c 1/16 oz. 75c; ½ oz. \$1.25; ¼ oz. \$2.00 ½ oz. \$3.50; 1 oz. \$6.00

Elk's Pride (Elk's Purple)—Blooms are darkest velvety purple. This remarkably rich and even tone is carried over the entire surface of the flower.

Pkt. 15c; 1/16 oz. \$3.00; 1/8 oz. \$5.50; 1/4 oz. \$10.00

Flaming Velvet—Deep, velvety blood-red. Very beautiful and showy.

Pkt. 15c; 1/16 oz. 60c; ½ oz. \$1.00; ¼ oz. \$1.50 ½ oz. \$2.50; 1 oz. \$4.50

PHLOX

Blooms practically all year. Plant in garden about 1/4 inch deep, and transplant seedlings or thin to 6 inches apart in 2 foot rows. Grows easily in most any type of soil, but likes plenty of water and sunshine. Excellent for bedding in gardens, planting in barren places, beautifying vacant lots and roadsides, and for bordering gardens. Also valuable for rock gardens. Phlox is relatively free from pests. Transplants easily.

Peninsular Mixture

Pkt. 15c; 1/16 oz. 30c; ½ oz. 50c; ¼ oz. 80c; ½ oz. \$1.25 1 oz. \$2.25; ¼ lb. \$7.50; 1 lb. \$25.00



Phlox—Peninsular Mixture

PINKS-CHINESE OR DIANTHUS

The Dianthus, or Chinese Pinks, are extremely popular because of their varied and lovely rich coloring. The flowers produce in clusters. They are hardy, heat resistant and do well in Florida. Fine for massing, bordering, and cut-flowers. Plant seed about 6 inches apart in 2 foot rows.

Medal Mixture — Flowers produced in clusters of double blooms, free flowering, in solid colors and combinations of white, pink, rose, maroon, scarlet and blush.

Pkt. 15c; 1/8 oz. 35c; 1/4 oz. 60c; 1/2 oz. \$1.00 1 oz. \$1.50; 1/4 lb. \$5.00



Dianthus-Medal Mixture

Sweet Wivelsfield — Multiflora type, remarkably free flowering and fragrant, with clusters of single blooms, sweet scented. Colors includes shades of white, pink, rose, scarlet, crimson, maroon, and salmon, some zoned and edged with contrasting shades.

Pkt. 15c; ½ oz. 35c; ¼ oz. 60c; ½ oz. \$1.00 1 oz. \$1.50; ¼ lb. \$5.00

Single Gaiety Mixed—Large lacy flowers in a brilliant range of colors, with very deeply cut petal edges. Blooms carry the attractive "pheasant eye" markings.

Pkt. 15c; ½ oz. 30c; ¼ oz. 50c; ½ oz. 75c 1 oz. \$1.25; ¼ lb. \$4.50

PORTULACA (See Sun Plant Page 48)

QUEEN ANNE'S LACE

Queen Anne's Lace has large heads made up of tiny white flowers, giving it a lace-like appearance. The flower clusters are borne on long stems which make this flower very suitable to mix with other cut flowers, giving the bouquet an airy appearance. Plant at least two feet apart in 4 foot rows as crowding stunts the growth of the plants. Valuable for backgrounds.

Pkt. 15c; 1/4 oz. 35c; 1/2 oz. 60c; 1 oz. \$1.00; 1/4 lb. \$3.00

SALVIA

Bonfire—Suitable for both bedding and borders of medium height. The plants bear spikes of very showy, intense scarlet flowers and make attractive bouquets. Sow seeds in flats or seedbed, transplanting to 3 foot rows, 18 inches in the row.

Pkt. 15c; 1/16 oz. 80c; $\frac{1}{18}$ oz. \$1.40; $\frac{1}{4}$ oz. \$2.25 $\frac{1}{2}$ oz. \$4.00; 1 oz. \$7.50

SCABIOSA OR MOURNING BRIDE

The showy, fragrant flowers of Scabiosa are borne on long, graceful stems over a long season. The colors are rich and varied—pure white, blue, red and variegated shades. As cut-flowers they are unsurpassed. Also useful for bedding.

Sunrise Mixture

Pkt. 15c; ½ oz. 35c; ¼ oz. 60c; ½ oz. \$1.00 1 oz. \$1.50; ¼ lb. \$5.50



Shasta Daisy-Single

SHASTA DAISY

Single and Semi-Double Giants—Beautiful flowers of pure glistening white, with bright yellow centers, Useful for cut flowers, and bedding. Sow seed in garden spacing 12 to 15 inches apart in 3 foot rows. Very hardy, but requires plenty of water.

Pkt. 15c; 1/16 oz. \$1.00; ½ oz. \$1.50; ¼ oz. \$2.50 ½ oz. \$4.50; 1 oz. \$8.00



Snapdragon-Giant Mixture

SNAPDRAGON OR ANTIRRHINUM

Giant Snapdragons — Successful only when grown through the winter and early spring months. The seeds are small and slow to germinate. They should be planted in a shallow box or seedbed of rich soil, in a cool, shady place and kept continuously moist until seeds sprout. Protect from ants. When 4 inches high, young plants may be transplanted 1 foot to 18 inches apart in rows 3 feet apart. As soon as the plants reach 8 inches in height they should be tied to light stakes until the side branches are well developed. Do best in rich loamy, well-drained soil with plenty of water. Most useful as cut flowers, as well as bedding.

Giant Mixture. (Rust Resistant). — This is a blend of the finest, outstanding shades of the tall growing (2 or 3 feet) type. Invaluable for cutting.

Pkt. 15c; 1/16 oz. 35c; ½8 oz. 60c; ¼ oz. \$1.00 ½ oz. \$1.50; 1 oz. \$2.50; ¼ lb. \$8.00

Giant Ruffled "Tetraploid" Mixture (Rust Resistant)—Large and attractive ruffled flowers in many colors and bi-colors some veined. Tall, erect plants (2½ to 3 ft.), strong and husky, with thick, dark green glossy foliage. Very long spikes.

Pkt. 15c; 1/16 oz. 60c; ½ oz. \$1.00; ¼ oz. \$1.50 ½ oz. \$2.50; ¼ lb. \$15.00

STATICE

Slender, wiry, bushy plants, bearing masses of flowers in delicate shades of lavender, rose, blue and white. Valued for drying, and makes an excellent long-lasting flower for bouquets, also attractive bedding plant. Sow seeds in the garden in 3 foot rows and thin to 18 inches in the row. Very easily grown. Requires sunny location.

Assorted Colors

Pkt. 15c; ½ oz. 30c; 1 oz. 55c; ¼ lb. \$1.50; 1 lb. \$5.50

Single Colors — Blue, Rose, Lavender, White or Yellow Pkt. 15c; ½ oz. 35c; 1 oz. 60c; ¼ lb. \$1.75; 1 lb. \$6.00

BE SURE TO INCLUDE STATE SALES TAX WHEN ORDERING FLOWER SEED. WE ARE NOT PERMITTED BY LAW TO ABSORB THE TAX.

Rate of Sales Tax. 10c to 35c—1c; 36c to 65c—2c; 66c to \$1.00—3c.

STRAWFLOWER OR HELICHRYSUM

Large, showy flowers of straw-like texture which retain their brilliant colors when dry, making them useful for decorative bouquets. The seed should be sown where plants are to bloom in 3 foot rows, and thinned out 18 inches apart. Plants grow 2 to 3 feet high and make a showy background.

Many Colors—A mixture containing fiery red, canary, silver, rose, salmon, and blue.

Pkt. 15c; 1/16 oz. 30c; ½ oz. 50c; ¼ oz. 75c; ½ oz. \$1.25 1 oz. \$2.00; ¼ lb. \$6.75



Strawflower or Helichrysum

SUNFLOWER (HELIANTHUS)

Easily grown and does best in a sunny location, it being a heat tolerant annual. Fine background for lower plantings, or can be grown in clumps. Successive sowings of seed can be made for cutting during the months of May thru September. Thin plants to 2 feet apart in 3 foot rows.

Sunshine Mixture — Single; Zoned bronze and red shades.

Pki. 15c; $\frac{1}{8}$ oz. 35c; $\frac{1}{4}$ oz. 60c; $\frac{1}{2}$ oz. \$1.00; 1 oz. \$1.50 $\frac{1}{4}$ lb. \$5.00

Sungold—Brilliant, golden yellow, flowers large and double, plants uniform height $3\frac{1}{2}$ to 4 ft.

Pkt. 15c; $\frac{1}{4}$ oz. 30c; $\frac{1}{2}$ oz. 50c; 1 oz. 75c; $\frac{1}{4}$ lb. \$2.50

SUN PLANT (PORTULACA) Blooms all Summer in Florida

A creeping, succulent plant, with odd, thick, fleshy stems and small, quill-shaped leaves, about 6 inches high, which thrives in sunny locations. Produces waxy flowers of varying colors which open only in sunshine early in the morning. It is excellent for edgings or borders. The seed is very fine; sow it thinly where plants are to grow, and do not cover. May be thinned 10 to 12 in. apart in 1 ft. rows.

Pkt. 15c; 1/16 oz. 45c; ½ oz. 75c; ¼ oz. \$1.30 ½ oz. \$2.25; 1 oz. \$3.75

SWEET PEAS EARLY FLOWERING SPENCER

The Early Flowering Spencer varieties are heat resistant and best adapted for Florida soil and climate. Sweet Peas do best in cool weather. They are heavy feeders and need very rich soil. Plant seeds one inch apart and cover two inches deep. When plants are five inches high, thin to three or four inches apart and furnish wire supports for vines to run upon. Fertilize and water frequently. Keep flowers cut off every day to prevent seed pods forming and to prolong the blooming period. Sunshine and fresh air are very essential.

Mrs. Herbert Hoover-Bright Blue.

Mars—Bright Crimson Red.

Shirley Temple—Rich Pink.

Hope-Pure White.

Lavanda—Pure Clear Lavender.

Ball's Orange Improved—Deep Orange.

Prices on each of the above: Pkt. 15c; 1 oz. 45c; 1/4 lb. \$1.10

Rainbow Mixture—Sweet Pea

Assortment of all popular shades of Early Flowering Spencer varieties.

Pkt. 15c; 1 oz. 45c; 1/4 lb. \$1.10; 1 lb. \$3.50

Zvolanek's Multiflowered Mixture—Sweet Pea

Vigorous and unusually free-flowering. On very long, strong stems, with many stems carrying 4 to 5 large flowers instead of the usual three or four smaller florets. This sweet pea is excellent for Florida.

Pkt. 15c; 1/4 oz. 35c; 1/2 oz. 60c; 1 oz. \$1.00 1/4 lb. \$3.00

TITHONIA OR FLORIDA SUNFLOWER

Makes ideal cut flowers, useful for backgrounds, and in shrubbery bays. Must have plenty of sun and a fertile soil. Plant seed in the garden and space plants 3 to 4 feet apart.

Torch—A bushy type with single, bright orange and scarlet flowers with yellow discs, borne on stout stems. Very early, heat resistant, and of easy culture with a long blooming season.

Pki. 15c; 1/16 oz. 35c; $\frac{1}{8}$ oz. 60c; $\frac{1}{4}$ oz. \$1.00 1/2 oz. \$1.75; 1 oz. \$3.00



Rainbow Mixture Sweet Peas

VERBENA

Verbenas are spreading, bushy, and cover themselves with heads of brilliant flowers in shades of blue, red, rose, lavender, white and scarlet. Seed should be soaked in water forty-eight hours in a refrigerator. Sow where they are to grow and thin to a foot apart in 2 foot rows.

Master Mixture

Pkt. 15c; 1/16 oz. 35c; 1/8 oz. 60c; 1/4 oz. \$1.00 1/2 oz. \$1.75; 1 oz. \$3.00; 1/4 lb. \$10.00

VINCA OR PERIWINKLE

A pretty ornamental plant, with dark green foliage and dainty, pink and white flowers. Useful for bedding, edging, and massing. Sow where the plants are to grow, and thin to about a foot apart each way. Does well in partial shade, and grows well on poor, sandy soil around beaches, where it is difficult to get other flowering plants to grow.

Pkt. 15c; 1/16 oz. 30c; 1/8 oz. 55c; 1/4 oz. 90c ½ oz. \$1.50; 1 oz. \$2.50; ¼ lb. \$8.00

ZINNIA

The zinnia is one of the most easily grown flowers for Florida, thriving in almost any soil and under almost all conditions.

While seedling plants of zinnias can be transplanted, they do best when the seed is sown directly in the garden where the plants are to flower. Seeds should not be sown, however, until the soil has warmed up and danger of low temperatures is past. For best results, provide a deep, rich soil and give plenty of water when needed especially in blooming time. For the best development of plants, they should be given ample room to allow free branching for sturdy plant development, which will assure the production of large, well formed, double flowers. Successive sowing can be made all through the summer and as late as the end of August in Florida.

Zinnias are practically free from all plant diseases and insects. Late in their growth, however, they may be subject to powdery mildew, but if the plants are properly spaced so that a free circulation of air is possible, this trouble should not be serious.

Pompon or Lilliput

Plants 11/2 feet high, dwarf, bushy and branching, bearing miniature flowers resembling pompon or button chrysanthemums. Charming for bedding. The small (1 to 1½ inches in diameter) double flowers are borne in great profusion, are small, compact, and symmetrical in shape. Excellent for cut flowers. Contains a large assortment of colors.

Pkt. 15c; 1/8 oz. 35c; 1/4 oz. 60c; 1/2 oz. \$1.00 1 oz. \$1.50; 1/4 lb. \$5.00; 1 lb. \$18.00

Peppermint Stick

Medium sized flowers with striped petals, red and white, red and yellow, pink and white, orange and yellow, purple and white. About one-third solid colors. An early blooming variety.

Pkt. 15c; ½ oz. 35c; ¼ oz. 60c; ½ oz. \$1.00

1 oz. \$1.50; 1/4 lb. \$5.00

Persian Carpet

Early and easy to grow, heat-loving and colorful. Miniature, fully double flowers of well rounded form or ball-shaped in a sparkling range of variegated colors. Resembles tiny dahlias. Plants dwarf and compact, in a nice color range, thrive on heat and yield a wealth of blooms on 6 to 8" stems.

Pki. 15c; 1/16 oz. 35c; 1/8 oz. 60c; 1/4 oz. \$1.00; 1/2 oz. \$1.75; 1 oz. \$3.00

Ortho Polka

A strikingly different Zinnia, similar to Peppermint Stick, but producing blooms almost double the size of the Peppermint Stick. The plants are large and vigorous, approximately 2½ feet tall. Colors are bright and varied, 70% of the blooms being striped or bi-colored, others solidly colored flowers of each hue.

Pkt. 15c; 1/16 oz. 45c; 1/8 oz. 75c; 1/4 oz. \$1.30

1/2 oz. \$2.25; 1 oz. \$3.75

ZINNIA—(Continued)

Giant Cactus Flowered

Blaze—Mandarin-red color, which changes to a deep warm orange as the flowers attain size. Large shaggy flowers 4 to 5 inches across, uniformly-double with graceful quilled and twisted flat petals. A strong grower, free flowering, uniform plants. A delightful cut flower.

Pkt. 15c; 1/16 oz. 35c; ½8 oz. 60c; ¼ oz. \$1.00 ½ oz. \$1.75; 1 oz. \$3.00; ¼ lb. \$10.00

Fantasy

The flowers of this group are double, of medium size, with shaggy petals that are tubular, curved or crested. Different from other Zinnias. An unusual range of colors.

Fantasy Mixture

Pkt. 15c; ½ oz. 35c; ¼ oz. 60c; ½ oz. \$1.00 1 oz. \$1.50; ¼ lb. \$5.00


Giant Hybrid Mixture—Plants are strong growing, free flowering, with long stems and good keeping quality for cut flowers. Flowers are very large with a variation in petal formation, mostly fluffy or ruffled, curled and quilled, free of the stiffness often associated with other zinnias. Large range of colors and color combinations.

Pkt. 15c; ½ oz. 35c; ¼ oz. 60c; ½ oz. \$1.00 1 oz. \$1.50; ¼ lb. \$5.00; 1 lb. \$18.00

Crown O' Gold

The petals of all the flowers are overlaid with a deep golden yellow at the base, with the individual flower color at the tips. Colors include various shades of salmon, peach and buff.

Pkt. 15c; ½ oz. 35c; ¼ oz. 60c; ½ oz. \$1.00 1 oz. \$1.50; ¼ lb. \$5.00; 1 lb. \$18.00



Zinnia-Special Crown O' Gold Mixture

Giant Dahlia-Flowered

The large flowers are fully double, with broad, undulating rays in magnificent solid colors. The plants grow 2 to 3 feet high.

Crimson Monarch—Crimson or red.

Wonder Rose—Light rose or pink.

Polar Bear-A large-flowered white Zinnia.

Price of each above
Pkt. 15c; ½ oz. 35c; ½ oz. 60c; ½ oz. \$1.00
1 oz. \$1.50; ¼ lb. \$5.00; 1 lb. \$18.00

Sunburst Mixture—Mixture of the finest colors.

Pkt. 15c; ½ oz. 35c; ¼ oz. 60c; ½ oz. \$1.00

1 oz. \$1.50; ¼ lb. \$5.00; 1 lb. \$18.00

Giants of California

Large double flowers of a distinctive graceful shape. The flowers are usually brighter than the Dahlia-flowered type and nearly as large, but more flat. Plants are vigorous, well branched, grow about 2½ feet tall and produce many long, strong flower stems each crowned with a beautiful flower. Perfect blend of color.

Giants of California Mixture-

Pkt. 15c; ½ oz. 35c; ¼ oz. 60c; ½ oz. \$1.00 1 oz. \$1.50; ¼ lb. \$5.00; 1 lb. \$18.00

GLADIOLUS BULBS

Rainbow Mixed

Enjoy beautiful cut flowers from your own garden. For continuous blooms plant bulbs every 10 days to two weeks. In North and Central Florida plant from August to October, and January to May. In South Florida plant from September to May. Set bulbs 4 inches deep and 6 inches apart in well-prepared and well drained beds.

Disease-free, well cured, graded extra large bulbs of good color assortment. We have exerted special effort, time and expense to furnish you these extra fine bulbs. No finer "Glad" bulbs for Florida can be secured any where at any price.

Gladiolus bulbs (2-21/2 inches).

1	dozen	bulbs (Exhi	ibition	Size)\$.85	postpaid
25	bulbs	(Exhibition	Size)		1.50	"
100	bulbs	(Exhibition	Size)		4.00	"
1000	bulbs	(Exhibition	Size)		35.00	**

STRAWBERRY PLANTS

All prices quoted are subject to change without notice.

We supply Maryland and Tennessee grown strawberry stock plants of the Missionary and Florida 90 variety from nurseries in those states. These northern grown stock plants are set out in multiplication beds in Florida from January through March. The young plants are usually set in the permanent field from September through November or early December, depending on weather conditions. The northern grown plants are usually available about November 1st and, when Florida grown plants are not available, are used by commercial growers for November setting and are also used to a great extent for home gardens. Care should be exercised never to allow the roots to dry out.

Strawberry plants may be set in rows three feet apart for single rows, or 4 to 5 feet apart for double rows, allowing 16 inches between double rows on each bed with 12 to 14 inches between plants in the row.

In setting strawberry plants, make holes deep enough to set the roots straight. Plants should be set so that the surface soil comes just to the crown.

Soil should be well firmed around the roots, and during dry weather each plant should be watered daily for several days, preferably in the late afternoon.

If the land is sour or acid, apply lime or hardwood ashes before setting plants. For control of insects and diseases see page 69.


See Fumigants on page 76 for Nematode (root knot) control.

After the plants are well established, 3 or 4 applications of high grade commercial fertilizer, such as 4-7-5 mixture, should be given the plants at about monthly intervals, being sure to make one heavy application of a 4-8-8 mixture when the plants begin to fruit.

MISSIONARY AND FLORIDA 90 (VIRUS-FREE)

		I	100 plants post	250 plants paid	500 plants F.O.B.	1,000 plants Nursery
MARYLAND	PLANTS		\$2.25	\$4.00	\$5.25	\$8.50
TENNESSEE			2.00	4.00	5.50	8.50
ARKANSAS	PLANTS	****	2.50	3.75	4.25	7.50





Beautiful lawns can be grown in Florida to enhance the appearance of your Florida home.

Lawn and Golf Course Grasses

All grass seed prices quoted are f.o.b. Plant City except lb. lots which are postpaid.

Write for prices on larger quantities.

Kilgore's Lawn Grasses are especially adapted to Florida conditions.

All grass seed prices quoted are subject to change without notice.

Bitter Blue-Stemmed St. Augustine and Centipede Grass are used only for lawns. The other grasses listed and described on this page are used for lawns, golf courses and for pastures. For information on "When to plant grass seeds in Florida" see page 6. For lawn and golf course fertilizers see page 79. For lawn grass pesticides and supplies see pages 67 and 81.

BERMUDA GRASS

Unhulled—For lawns sow 2 to 4 pounds per 1000 square feet. For pasture, sow 25 pounds per acre.

Hulled—For lawns sow 1½ to 2 pounds per 1000 square feet. For pastures, sow 15 to 25 pounds per acre.

Bermuda grass is of fine texture, can be maintained green practically the year around, is a rapid grower and produces a lush, green carpet of turf though it requires careful attention in maintenance. Frost will turn it brown for a short period, but it soon regains its green color. It prefers loam or clay soil but will grow well on sand or muck. 20 to 30 pounds per 1000 square feet of a complete fertilizer such as Vigoro or Vertagreen should be applied twice a year. Frequent weeding is desirable. A thorough watering every 5 to 7 days during periods of drouth should be sufficient. The grass may be clipped closely (34 to 1 inch) and frequently. Bermuda will not grow in shade. It spreads by surface runners and by underground root stalks. Seed can be planted from March to November.

Unhulled Bermuda Grass

1 lb. 70c; 10 lbs. \$3.80; 25 lbs. \$9.00; 100 lbs. \$35.00

Hulled Bermuda Grass

The almost watertight husks of the seed have been removed, facilitating quicker, surer results, the seed germinating more uniformly and in a much shorter time with proper moisture conditions.

1 lb. 85c; 10 lbs. \$5.60; 25 lbs. \$13.50; 100 lbs. \$53.00

BAHIA GRASS

For Pasture See Page 49. For lawns, sow 100 to 150 pounds per acre or 5 to 6 pounds per 1000 square feet.

Pensacola Bahia Grass

This variety of grass is sometimes used for low maintenance lawns. It is of fine texture and will remain green most of the winter (longer than Centipede grass). Will tolerate dry or infertile soils. It is somewhat difficult to mow, especially when rapid growing seed shoots develop during the summer, but should not be mowed closer than 1½ to 2 inches. It requires fertilization in mid-spring and late summer, with a complete fertilizer such as Vigoro or Vertagreen. It is susceptible to damage from salt spray, but is relatively free of insect and disease troubles. Spring and early summer are the best periods for planting. Seeds are somewhat slow to germinate, sometimes requiring several weeks. The seedbed should be kept moist until germination occurs. Sow seed at the rate of 2 to 3 pounds per 1000 square feet. "Common" Bahia should never be used for lawns.

1 lb. 70c; 10 lbs. \$4.30; 25 lbs. \$10.25; 100 lbs. \$40.00 Paraguayan Bahia Grass. (Texas grown turf type) is the newest and most highly recommended of all the Bahias—a short narrow leaf perennial making a dense sod.

1 lb. \$1.35; 10 lbs. \$10.50; 25 lbs. \$25.25 100 lbs. \$100.00

Argentine Bahia Grass

Makes a semi-erect grass generally more vigorous than other Bahias with leaves wider than Pensacola but slightly narrower than Common.

1 lb. 85c; 10 lbs. \$5.30 25 lbs. \$12.75; 100 lbs. \$50.00


DICHONDRA

For lawns sow 1 pound per 500 square feet.

Dichondra is one of the few ground cover plant materials that are hardy throughout Florida and may be grown with equal success throughout the state. Since the height of this plant is seldom more than 1 or 2 inches, there is no need of mowing or shearing. It also may be tramped as well as any of the lawn grasses.

 $\frac{1}{2}$ oz. 75c; oz. \$1.25; $\frac{1}{4}$ lb. \$3.75; 1 lb. \$6.75 (Continued on next page)





COMMON RYE GRASS

For Lawns Sow 350 to 400 Pounds Per Acre. Or 10 Pounds Per 1000 Square Feet. For Temporary Winter Pasture Sow 50 Pounds Per Acre.

This is valuable and desirable fall and winter grass for Florida, but it does not make a permanent lawn. It dies out in warm weather or late spring and early summer. For quick covering of lawns and golf courses in winter it is unsurpassed. Do not plant in Florida before October or later than March. It will make a vigorous, luxuriant growth throughout the fall, winter and early spring but must have plenty of water. A common practice in Florida is to seed Italian Rye grass on other brown lawns or in brown unsightly spots of old lawns in the fall and winter for a velvety green appearance. Plant ½ lb. to 100 square feet right on top of old lawn after loosening the surface. Keep soil moist until seed sprouts.

1 lb. 50c; 10 lbs. \$1.75; 25 lbs. \$3.75; 100 lbs. \$14.00

BITTER BLUE-STEMMED ST. AUGUSTINE GRASS

Use one bushel of sprigs per 200 to 300 square feet.

This is one of the most popular lawn grasses for Florida. It is quick to establish, rather coarse in texture, but easy to mow, and remains green the year around. It should be cut at a height of 2 inches. Severe cutting back is quite harmful. It should be fertilized two or three times a year at the rate of 20 pounds per 1000 square feet with a complete fertilizer such as Vigoro or Vertagreen. St. Augustine is our best shade grass and will grow on alkaline as well as slightly acid soils. It is very salt tolerant and can be grown along the beaches where a good soil base is provided. It can be grown by vegetative propagation only. Sprigs should be planted in well prepared soil, setting about 6 X 9 inches apart. For chinch bug control in St. Augustine lawns see page 57.

1 lb. \$2.50; 5 bu. \$2.40 per bu.; 10 bu. \$2.25 per bu.

Not prepaid. Available April to November. Shipped only by express direct from a Florida nursery.

For further information on lawns, write the Florida Agricultural Experiment Station, Mailing Department, Gainesville, Florida, for a free copy of Bulletin 518 entitled "Lawns in Florida" and for Circular S-96 entitled "Insects and other Pests of Lawns and Turf."

CARPET GRASS

For Lawns Sow 100 to 150 Pounds Per Acre or 2 to 4 Pounds per 1000 Square Feet. For Pastures 25 Pounds Per Acre.

Carpet grass is particularly well adapted to low moist soils and does not require much fertilization, but needs plenty of moisture. This grass is not suited to drouthy conditions or to sweet soils with a high pH. It burns brown in the winter, is not at all salt tolerant nor will it stand much shade. Keep mowed to a height of 2 inches. Plant in the spring from March to July, or in fall from October to December.

1 lb. 85c; 10 lbs. \$5.00; 25 lbs. \$12.00; 100 lbs. \$47.00

CENTIPEDE GRASS

Sow 4 to 6 ozs. of Seed per 1000 Square Feet. Use one bushel of sprigs per 300 to 400 square feet.

This is considered the best low maintenance grass for Florida. It is adapted to a wide range of soils including high sand hills, and will tolerate infertile acid soils, can get by on a minimum of fertilizer and water, and is relatively free of insect pests and diseases. A light spring and/or fall application—10 to 15 pounds per 1000 square feet, should be sufficient for best results. Do not over-fertilize. The fertilizer should be watered in. In periods of extended drouth a thorough watering every 5 to 7 days should be sufficient, wetting the soil 4 to 6 inches deep each time. Grass seed germinates in a period from 2 to 7 months and is easy to mow to a height of 1½ inches. However, it turns brown during the winter, does not tolerate shade and salt spray as well as St. Augustine, and does not grow well in alkaline soils.

Centipede Grass Seed

1 oz. \$1.00; ½ lb. \$3.95; ½ lb. \$7.75; l lb. \$14.95

Postpaid

Centipede Grass Sprigs or Roots

Centipede sprigs or root runners 3 to 5 inches long should be planted in rows 10 to 12 inches apart, 4 to 5 inches apart in the rows. Insert sprigs in the ground, leaving about an inch showing. Give frequent watering until plants are rooted firmly.

1 lb. \$2.25; 5 bu. \$2.15 per bu.; 10 bu. \$2.00 per bu.

Not prepaid. Available April to November. Shipped only by express direct from a Florida nursery.

Write Mail Order Department, Kilgore Seed Co., Plant City, Fla., for instructive folder on "Centipede Grass Lawns."

PASTURE GRASSES

All prices quoted are f.o.b. Plant City except 1 lb. and 2 lb. lots which are prepaid.

Write for prices on larger quantities.

All prices quoted are subject to change without notice.

To get better growth and increase soil nitrogen, it is essential to inoculate seed of all leguminous crops—clovers, cowpeas, crotalaria, hairy Indigo, lupines, peanuts, sesbania, etc. (See page 82.) Bermuda, Carpet and Italian Rye Grass are used for pasture as well as lawns. (See pages 51 and 52.)

BAHIA GRASS FOR PASTURES

Sow 10 to 15 pounds broadcast per acre.

Bahia grass is established with seed planted ¼ to ½ inch in depth, during any month of the year in Florida. In order to insure better germination all Bahia grass seed is scarified before delivery to our customers. Widely adapted. Will grow in dry soils because of deep root system. Makes one of Florida's best permanent pasture grasses.

Common Bahia. (Scarified.) A low growing perennial, spreading by short, stout, woody runners which root heavily with large fibrous roots forming tough sod, even on drouthy, sandy soils. Leaf blades shorter and coaser than Pensacola but usually less than ½ inch wide.

1 lb. 50c; 10 lbs. \$2.30; 25 lbs. \$5.75; 100 lbs. \$22.00

Pensacola Bahia. (Scarified.) Narrow leaf, tall, erect-growing strain of Bahia grass, very hardy, claimed to stand cold weather conditions better than Common. Pensacola variety seed usually germinates quicker and stronger than Common. Succeeds on wide range of soils. Has heavy root system and withstands drouth better than other pasture grasses. For best results fertilize with 500 lbs. per acre of a complete fertilizer. Prevents erosion and provides very palatable pasturage. Withstands heavy grazing and is very resistant to trampling by livestock.

1 lb. 70c; 10 lbs. \$4.30; 25 lbs. \$10.25; 100 lbs. \$40.00

Argentine Bahia

Makes a semi-erect grass generally more vigorous than other Bahias, with leaves wider than Pensacola but slightly narrower than Common. Adapted to all kinds of soils, from high dry sand to low wet areas. Just as frost-resistant as Pensacola. For best results sow seed in late fall or early winter one-half to one-inch deep.

1 lb. 85c; 10 lbs. \$5.30; 25 lbs. \$12.75; 100 lbs. \$50.00

SWEET SUDAN GRASS

Sow 10 pounds in drills or 20 pounds broadcast per acre.

Resembles common Sudan in plant habit and yield, but is sweet and juicy, more productive and resistant to several foliage diseases, which attack common Sudan. Grazing tests have indicated that cattle prefer Sweet Sudan to common Sudan. Sow from March to Sept.

1 lb. 50c; 10 lbs. \$1.50; 25 lbs. \$3.25; 100 lbs. \$12.00

BEGGARWEED (Florida Clover)

Sow 10 pounds per acre

Perfectly adapted to Florida sandy soil. Makes fine pasture, nutritious hay. For best quality hay, cut crop when 3 to 4 feet high, or at beginning of blooming period. Sow from April to August. Cover seed lightly, inoculate seed with Nitragin. (See page 82.)

1 lb. \$1.10; 10 lbs. \$7.80; 25 lbs. \$19.00; 100 lbs. \$75.00

ALFALFA-HAIRY PERUVIAN

Sow 20 pounds per acre.

A valuable forage and hay crop for Florida. Does well in central and south Florida where it is used extensively by dairy and cattle farmers. Does best on moist, dark sandy soils (not subject to flooding). Also does well on muck. Soil should be tested before starting seedbed and sufficient limestone applied several months in advance of seeding. The pH should be brought up to 6.5 or 6.8. Extra lime can be applied after the seedbed has been prepared if pH is too low. After plowing, firm seedbed by disking and cultivating. Broadcast, 1000 pounds per acre of fertilizer high in phosphoric acid and potash blus 20 pounds of borax ahead of seeding. Inoculate seed with double recommended rate of inoculant. Use cultipacker type of seeder, and seed from September to January at rate of 20 pounds per acre, when soil is moist. After seeding apply 500 pounds of a phosphoric acid-potash fertilizer broadcast in 60 to 90 days. (See page 82 for inoculants.)


1 lb. 70c; 10 lbs. \$3.90; 25 lbs. \$9.25; 100 lbs. \$36.00

ALYCE CLOVER

Sow 10 to 15 pounds per acre

Used as cover crop, soil builder, for hay, and as pasture plant. Best time for planting is during spring. Must be planted on well drained soil free of contamination with root knot. Seed should be planted on well prepared ground and covered very lightly. Pack soil after seeding. Makes palatable hay. Cut for hay when it begins to flower. A good soil builder. Most economical fertilizer application is 200 to 400 pounds per acre of 0-16-16 or 0-14-14 mixture. Seed should be inoculated. (See page 82.)

1 lb. 50c; 10 lbs. \$1.80; 25 lbs. \$4.00; 100 lbs. \$15.00



White Clover Pasture

Photo Courtesy of Florida Experiment Station

CLOVERS

All prices quoted are subject to change without notice.

Prices quoted are f.o.b. Plant City except 2 lb. lots or less which are postpaid.

Write for prices on larger quantities, indicating variety and quantity desired.

All clovers should be inoculated. (See page 82.)

ANNUAL SWEET CLOVER

Sow 8 to 12 pounds per acre of inoculated seed alone, or 3 to 5 pounds per acre in mixed clover plantings.

Sow in Florida from Sept. through Jan., although from Oct. to Nov. 15 sowing is best. Makes a rapid growth, and widely adapted, growing best on moist soils. Does not tolerate very close grazing. Grazing reaches peak during March and April. Liming soil heavily and applying 400 to 600 pounds of a fertilizer high in Phosphoric Acid and Potash at planting time is recommended.

HUBAM. (Sweet). A most valuable legume crop for grazing and soil improvement. It makes rapid and luxuriant growth and can be used as a winter cover crop. Excellent for silage and desirable for bees.

1 lb. 60c; 10 lbs. \$2.50; 25 lbs. \$5.75; 100 lbs. \$22.00

Floranna (Sweet). This variety developed by the Florida Agricultural Experiment Station from hardy strains found volunteering is best adapted to Florida soil and climatic conditions. It outyields and resists drouth and cold weather better than other varieties.

Write for Prices

WHITE CLOVER (Louisiana grown)

Formerly listed as "White Dutch"
Sow 4 to 6 pounds per acre alone, or 2 to 3 pounds per

Recommended principally for mixture in perennial pastures. Succeeds best on moist ground and during wet seasons. Sow in the late fall and early winter. Fertilize with 400 to 500 lbs. per acre of 0-14-10 mixture. Furnishes abundant grazing during most of the year.

1 lb. \$1.50; 10 lbs. \$11.30; 25 lbs. \$27.75; 100 lbs. \$110.00

LADINO CLOVER (Certified)

This is a large form of White Clover. It grows rapidly and spreads by creeping stems that root at the nodes. Sow in late fall or early winter. Used for soil improvement, pasture and hay. Does best on moisture retaining soils.

1 lb. \$1.15; 10 lbs. \$8.30; 25 lbs. \$20.25; 100 lbs. \$80.00

KENLAND RED CLOVER (Certified)

Sow 4 to 6 pounds per acre.

This clover is rapidly gaining in popularity in Florida as a luxuriant hay and pasture crop for cattle and hogs. It has a longer active growing season than some other legumes. Fertilize with 400 to 600 lbs. per acre of a 0-14-10 or 0-10-10 fertilizer mixture just before planting time. Seed from September to December. Grows well with Bermuda and Bahia grass. For hay, cut when in full bloom. It makes good growth in April, May and June when White Dutch and Crimson clover don't do so well.

1 lb. 90c; 10 lbs. \$5.80; 25 lbs. \$14.00; 100 lbs. \$55.00

COWPEAS

CULTURE

Table varieties are sown in drills three to four feet apart. Fertilize up to 300 lbs. per acre. Inoculate seed with Nitragin. (See page 82.) For insect control see page 66.

TABLE VARIETIES

Sow 20 to 45 pounds per acre in drills with table varieties. In the garden $\frac{1}{2}$ to $\frac{3}{4}$ pounds to 150 feet row.

TEXAS CREAM 40 (Improved Bush Conch). (62 days.) This variety of Conch pea is very early. The vigorous upright growing plants make a bush about 24 inches tall. Pods are oval, medium green, straight, and closely filled with light green, tender peas. Seeds white.

1/4 lb. 20c; 1 lb. 60c; 2 lbs. \$1.10 15 lbs. (pk.) \$4.25; 60 lbs. (bu.) \$16.00

ACRE (Running Conch). (80 days.) This variety makes a long spreading vine, matures relatively early for a running variety, and has an exceedingly long bearing period, bearing an immense crop. It does well for late planting, and makes a good, late summer crop.

1/4 lb. 15c; 1 lb. 35c; 2 lbs. 60c; postpaid Write for prices on larger quantities.

CALIFORNIA BLACKEYE NO. 5 (Early Wilt-Resistant Ramshorn). (65 days.) It is resistant to nematodes (root knot) and cowpea wilt. Very early and prolific, making a great number of 10 to 12 inch long pods on an upright vine. Uniform in size, growth and maturity, and flavor.

1/4 lb. 20c; 1 lb. 55c; 2 lbs. \$1.00 15 lbs. (pk.) \$3.15; 60 lbs. (bu.) \$12.00

CALIFORNIA BLACKEYE No. 7 (Giant Wilt-Resistant Ramshorn). (70 days.) A vigorous heavy yielding pea. About 10 days later than the Early Wilt-Resistant Ramshorn Blackeye. The peas are much larger than the regular Blackeyes and can be used both green and dried.

1/4 lb. 20c; 1 lb. 55c; 2 lbs. \$1.00 15 lbs. (pk.) \$3.15; 60 lbs. (bu.) \$12.00

ALABAMBA CROWDER. (70 days.) The bush is of the semi-runner type. The large, straight pods average about 8 to 9 inches in length and are of very attractive appearance. A sure cropper, and a heavy yielder.

1/4 lb. 20c; 1 lb. 60c; 2 lbs. \$1.10 15 lbs. (pk.) \$4.25; 60 lbs. (bu.) \$16.00

PURPLE HULL NO. 49. (65 days.) This variety was introduced by the Texas Agricultural Experiment Station. It combines purple pod color, buffeye seed and high production with earliness and erect, bunch-type growth. Pods are long, well filled with large peas. Excellent eating quality.

1/4 lb. 20c; 1 lb. 65c; 2 lbs. \$1.15; 15 lbs. (pk.) \$4.75 60 lbs. (bu.) \$18.00

FIELD VARIETIES

Sow 1 to $1\frac{1}{2}$ bushels per acre broadcast with field varieties. A bushel weighs 60 pounds.

BRABHAM. Is resistant to wilt and root knot, makes fine, vigorous growth of vines, is early, and when planted in March, April or May good cuttings of hay can be made. Also a very good soil builder.

Write for Prices

COMBINE. Excellent for soil improvement, hay, and wild bird feed. Seed very small.

Write for Prices

MIXED. Mixed Cowpeas produce an exceedingly heavy growth of vines for forage or soil improvement.

Write for Prices

FIELD SEED

All prices quoted are subject to change without notice and are f.o.b. Plant City except 2 lb. lots or less which are postpaid. Write for prices on larger quantities.

For information on "When to Plant Field Seed in Florida" see page 6.

BENE (SESAME)

Sow 3 to 5 pounds per acre drilled.

Produces flowers, followed by seed pods, which scatter oily seeds in profusion, attracting quail and other game birds. Valuable for poultry feed. Plant from March to June, drilling seed thinly in rows 3 to 4 feet apart. We have only the native giant or tall variety growing to a height of from 4 to 6 feet, and rather spreading.

¹/₄ lb. 25c; 1 lb. 80c; 10 lbs. \$4.30; 25 lbs. \$10.25 100 lbs. \$40.00

CHUFAS

Plant 1 to 1½ pecks (10 to 15 lbs.) per acre.

(90 days.) This is a species of ground nut, easily grown, a useful hog fattening crop. Plant from April to August. Cultivation same as for bunch peanuts. Can be left in the ground until time to turn in hogs who do harvesting. On sandy loam soil best adapted for Chufas, plant in 2-ft. rows with plants 12 inches in the row, or in 3-ft. rows with plants 6 inches in the row.

1 lb. 50c; 2 lbs. 90c; 10 lbs. (pk.) \$2.25 40 lbs. (bu.) \$8.00

CROTALARIA

Sow 10 to 20 pounds of seed per acre.

Crotalaria is particularly desirable as a cover crop for groves, truck and farm lands, does well on all soils except muck. Is a legume, thus adding a large amount of nitrogen to soil. Is especially valuable because it is immune to nematodes causing root knot disease. Furnishing abundance of green material for humus. Sow broadcast March to June, on well prepared land. Use an inoculant. (See page 82.)

EARLY SPECTABILIS. This vigorous growing, long leaved variety is not so fibrous as others, the growth is much heavier, and the stems are hollow. As a result it rots much quicker when plowed under. Spectabilis does well on high ground, but is best adapted to the better grades of sandy soils. Leaves, stems and seed of this Cortalaria plant, green or dry, are poisonous to poultry and livestock.

1 lb. 60c; 10 lbs. \$2.80; 25 lbs. \$6.50; 100 lbs. \$25.00



Crotalaria—Early Spectabilis

HEGARI

See Sorghum, page 58

INDIGO

(Indigofera hirsuta)

Sow three to five pounds per acre in 30 inch drills, 6 to 10 pounds broadcast.

A three purpose legume, used as a summer cover crop, for hay and for grazing. Used extensively as a cover crop by citrus growers in south Florida, as a grazing crop or for hay by cattlemen and as a cover crop by general farmers. Its feeding value compares fairly well with alfalfa, making a palatable and nutritious hay. It makes an ideal summer cover crop for citrus groves, harboring no troublesome insects, and reseeds itself year after year. It's very resistant to drouth. The plants grow four to seven feet in height, and produce an abundance of hay and grazing. If grazed or harvested for hay before the stems become woody a second or third growth may be expected.

Seed may be planted not over an inch deep, from the middle of March to the later part of May, but early seedlings are preferable. Heavier seedings are desired for green manure or forage, lighter seeding for seed. On poor soil 300 to 500 lbs. per acre before seeding of a 0-10-10 or 0-14-10 fertilizer mixture is desirable. It will grow on a wide range of soils, but well drained sandy loam soils are best. It is sensitive to cold, is killed by first hard frost and is best suited to the section of Florida from Gainesville south, although the early strain can be grown successfully in north Florida and as far north as central Georgia. Highly resistant to root knot, but in general it is reasonably free from disease and insect injury. Use an inoculant. (See page 82.)

COMMON INDIGO

Write for Prices

EARLY INDIGO

Write for Prices

LUPINES

60 to 80 pounds with yellow and 70 to 90 pounds with blue per acre broadcast, or 50 to 60 pounds per acre in drills with a grain drill. Inoculate the seed. (See page 82.)

Lupines are adapted to north and central Florida as a winter legume cover crop. They are not affected by cold and because of their heavy yield make an ideal green manure crop. Plant from late September to December, preferably in late September or October, in well prepared soil, using between 250 to 300 lbs. of Superphosphate per acre.

BITTER BLUE LUPINE. This variety has rapidly grown in popularity as a soil builder. Normal growth is about waist high and normal weight of green matter 20-30 tons per acre. It produces about three times as much growth as Winter Peas, Hairy Vetch, etc. It protects the soil from erosion and restores humus and plant food removed by peanuts and other crops. Peanut and corn yields following Bitter Blue Lupine have been greatly increased. A good winter cover or green manure crop, but poisonous to livestock.

1 lb. 40c; 10 lbs. 95c; 25 lbs. \$1.85; 100 lbs. \$6.50

BORRE SWEET (Blue). This variety is the pasture and feed counterpart of Bitter Blue Lupine. Both plant and leaf are sweet to the taste and are relished by all kinds of livestock. Makes about the same growth and tonnage as Bitter Blue Lupine. Stock can be turned in on Sweet Blue Lupine as soon as it is about a foot high. The plants have a high protein value averaging approximately 18% of the dry matter.

1 lb. 50c; 10 lbs. \$1.20; 25 lbs. \$2.50; 100 lbs. \$9.00

(Continued on next page)

All prices quoted are subject to change without notice and are f.o.b. Plant City except 2 lb. lots or less which are postpaid. Write for prices on larger quantities.

LUPINES (Continued)

SWEET YELLOW LUPINE. (White seeded.) More readily grazed by cattle than the Sweet Blue strains, and will recover more rapidly after grazing. More tender and less resistant to excessive cold than the Blue, and thus the Yellow Lupine is more particularly adapted to the southern portion of the Lupine belt than to the more northern areas.

1 lb. 55c; 10 lbs. \$2.00; 25 lbs. \$4.50; 100 lbs. \$17.00

MILLET

Sow 6 to 8 pounds per acre in rows 2 feet apart. Sow 20 to 25 pounds per acre broadcast for pasture.

CATTAIL OR PEARL. This is one of the best yielders of green forage and hay, continuing to grow and produce through the entire season if cut frequently enough to prevent its going to seed. Relished by all kinds of stock. Plant from March until September.

1 lb. 45c; 10 lbs. \$1.30; 25 lbs. \$2.75; 100 lbs. \$10.00

STARR (Certified)

A new variety of millet bred to last longer and be easier to manage. It appears to be the best of all summer temporary grazing crops for dairy cattle, beef cattle, and hogs on the lighter soils. Leafier than common millet and produces more tonnage. Apply up to 100 pounds of nitrogen per acre at planting time if soil fertility is low.

1 lb. 50c; 10 lbs. \$1.70; 25 lbs. \$3.75; 100 lbs. \$14.00

OATS

Sow 5 pecks (40 lbs.) per acre in drills; broadcast 2 to $2\frac{1}{2}$ bu. per acre. A bushel weighs 32 pounds.

Plant in drills 1½ feet apart and cover seed 2 to 3 inches deep. If the seed is broadcast, use tooth harrow. The best time to plant oats in Florida is October and November, but can be planted to January. Land should receive 300 to 400 lbs. per acre of complete fertilizer at planting, followed by a side-dressing of 50 to 200 lbs. Nitrate of Soda or its equivalent after the crop is half grown.

VICTORGRAIN 48-93. (120 days.) An early high yielding cold resistant variety, with strong, stiff straws that will withstand wind and storms. Resistant to most races of smut and rust, also tolerant to strains of blight. Victorgrain 48-93 is an improvement over the original strain and is highly recommended for Florida.


1 lb. 35c; 2 lbs. 50c; 8 lbs. (pk.) 75c; 32 lbs. (bu.) \$2.00



Victorgrain 48-93

SEMINOLE (new) (Disease resistant). (110 days.) This new, early maturing oat has been increased on Experiment Station Farms in Florida, and seed is now available. It is resistant to different races of Crown rust and to smut. Due to its earliness it escapes damage from stem rust. A short variety with strong straw and not subject to lodging. The grayish-white grain is larger, plumper and of higher quality than Floriland. Makes a very early, upright growth of medium large leaves. Produces an abundance of early forage and is good for early and mid-winter forage. Also excellent in grain yield. Because of its earliness it is well adapted to central and south Florida.

1 lb. 35c; 2 lbs. 50c; 8 lbs. (pk.) 75c; 32 lbs. (bu.) \$2.00



Seminole

FLORILAND. (115 days.) Floriland has an intermediate to upright habit of growth. Floriland is early, heading approximately 10 days earlier than Southland. At maturity the variety is slightly taller and more uniform in appearance than Southland. The kernels are red, midsize, midplump and midlong. They have numerous non-twisted awns and abundant basal hairs of medium length. Floriland is highly resistant to prevalent races of crown rust and victoria blight.

1 lb. 40c; 2 lbs. 55c; 8 lbs. (pk.) 85c; 32 lbs. (bu.) \$2.25

PEANUTS

Plant 1 bushel of seed per acre. A bushel weighs 25 lbs., except Valencia 22 lbs. per bu.

In addition to the crop of nuts for hogs, peanuts yield a large quantity of nutritious dry forage. Dixie Runner peanuts are by far the best when grown to fatten hogs, as they can be left in the ground until eaten.

It is better to shell the seed that is to be planted. If planted in corn, a bushel will plant 4 acres, but where planted solid it requires a bushel of seed per acre, in 3-foot rows, dropping seeds in hills a foot apart. Plant from March to July. For best results inoculate peanuts with Nitragin. (See page 82.)

Most shelled peanuts are treated so they will keep better and longer, and also to protect them from rot and disease after they are planted and to increase the stand. Do not use Nitragin inoculation with treated shelled peanuts.

For Insect and Disease Control See Page 68.

VALENCIA. (100 days.) This variety is adapted for growing in light sandy soils. It fills the demand for a peanut that is large and contains more nuts than the common varieties. Pods are large, well filled, usually containing 3 to 4 nuts.

1 lb. 60c; 5½ lbs. (pk.) \$1.85; 22 lbs. (bu.) \$6.75

(Continued on next page)



All prices quoted are subject to change without notice and are f.o.b. Plant City except 2 lb. lots or less which are postpaid.

Write for prices on larger quantities.



Dixie Runner

bixie runner. (120 days.) This variety produces high yields like Spanish peanuts. Four tests averaged 53 per cent higher yields than the Florida Runner. Remains dormant after maturity like "hogging off." Is a vigorous grower with more prostrate branches. Kernels are like Florida Runner in size, shape, markings and color. Pods are somewhat lighter, shorter, thicker at the waist and more cylindrical in shape. In addition to high yield and dormancy characteristics, this variety possesses a valuable quality for shellers, shows less than 1 per cent concealed damage, or decay of the kernels.

1 lb. 50c; 61/2 lbs. (pk.) \$1.50; 25 lbs. (bu.) \$5.50

Shelled and treated Dixie Runners. One bushel of good Dixie Runners should yield about 15 lbs. shelled peanuts. Plant 25 lbs. per acre.

1 lb. 65c; 10 lbs. \$3.30; 25 lbs. \$7.75; 100 lbs. \$30.00

DWARF ESSEX RAPE

Sow 4 pounds in drills or 8 to 10 pounds broadcast per acre.

For early spring, fall or winter planting to give quick green forage or grazing for hogs and poultry. While mostly planted in fall and early winter, it may be planted in early spring, but it will not stand much warm weather. It makes a quick succulent growth. It can be planted in rows like turnips, or broadcast covering by a light harrowing. It is ready for grazing in eight to ten weeks after planting. After cutting, or grazing, it keeps coming out again. Rape can besown with fine results any month from September to March. It is a splendid green feed and forage crop relished by all kinds of stock and especially by milk cows.

Pkt. 10c; 1 lb. 55c; 10 lbs. \$2.30; 25 lbs. \$5.25 100 lbs. \$20.00

RICE

Sow 1/2 bushel (22 lbs.) per acre.

Rice makes fine feed for both stock and poultry. Prepare the ground thoroughly, drill ½ bushel per acre in 2½-foot rows and cultivate a few times. Should be planted in March or April so that it may mature at a time to miss the rice-bird season. If planted in June and July it will also miss the bird season.

BLUE BONNET (Gopher). A popular variety for Florida, generally known as Upland Rice, but will also do well on low land. Grains heavy, broad, and flat.

1 lb. 35c; 2 lbs. 60c; pk. \$1.60; bu. \$6.00

ZENITH (Pearl). Grown extensively in Florida. It produces a heavier, taller, more vigorous stalk with heavier heads than other varieties of this type. Does not shatter so badly as other varieties. Produces long, slender grains. Best adapted for low wet land.

1 lb. 35c; 2 lbs. 60c; pk. \$1.60; bu. \$6.00

RYE

Sow 3 pecks in drills; broadcast 1 to ¼ bushels per acre. A bushel weighs 56 pounds.

Rye can be planted from September to January but October, November and December are the best months in which to plant for seed. Earlier planting is recommended for winter and early spring grazing. Fertilization is the same as for oats. (See page 52.)

ABRUZZI. One of the best for Florida, producing a heavier tonnage of green feed and grain than most other varieties. Tall and upright, and a heavy yielder.

1 lb. 40c; 2 lbs. 75c; 14 lbs. (pk.) \$1.40 56 lbs. (bu.) \$5.00

FLORIDA BLACK. A standard, upright, tall growing variety. Northern and western varieties of rye have a tendency to fall over on the ground and are not suitable for grazing or cutting in Florida. Florida Black is upright in growth and is exceptionally well suited for early grazing, making pasturage quickly, and it withstands cold very well. Highly recommended by the Florida Agricultural Experiment Station.

1 lb. 40c; 2 lbs. 75c; 14 lbs. (pk.) \$1.40 56 lbs. (bu.) \$5.00



Florida Black



All prices quoted are subject to change without notice and are f.o.b. Plant City except 2 lb. lots or less which are postpaid. Write for prices on larger quantities.



Beefbuilder

SORGHUM

Sow 8-10 lbs. to 1 bushel (50 lbs.) per acre.

The uses of Sorghum are many; green feed, hay, pas-

turage, silage, syrup, etc.


It is common practice to sow Sorghum and Cowpeas together for hay and pasturage purposes using about ½ bushel Sorghum and 1 bushel of Cowpeas per acre, sown broadcast. For forage, use 1 bushel of seed per acre. For grain Sorghums apply 300 to 500 lbs. of a complete fertilizer mixture per acre. Plant from March to August.

BEEFBUILDER (For Forage). A vigorous, midseason hybrid developed by Asgrow for the grower who wants a maximum tonnage of high quality feed in a relatively short growing period, compared with late open pollinated varieties. Makes excellent recovery for second crop in Florida.

1 lb. 55c; 10 lbs. \$2.50; 25 lbs. \$5.75; 100 lbs. \$22.00

AMAK R-10 (For Grain). A widely adapted, early maturing Combine type developed by AMAK to produce a heavier yield of quick drying grain in areas where Martin, which it resembles, can be used. It has consistently outyielded Martin. Combine 7078 and other leading open pollinated varieties in trials conducted throughout the sorghum belt.

1 lb. 55c; 10 lbs. \$2.30; 25 lbs. \$5.25; 100 lbs. \$20.00



Amak R-10

HEGARI. Early Dwarf Type. (For grain and forage.) (100 days.) Grows 2 to 5 feet depending on type of soil, with large heads of white grains which do not shatter. Stands dry weather exceptionally well. Sow ten pounds per acre for grain, one bushel (50 lbs.) per acre broadcast for forage.

1 lb. 40c; 10 lbs. 95c; 25 lbs. \$1.90; 100 lbs. \$6.50

HEGARI. Regular Type. (For forage and cover crop.) (115 days.) This strain is slower to seed but makes more forage quicker than the Early Dwarf Type, and also is better for cover crop. Sow 30 to 40 lbs. per acre broadcast for forage or for a cover crop.

1 lb. 40c; 10 lbs. 95c; 25 lbs. \$1.90; 100 lbs. \$6.50

SILO KING (For Forage). A widely adapted, early maturing, vigorous hybrid developed to replace Atlas. Produces a heavier yield of sweet, high-nutrient, palatable feed for live stock. Makes excellent recovery in areas where a second crop can be produced. Tolerant of heat and drought.

1 lb. 55c; 10 lbs. 2.50; 25 lbs. \$5.75; 100 lbs. \$22.00



Silo King

TEXAS SEEDED RIBBON CANE. (For syrup, ensilage, forage and green feed.) The plants grow ten to twelve feet tall, and the thick stalks contain an abundance of juice and also produce plenty of leaves for fodder. Not only valuable for syrup-making, but will make a large amount of forage and green feed for livestock.

1 lb. 55c; 10 lbs. \$2.30; 25 lbs. \$5.25; 100 lbs. \$20.00

WINDBREAKER (For Windbreaks). A sterile hybrid. Excellent protection. No volunteers. Produces quick, vigorous, leafy growth on a sturdy stalk to height of 5-7 ft. Gives excellent wind protection for low-growing crops, good quality for ensilage. Windbreaker is sterile and will not make seed unless other sorghums are close enough to provide pollen. Can be left to full maturity and turned under for soil improvement, without danger of volunteer plants in next crop.

1 lb. 55c; 10 lbs. 2.50; 25 lbs. \$5.75; 100 lbs. \$22.00

SHALLU OR EGYPTIAN WHEAT

Sow 8 to 10 pounds per acre for grain. 25 to 30 pounds for cover crop.

This crop is being grown extensively and very successfully as a summer green manure or cover crop in many sections of Florida. Grows rapidly and makes a great amount of vegetable matter to plow under in a relatively short time. Not a legume. If allowed to go to seed for a grain crop like other sorghums, makes large, loose, bushy heads covered thickly with small grains. Grain is of high feeding value and can be fed to chickens, hogs and cattle as desired. For grain sow seed thinly in 3 to 4 foot rows, leaving 3 to 4 plants every 3 feet of row. Plant from March to August.

1 lb. 50c; 10 lbs. \$1.55; 25 lbs. \$3.40; 100 lbs. \$12.50

SESBANIA

Sow 30 to 40 pounds per acre.

Sesbania is a most valuable summer cover crop to grow on land that is more or less covered with water. Unlike other cover crops, Sesbania will grow in water, and will grow well in the hot weather of midsummer on low, wet land. It is valuable not only as a green manure cover crop, but also as a means of killing out Bermuda grass, nut grass, and other weeds. In addition, being a leguminous crop, it fixes atmospheric nitrogen in the soil. For seed inoculant. (See page 82.)

We recommend scarified Sesbania seed for quicker and higher germination. Sown in late May, June and July. Just before blooming, the crop may be plowed under and the stalks will rot or decay rapidly. In some sections Sesbania is grown for pole bean sticks. For this purpose it should be sown 12 to 15 lbs. per acre.

Write for prices.

SUNFLOWER

Sow 6 pounds per acre.

MEDIUM CALIFORNIA. The Medium California makes three to four times as much seed as ordinary varieties. It is used extensively as a windbreak for beans. Plant from March to September. Plant seed in drills three feet apart, thinning out to one stalk in the hill when four to six inches high.

1 lb. 55c; 10 lbs. \$2.30; 25 lbs. \$5.25; 100 lbs. \$20.00

VELVET BEANS

Sow 2 pecks (30 lbs.) per acre.

Velvet Beans are good for groves and truck lands in building up to humus; good pasturage for cattle and hogs; and fine land renovators. They are especially valuable because they are practically immune to nematodes causing root knot disease. Plant from March to August. Use Nitragin. (Page 82.)

90-DAY. (90 to 100 days.) This variety makes a very luxuriant growth and an abundant amount of foliage. for pasturage and soil improvement.

Write For Prices.

PLANT POPULATION PER ACRE AT DIFFERENT SPACINGS

Inches apart	Number of plants	Inches apart	Number of plants	Inches apart	Number of plants
12 x 3	174.000	30 x 4	52,300	48 x 24	5,450
12 x 12	43,560	30 x 6	35,000	48 x 36	3,500
18 x 3	116,000	30 x 12	17,500	60 x 36	2,900
18 x 12	29,000	36 x 3	58,000	60 x 48	2,000
24 x 6	43,560	36 x 12	14,500	60 x 60	1,700
24 x 0 24 x 12	21,780	36 x 18	9,680	72 x 36	2,420
24 x 12 24 x 18	15,500	36 x 24	7,260	72 x 48	1,815
24 x 16 24 x 24	11,000	36 x 36	4,840	72 x 60	1,450



VIGORPAK SEED. Your insurance against loss in germination and vigor

Loss from pests

Waste from mixture

PESTICIDE SECTION

(Insecticides, Fungicides, Fumigants, Rodenticides, Weed Killers, Spreaders, Stickers, Emulsifiers—Pages 60 to 78.)

The information and recommendations contained in this publication are generally based on the recommendations of the Florida Agricultural Experiment Stations, the USDA, and experiments conducted by The Kilgore Seed Company and other agricultural chemical companies.

Every attempt has been made to provide such information in a clear and concise form for the benefit of our growers. Since constant changes are taking place in the field such as the introduction of new pesticides, new insect and disease problems; such as insects becoming resistant to some of the established pesticides, it is difficult to keep any annual publication strictly up-to-date. We therefore invite you to contact your local Kilgore Seed Company representative. He will be glad to assist you whenever you are facing new problems.

The use of pesticides is now just as necessary as that of fertilizer and various modern cultural practices. Many factors are involved in the use of pesticides in relation to obtaining the results desired, such as the choice of the proper pesticide for the control of the various diseases and insects and the time and method of application. Satisfactory control cannot be expected from poor or improperly timed applications. For best results the grower should adhere to a regular preventative spray or dust program. It is often impossible to gain satisfactory control after heavy infestations of either insects or diseases are established.

The Food and Drug Administration, through Public Law 518, commonly called "The Miller Amendment," established tolerances for the various pesticides on the various vegetable crops which govern the amount of residue of the pesticide which may legally and safely remain on vegetables at harvest time. It is very important that these tolerances are not exceeded in controlling insects and diseases. Therefore, control must be started early and the crop kept clean so as to avoid applying excessive amounts of pesticides close to maturity or harvest.

See page 61 for more detailed information on the Miller Amendment.

As a rule, it is much cheaper to use a combination spray or dust when one must contend with both insects and diseases, as one application may do both jobs. Most insecticides and fungicides are compatible and can be used together in sprays or dusts. Although we manufacture numerous combination dusts, only the more important are listed in this catalog. If you do not find the combination listed which you want, write to The Kilgore Seed Company, Plant City, Florida, or call at any of our stores. You can be assured that you will receive fresh and perfectly blended dusts formulated to meet your particular requirements. Only highgrade technical materials are used by us in producing the various pesticidal compositions that you may desire.

SANITATION—Serious losses may result from neglect of a few simple rules. Diseases will spread readily from sick to healthy plants by various means, depending upon the nature of the disease. Spores of many species of fungi are air- or wind-borne, others may be scattered over fields in rain or irrigation water. Insects, particularly those of the sucking type, such as aphids and thrips disseminate diseases. Certain

cultural operations, such as pruning, picking, transplanting, etc., may scatter diseases, especially those caused by bacteria and viruses (mosaic being a specific example of the latter).

Considering methods of infection, it becomes obvious that the source of the disease should be destroyed. Affected plants, as well as decayed fruit, should be removed from the field and burned. Cultivation should be clean and frequent. Mechanical injury to plants by plowing, spraying or picking operations should be avoided as much as possible. Crop rotation is essential. The crop rotation program should provide that no host crop of the responsible disease is grown for several years. This method often proves cheaper than an attempt to control the disease chemically. As weeds serve as host plants to many insects, which may be carriers of fungi, viruses and other undesirable organisms, the necessity for their eradication certainly becomes evident. Whenever practical, soil moisture should be kept fairly constant. Corrective measures should be adopted where soils are either too acid or too alkaline.

MOSAIC—Mosaic disease is caused by virus organisms in the sap of the affected plants. The viruses may be transmitted in various ways from diseased to healthy plants. Insects undoubtedly play an important part in the transmission of the disease, although it may be spread by other means, such as pruning operations, bruising of plants by workers or mechanical equipment, picking fruit, the use of tobacco and by other means.

The symptoms of mosaic vary slightly in different plants, but in general, certain broad characteristic indications can be noted in most all infected plant types. There is usually a stunting or dwarfing with accompanying pattern of light yellowish areas merging with green patches, giving a "mottled" effect. Often there is a roughening of the surface, producing, in some cases, a warty appearance. Edges of leaves may cup downward. The fruit also may be mottled and warted.

Mosaic has become costly and destructive on many crops, especially peppers, tomatoes, cucumbers and squash. This disease apparently does not carry over in the soil, provided cover crops are turned under long enough to be decomposed before planting. It is carried from season to season in wild host plants (weeds) in and around the field, and is transmitted to the plants in the field by insects, particularly aphids and thrips.

Pepper and tomato seedbeds should have clean cultivation two or three weeks before planting the seed. The seedbeds should not be close to ditch banks or fence rows, unless the weeds near the beds are destroyed. Smoking or other use of tobacco should not be permitted while working in the seedbeds or handling plants. Tobacco mosaic has been known to be carried on tobacco. If plants become affected with mosaic in the seedbeds, do not set them in the field. Once the plant get the disease there is no known cure.

The State Experiment Stations recognize the severity of the problem and they are now very much engaged in the attempt to develop effective control measures by chemical or biological means. A serious effort is also being manifested in the development of resistant varieties of various vegetables commonly susceptible to virus diseases. In the event of success, such new varieties would hardly be available commercially for several years, and we would again like to stress the importance of sanitary practices and effective insect and weed eradication as the only means of crop damage reduction at the present time.

THE MILLER AMENDMENT . . . As it Concerns the Grower

THE MILLER AMENDMENT is a law which affects all growers who use pesticide sprays and dusts in combating insects, weeds, rodents and plant diseases. It sets up procedures for the Food and Drug Administration in Washington to use in establishing official limits as to how much of the spray or dust may remain on a crop.

"PESTICIDE RESIDUES" means the amount of pesticide chemical left in or on the crop when it is harvested.

A "TOLERANCE" is simply the amount of pesticide which scientists have determined may safely remain as a residue on the food crop without injury to the consumer. The "tolerance" is specific for the pesticide and the crop. It is set by regulation.

PESTICIDE SPRAYS AND DUSTS are essential to assure the continued supply of high quality foods. The law is designed to permit the effective use of these sprays and dusts in the production of food without hazard to the consumer. In order to accomplish this, the USDA and the various State Experiment Stations establish a safe interval between the last application of the pesticide and harvest or other restrictive guides. The safe interval between last application and harvest is usually expressed as "days" or by such phrases as "Do not apply after edible parts begin to form," etc.

The law thus enables the grower to know with reasonable assurance what sprays and dusts he can use on his crops and how he must time his last application to avoid any excessive residue. All fruits, vegetables and grains are covered by the law. For your aid we have included on pages 62 and 63, a chart which states the number of days between last application and harvest and other restrictive information for most of the pesticides for which tolerances have been established.

Information regarding tolerances and safe intervals which may be established after publication of this

catalog will be made available to all our sales personnel and we urge all growers to approach them at any time for this information.

All pesticides used in agriculture are reviewed by the FDA and are either

- (1) Given a tolerance expressed in parts per million (ppm) for specific crops;
- (2) Considered as safe chemicals (A limited number of pesticide chemicals have been officially declared safe and do not require either a tolerance or exemption. These chemicals are sulphur, lime, lime-sulphur, potassium polysulfide, sodium carbonate and sodium polysulfide);
- (3) Exempt from the requirements of a tolerance (Certain pesticides are exempt from the requirement of a tolerance because when used according to prevailing procedure there will be no hazard to public health. When applied to growing crops in accordance with good agricultural practices, the following pesticide chemicals are exempt from the requirement of a tolerance: Bordeaux Mixtures, Copper Acetate, Basic Copper Carbonate, Copper-Lime Mixtures, Copper Oxychloride, Copper Silicate, Copper Sulphate or Basic Copper Sulphate, Copper-Zinc-Chromate, Cuperous Oxide and Copper-A Compound, MGK-264, Petroleum Oils, Piperonyl Butoxide, Piperonyl Cyclonene, N-propyl Isome, Pyrethrins, Rotenone, Ryania and Sabadilla. These materials may be used near harvest, but are not exempt from the requirement of a tolerance when applied to a crop at the time of or after harvest.);
- (4) Given a Zero tolerance (A Zero tolerance means that no amount of pesticide chemical may remain on an agricultural commodity when it is offered for shipment. All chemicals other than those which are considered as safe, exempt or which have had a tolerance established for specific crops, must be considered as having a Zero tolerance.)

Safe Combinations of INSECTICIDES AND FUNGICIDES

Insecticides	Trade Names	Can be used with
AGRIMYCINSTREPTOMYCIN		Neutral Copper compounds, carbamates and most insecticides in wettable form.
PARATHIONMALATHION	(Vapophos) (Kilphos)	DDT, DDD, methoxychlor, chlordane, benzene hexachloride, lindane, toxaphene, aldrin, dieldrin, dithane, "Parzate," "Zerlate," "Fermate," "Manzate," Kryocide and sulphur. Most copper compounds can be used with parathion and malathion for spraying mixtures if mixed freshly and used immediately.
DDT DDD METHOXYCHLOR CHLORDANE BENZENE HEXACHLORIDE LINDANE TOXAPHENE ALDRIN DIELDRIN	(RHothane)("Marlate")	Each other, and also with nicotine, sulphur, copper, dithane, "Parzate," "Zerlate," "Fermate," "Manzate," and "Karathane."
TEPP	(Vapotone)	DDT, DDD, methoxychlor, chlordane, benzene hexachloride, lindane, toxaphene, sulphur.
NICOTINE	#= ## = # = 0	DDT, toxaphene, sulphur, copper, dithane, "Parzate."
KRYOCIDE		Sulphur, copper, dithane, "Parzate," Parathion.
Do not use emulsions with sulphur.		

PESTICIDE

								_												т—		-								$\overline{}$
	ALDRIH	ALLETHRIN	ARAMITE	BHC	CALCIUM ARSENATE	CHLORODANE	CHLOROBENZILATE	COPPER-A-COMPOUND	DDT	DIAZINON	DIBROM	DIELDRIN	DN (Dinex)	DYRENE	ENDRIN	ПРХ	ETHION (Nialore)	FERMATE (Ferbam)	FUNGICIDE 658	GLYODIN	GUTHION	HEPTACHLOR	KARATHANE	KELTHANE	KILCOP '53"	KRYOCIDE	LEAD ARSENATE	LIME SULPHUR	LINDANE	MALATHION
AVOCADOS				В				н	30										н					٨	н	$\lceil \rceil$	H, J		В	7
BEAN5	В	н	B, F	м	В	В		н	5	7		D				21		4						7 1	н	н, л		н	В	1
BEEYS	14				K ,(H)	к		н	к	14		30										D			н	Н, Ј				7
BROCCOLI	14	н	В	к	В	В		н	7	5	4	30			8							D			н	н, Ј			В	3
BRUSSELS SPROUTS	14	н		К	В	В		н	В		4	30			В							В			н	н, л			В	7
CABBAGE	21	н	В	к	В	В		н	7	7	4	21			35		'	60				В			н	Н, Ј			В	7
CABBAGE, CHINESE								н	14		4														н					7
CANTALOUPE	7		В	м	н	E	н	н	E			В							н			В	7	2	н	Н, Ј		н	H, J	1
CARROT	(7)			с	(H)	(H)		н	(H), F			(21)						7							н	н, Ј			G	7
CAULIFLOWER	21	н	В	К	В	В		н	7	5		21			В							В			н	Н, Ј			В	7
CELERY	D		К	К	В	В	\Box'	н	14	10	4	D		H, J				Н, Ј	н			D			н		В		В	7
CITRUS	30		A	В	U	14	н	н	30	21		30	12		Ĺ'	30		A	н			D		7	н	н		н	В	7
COLLARD	14	н	'	к	к	В	'	н	14	<u> </u>	4	21	Ĺ'	Ĺ'											н	H, J	Ľ	<u>_</u> '	к	7
CORN, SWEET	D	$oxed{oxed}$	B, F	с	\perp	н,1	<u> </u>	м	1 1	<u> </u>			<u> </u>	Ĺ'	45	14			м	<u> </u>		D			м.	м	<u> </u>	Ĺ_'	с	5
CUCUMBER	7	<u></u>	В	м	В	E	<u> '</u>	н	E	7		7	<u>'</u>	<u>'</u>	14		<u></u>		н	<u> </u>		В	7	2	н	н, л		н	H, J	1
EGGPLANT	3	$oxed{igspace}$	В	В	В	В	_'	н	5	<u> </u>	<u> </u>	7	<u> </u>	<u>'</u>	14		<u> </u>	<u> </u>		<u> </u>			<u> </u>	2	н	Н, Ј	н, Ј	<u> </u>	В	3
ENDIVE	30		<u> </u>	<u> </u> '	<u></u>		<u></u>	н	14		4	21	<u></u>	<u> </u>											н			<u></u> '	<u></u> '	7
KALE	14	н	<u> </u>	к	к	В	'		21	<u> </u>	4	21	Ĺ'	<u> </u>												Н, Ј			В	7
KOHL RABI	14	н	<u> </u>	k	В	В	<u> </u>		В	<u> </u>		30	<u> </u>	<u> </u>	<u> </u>		<u> </u>	<u> </u>	<u> </u>	<u> </u>					<u> </u>	H, J	<u> </u>		В	7
LETTUCE	30	н	K	ĸ	<u> </u>	В	<u> </u>	н	7, J	10	4	21	Щ'	<u></u>	<u> </u>			30		<u> </u>					н	H, J	<u></u> '	<u></u> '	В	7
MANGOES	21	igspace	<u> </u>	<u> </u> '	_'	<u> </u>	<u> </u>	н	30	<u>'</u>	<u> </u>	45	<u> </u>	<u> </u>	<u></u>		<u></u>	<u> </u>	<u> </u>	<u> </u>	<u> </u>		<u> </u>		н	<u> </u>	<u></u> '	<u> </u>	В	2
MUSTARD	14	н	↓_	м	<u> </u> '	<u> </u>	<u> </u>	M	14	⊥_′	4	21	<u> </u>	<u> </u>			<u></u>	<u> </u>	м	<u> </u>			└	\sqcup	м	H, J	<u></u> '		К	7
OKRA	<u> </u>	<u> </u>	В	<u> </u>	_'	В	<u> </u>		7	<u> </u> '	<u> </u>		<u> </u>		<u> </u>		<u> </u>	4	<u>'</u>	<u> </u>			<u></u> '	\sqcup	<u> </u>	H, J	<u> </u>	<u></u> '	G	В
ОИЮИ	G,D	<u> </u>	 _	C L	<u> </u>	L, D	<u> </u>	н	L	10	<u> </u> '	L, 14	<u> </u>	<u> </u> '	<u> </u>	21,1	l, D	<u> </u>	└	<u> </u>	<u> </u>	5	<u> </u>	\sqcup	н	<u></u> '	<u></u> '	<u></u> '	Ĺ	3
PARSLEY	_	igspace	<u></u>	<u> </u>	<u> </u>	<u> </u>	<u> </u>	<u> </u>	<u> </u>	<u></u> '	<u> </u>	<u> </u>	<u></u>	<u></u> '	<u></u> '		<u></u>	<u> </u>	<u> </u>	<u> </u>			<u> </u>			<u></u> '	<u></u> '	<u> </u>	<u> </u>	7
PARSNIPS	7	<u> </u>		<u> </u>	<u> </u> -	<u> </u>	<u> </u>		H, F	<u> </u> '	<u> </u>	21	<u>'</u>	↓ _'	<u> </u>	<u> </u>	<u></u> '	1	<u> </u>	<u> </u>		<u> </u>	<u></u> '	1	<u> </u>	<u></u>	<u> </u>	└	<u>'</u>	7
PEAS, ENGLISH	3	<u> </u>	 	В	_'	15, F	<u> </u>	н	B, 1	<u> </u> '	<u> </u>	<u> </u>	_	<u></u>	1	<u> </u>	<u> </u>	1	<u> </u>		\sqcup	В	<u></u>		н	Н, Ј	<u> </u>	<u> </u> '	В	4 3
PEAS, SOUTHERN	D	<u> </u>	\vdash	M	<u> </u>	В	<u> </u>	н	5	<u></u> '	<u> </u> '	<u> </u>	<u></u>	<u> </u>	<u></u>	 '	1	<u> </u>	<u> </u>	<u> </u>	1	10	<u> </u>	\sqcup	н	H, J	<u> '</u>	н	'	
PEPPER	3	<u> </u>	В	В	H, J	 	ļ'	н	3	5	ļ	7	 '	—'	14	<u> </u>	<u> </u>	60	<u></u> '	<u> </u>	1	D	<u> </u>	2	н	Н, Ј	н, з	<u> </u>	8	3
POTATOES, IRISH	н	igspace	н	С	н	Н	ļ'	H	н	14	4	21	 	н	3	<u> </u>	<u> </u>	н	н	<u> </u>	<u> </u>	н	<u> </u>	<u> </u>	н	н	н	<u> </u>	С	н
POTATOES, SWEET	н	ـــ	 	С	ليل	D		+-	н	<u> </u>	-	21	-	<u></u>	<u> '</u>	 '	<u> </u>	1	1		1	30		<u></u>	<u> </u>	 '	<u></u>	<u> </u>	С	
PUMPKIN	7	 	 	м	н	E	<u> </u>	н	E	<u> </u>	<u> </u>	В	-	 '	<u> </u>	<u> </u>	 '	 	н	<u>—</u> '	 	В	7	2	н	<u> </u>	<u></u>	<u> </u>	н	3
RADISH	(7)	н	₩	С	₩'	В	<u> </u>	н	н	<u> </u> '		(21)	-	<u> </u>	 '	1-	 '	<u> </u>	<u></u>	<u> </u>	<u></u>	К	<u> </u>	<u> </u>	н	Н, Ј	<u> </u>	1	<u> </u> '	7
RUTABAGA	14	н	 	С	(H).K	В	<u></u>	1	н	<u> </u>	<u> </u>	30	1	<u> </u>	<u></u>	<u>—</u> '	<u>—</u> '	<u> </u> '	<u> '</u>	<u></u>	<u> </u>	(15)	<u> </u>	<u> </u>	<u> </u>	Н, 3	<u>'</u>	<u> </u>	<u> '</u>	3
SPINACH	14	 	K	K	K		<u> </u>	н	14	10	4	21	-	<u> </u> '	 '	<u></u>	<u> </u>	<u></u>	<u></u>	<u> </u>	<u> </u>	<u> </u>	<u> </u>	1_'	н	<u> '</u>	<u> </u> '	<u> </u>	К	7
SQUASH	7	-	В	м	н, ј	+-	-	н	E		<u> </u>	7		<u> </u>	14	 '	<u> </u>	<u></u>	н	<u> </u>	1	В	7	2	н	Н, Ј	<u>'</u>	н	н, ј	1
STRAWBERRIES	В	-	В	В	В	7,1	-	A	5	5	4	В 21		-	A	<u> </u>	<u> </u>	14	<u> </u>	<u> </u>	<u></u>	В	21	2	A	В	В	<u></u>	В	1
SWISS CHARD	14	-	Ļ	 	 '	 	-	1	21		4	21		<u></u>	<u></u>	 	1	<u></u>	<u></u> '	-	<u></u>	<u></u> '	<u> </u>	<u></u>	<u> </u>	—'	<u> </u>	<u> </u>	к	7
TOMATO	3	 	В	B	Н, Ј	14	 	н	3	3	-	7		н	14	3		14	н	-	<u> </u>	D	 	2	н	H, J	H, J	1	В	1
TURNIP	(14)	н	 	, M	,H), K	+	-	м	14	<u> </u>	-	30		<u> </u>	<u></u>	<u> </u>	<u> </u>	<u> </u> '	M	<u> </u>	<u> </u> '	(15)	<u> </u>	<u></u> —'	м	н, ј	<u> </u>	<u> </u>	<u> </u>	3
WATERMELON	7	ـــ	В	M	<u> </u> '	E		н	E	 	↓_	В	_	<u> </u>	<u></u>	<u>'</u>	<u></u> '	<u> </u>	н	<u> </u>	<u> </u>	В	7	2	н	H, J	<u> </u>	н	н	1
					Щ'	↓	\perp	\perp		Ш'	<u> </u>	<u> </u>	↓		Щ'	<u> </u>	<u> </u>	'			Ш'	<u></u> '	<u> </u>	⊥_'		<u> </u>	<u> </u>	\perp		

CODE LETTERS

Apply when no fruit is present. This usually means after harvest of such crop as citrus, apples, peaches, etc.

Do not apply after edible parts start to form.

Do not use—causes residual odor or taste to remain in or on edible parts of the crops.

D—For vegetable and other annual crops—preplanting soil treatment only, for fruit trees and perennials—soil treatment only.

E—Do not use—reduces yield.

F—Do not feed treated crops to livestock.

G—Seed treatment only.

H—No time limitation.



INTERVAL CHART

Part				_		_																								
	MANEB	METHOXYCHLOR	METHYL PARATHION	NICOTINE	ORTHOCIDE (Capien)	OVEX	PARATHION	PERTHANE	PHALTAN	PHOSDRIN	PHYGON XL	PYRETHRINS	ROTENONE	RYANIA	SPERGON	STREPTOMYCIN (Agri-Mycin)	sulfOCIDE	SULPHUR	SYSTOX (Demeton)	TDE	TEDION	ТЕРР	THIMET	THIRAM (Thylote)	THIODAN	TOXAPHENE	TRITHION	ZINEB	ZIRAM (Zerlate)	
	-	_			н		21																			м				AVOCADOS
	_	7	21	3	u l					1	7	н	н		G	B, F	н	н	14	В				G		5	7, F	7	4	BEANS
	-				-		-					н						н						G			21		(H), 7	BEETS
		-	21	-			7	3		,			н		G			н	21	В		3		С	\neg	В		7	7	BROCCOLI
			-					_	-	\vdash								н				3			\neg	8		7		BRUSSELS SPROUTS
	-						-		-													3				7, 3		7	7	CABBAGE
	7	-	-	7			'	3							-	-			21			-	_				_		<u> </u>	CABBAGE, CHINESE
			21					_	-				_		-				21			3		-1			_	-	<u></u>	
	-			3			-		-						-	-		\vdash	21					-				\vdash	├─	
	н				(H)	<u> </u>	-			-					-	-				\vdash									-	
		7	21	7			7	3	-	3		Н	_		G		_		21	В				\vdash			_	-		-
Mathematical Region of the content	H.J		15	7		<u> </u>	7		\vdash	7	н	н			-	K	-					-		н				-	П, 3	
				3	H, I	14	14	_	^	_					_		-	Н	21	14	Α	-					14	Н .	⊢	
		14		7			10		-	3		н	_		-	-					_	3				 		1		
	H,F	7		3	н, 1		3	ļ		1			н	н	G	_	_	н		H, I				G		B, 1		-	-	
Mathematical Reservation of the content of the co	н	1		3	н		3	<u> </u>		1		н	Н		G	В		M		1		3		-		м	_	н	-	CUCUMBER
7 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	н	1		3	н		3			5			Н		G			н	_	1		3		G		5	7	н	н	EGGPLANT
	7 J						10								G			н				3		G				7	_	ENDIVE
		14		7			21			3		н			G			н				3				_		7		KALE
Part		1		7			21	3										н		8		3					ļ	В	7	KOHL RABI
	7 J	14	21	7	н		10	4		2		н			G			н	21	В		3		G		7, 3	_	5	$oxed{}$	LETTUCE
Note					н		21											н												MANGOES
Mathematical Region of the content				7			10			3								н				3				К		7		MUSTARD
				3			3								G			н				3				В				OKRA
Parison of the content of the cont	н	(7),L			н		15			1								н		H, L				G		H, L		7		ONION
Passing Pass		 	\vdash	+	1		\top	1		T					G			н												PARSLEY
Pack of the content				_			1		-						+			н								(H)				PARSNIPS
Part		7	15	+-	-	\vdash	10	-		1	1		н	1	G	1		Н						G		В	7		7	PEAȘ, ENGLISH
H		1		 	\vdash		+		+	+,			н		G	+-	н	н					1	G		5				PEAS, SOUTHERN
H H S S S S S S S S S S S S S S S S S S	-		75	+-	н	+-	+-	+	+-	+		\vdash	н	\vdash		В		н	T	1		3		G		5	7	н	н	PEPPER
H H G G G G G G G G G G G G G G G G G G		 	-	+-	+	\vdash	+		Н	+	н	н	н	\vdash	+-	G	+-	н	21	н	\dagger		D		н	н	н	н	н	POTATOES, IRISH
No. No.	<u> </u>	-	-	3	<u> </u>	-	_	+	+	+	-	+"	+-	+-	+	+-	+-	Н	\vdash	\vdash		1	-			н	\vdash	†	+	POTATOES, SWEET
T	-	+	┼	\vdash	н	╁	-		+	14	-	+-	+-	+	G	+	+	M		1	_	3		G		M	\vdash	н	н	PUMPKIN
T		\vdash	 	+	 "	\vdash	+	+-	+-	+-	+-	+-	+	+-	+	+	+	+	T	+	\vdash	\vdash	\vdash	G		<u> </u>	T	н	н	RADISH
7, J 14 21 7 H 10 7 A H H G H H G H H G H H G H H G H H G H H H G H H H G H H H G H H H G H H H G H H H G H H H G H H H G H H H H G H	-	+	+	3	+-	+-	_	+-	+	+	+	+ "	+-	+	+-	+-	+	+		н	1	+	-	\vdash	t	(H)	+	1	\top	RUTABAGA
H 1 21 7 H 10 7 A N N N N N N N N N N N N N N N N N N	-	7	-	+-	 —	+-	+	+	+-	+-	-	-	+	+-	-	+-	+	+-	+	+	+	-	+-	6	-	T	+-	7	+	
H 1 3 H 3 14 H 7 0 N 7 1 3 N A B 3 7 7.J STRAWBERRIES 14 7 N N N N N N N N N N N N N N N N N N	\vdash	1-	21	+-	+	-	+	7	-		-	+-	+	+	+-	+	+	+	+	+	+	+ -	+	+	+-		1 21	+-	+	
14 3 H A 3 1 1 3 H N	н	1	╀-	3	+	┼-	+-	-	+-	+	-	+	+	+-	G	+-	+	+-	+-	+	-	+	+	Ť	-	+-	+-	+-	+-	
H 1 15 3 H 3 1 H H H B B H 1 M H 1,J 7 H H TOMATO 14.(1) 7 10.(15) 3 G H K 3 G M 1 3 G M 5 H H WATERMELON	_	14	_	3	H	A	3	╁		1	3	Н	H	+-	-	+-	-	-	7.3	5	-	3	+-	6	 ^	+ "	+,	+		
H 1 15 3 H 3 1 H H H H B B I I M K 3 M 5 H H WATERMELON	-	-	<u> </u>	7	1_	\perp	1	_	4	-	-	-	-	+-	+-	+-	+-	+-	+-	+-	+-	+-	-	-	\vdash	, .	+	+		
114.(1) 7 10.(15) 3 G M 1 3 G M 5 H H WATERMELON	н	+	+-	3	Н	_	3	1	1	1	н	H	H	+-	-	+-	+	+	-	+-	-	+-	+-	+	-	+ "	+ 7		+-	
		14. (1)	1_	7	_		10, (1	5)	\perp	3	_	\perp	_	+-	-	+	+-	+	+-	+-	-	+-	+-	+-	+	+	+	+	+-	"
	н	7	1	3	н		3	\perp	\perp	14	1	Н	н	_	G	+	+-	M	+	+-	+	3	+	1 6	-		5		+	TYATERMELON
						\perp								\perp	1_				╽		_		_		<u></u>			\perp		

I—Do not feed treated crops to dairy animals or to animals being finished for slaughter.
 J—Remove excessive residues at harvest.
 K—Apply during the seedling stage only, or do not apply after the seedling stage.

L—Do not apply to green onions.
 M—Do not use—Phytotoxic to this crop in Florida.
 ()—Parentheses indicate root consumption. Numbers indicate minimum days between last application and harvest.

SEED TREATMENT

DAMPING-OFF CONTROL

These recommendations are for the control of various soil fungi which cause damping-off.

5011 Tuligi Willell Guabo		Oz. Per	Tspns Per Lb.
Beans, Lima	Sporgon or Delsan	2	1/3
Beans, Lilla	Spergon or Dolson	2	1/3
Beans, Snap and Pole	Spergon of Delsan	2	$\frac{1}{3}$
Peas, English	Spergon or Deisan	2	
Cowpeas	Spergon or Delsan	2	1/3
Corn	Arasan or		
	Ortho Seedguard	2	1/4
Broccoli	Arasan		$\frac{1}{2}$
Cantaloupe	.Spergon or Semesar	n	1
Carrot	Spergon		3/4
Celery	.Spergon or Semesar	1	$\frac{3}{4}$
Cabbage,	-		
Cauliflower, etc	.Arasan or Semesan		$1/_{2}$
Cucumber	.Spergon or Semesai	n	1
Eggplant	Semesan		$3/_{4}$
Escarolle	Semesan		1
Lettuce	Semesan		1
Pepper	Semesan		1/2
- · ·	Orthocide 50-W	4	
Potato	Gibberellic Acid (s	ee page	80)
Spinach	Spergon or Arasan		$1/_{2}$
Tomato	Semesan		1
Watermelon	Melon Seed Treatr	nent	_
W a cellifetoii	(see page 76)	110110	
	(bee page 10)		

The seed and seed treatment material should be thoroughly mixed together in a rotary drum mixer until all the seed is uniformly coated. In most cases this will take from three to five minutes at a speed of forty revolutions per minute. Do not use more material than recommended. Screen off excess dust.

SEED-BORNE DISEASE CONTROL

Bichloride of Mercury (Corrosive sublimate) treatment: Cucumbers: Dilute 1:1000, immersing seed for 10 minutes.

Less hard seeds: Dilute 1:1000, immersing seeds for 5 minutes.

For small quantities of seeds, dissolve 1 tablet in 1 pint of water.

For larger quantities, dissolve 1 oz. of the powder form in 7½ gallons of water. It is advisable to dissolve the powder in a small quantity (a pint or quart) of boiling water first.

Place seeds in loosely woven cloth bags (about half full). Submerge bags in solution and stir with a stick to loosen adhering bubbles. After soaking the proper length of time, remove bags and drain thoroughly. Seed should be washed for 15 minutes in several changes of water. Then spread in thin layers to dry. Do not prepare solution of bichloride of mercury in metal containers. Only glass, earthenware or wooden vessels should be used. In working with the compound, the greatest caution should be exercised, as it is extremely poisonous.

Hot Water Treatment:

Black Rot. Cabbage seed which is not known to be free from the black rot organism should be soaked in water held at 122° F. for 25 minutes (cauliflower and broccoli, 18 minutes). Use an accurate thermometer and hold the temperature within 1° F. of that specified to control the disease without injuring the seed. Fill a cheesecloth bag two-thirds full of seed, tie the top, immerse it in the hot water and then stir it enough to remove the air. After treatment, dip the seed in cold water, drain and spread in a thin layer to dry.

Use new land for seedbeds or old land in which cabbage or other related crops have not been grown for two years.

Do not use plants from a plant bed in which black rot has been found.

Bureau of Standards Household Measurements

2	Tsp.—1	ounce	1 Cup—8 ounces 2 Cups—1 pint	2 Pts.—1 quart 4 Qts.—1 gal.
16	Tbs.—1	cup		

DILUTION TABLE FOR INSECTICIDES AND FUNGICIDES

AND	ONOICID	LJ	
Material Av	erage amounts	needed to	make spray
Agri-mycin 100 at	50 Gals.	4 Gals.	1 Gal.
200 p.p.m. level	9:34 oz.	4 tbsp.	1 tbsp.
Aldrin 2 Spray	½ pt.		
Alltox 40-W	2 lbs.	8 tbs.	2 tbs.
Alltox Spray No. 4.8	1 1/4 pts.	33/4 tbs.	4 tsp.
Aramite	3/4 lbs.	4 tbs.	I tbs.
Arsenate of LeadBlack Leaf 40	1½ lbs.	10 tbs.	2½ tbs.
C-M-Z Bean Spray	1 pt. 5 lbs.	8 tsp. 20 tbs.	2 tsp. 5 tbs.
C-M-Z Spray Improved	3 lbs.	14 tbs.	3½ tsp.
C-M-Z Garden Spray	21/2 lbs.	8 tbs.	2 tbs.
Copper-A Compound	2 bls.	8 tbs.	2 tbs.
DDT 25% Emulsion	1 pt.	5 tbs.	4 tsp.
DDT 50% Wettable Diazinon 25-E	1 1/4 lbs.	8 tbs.	2 tbs.
Diazinon 25-W	½ pt. ½ lb.	4 tsp. 4 tbs.	1 tsp. 1 tbs.
Dieldrin Emulsion (1.5)	11/4 pts.	33/4 tbs.	4 tsp.
Dithane D-14 (Nabam)	1 qt.	5 tbs.	4 tsp.
(Zinc Sulphate)	6 oz.	3 tsp.	3/4 tsp.
Dithane M-22 (Maneb)	3∕4 lbs.	5 tbs.	4 tsp.
Dyrene	1 lb. 1 lb.	5 tbs. 5 tbs.	4 tsp.
Endrin	6 oz.	3 tbs.	4 tsp. 2 tsp.
Fermate		9 1/3 tbs.	7 tsp.
Fungicide 658	1 lb.	5 tbs.	4 tsp.
Gamtox 10 Wettable	11/4 lbs.	8 tbs.	2 tbs.
Heptachlor—2 Spray	1 pt.	5 tbs.	4 tsp.
Improved Isotox Garden Spray	1	0 4	9 4
Isotox Spray No. 200	l pt. ½ pt.	8 tsp. 4 tsp.	2 tsp.
Isotox Wettable No. 25	½ lb.	4 tbs.	1 tsp. 1 tbs.
Karathane WD	1/4 lb.	2 tbs.	1½ tsp.
Kelthane W	1 lb.	5 tbs.	4 tsp.
Kilchlor 40-W (Chlordane)	11/4 lbs.	6¾ tbs.	4 tsp.
Kilcop "45" Kilcop "53"	Consult la		21/ 11-
Kilphene 6	2 lbs. 1 pt.	10 tbs. 5 tbs.	2½ tbs. 4 tsp.
Kryocide	3 lbs.	13 tbs.	3 1/3 tbs.
Malathion "50" Spray	1 gt.	5 tbs.	4 tsp.
Malathion 25% Wet	1 lb.	7 tbs.	5 tsp.
Marlate 50 Methoxychlor	11/ 11	22.1	
Spray Manganese Sulphate	1 ½ lbs. 1 ½ lbs.	11 tbs. 6 2/3 tbs.	8 tsp.
Manzate (Maneb)	3/4 lbs.	3½ tbs.	5 tsp. 4 tsp.
Oil Emulsion (NP 90)	3 qts.	15 tbs.	3½ tbs.
Orthotran 50W	⅓ lb.	21/2 tbs.	2 tsp.
Ortho Streptomycin		4 .1	
(200 PPM) Orthocide 50-W	8 oz. 2 lbs.	4 tbs.	l tbs.
Orthol Garden Spray	1½ qts.	12 tbs. 7½ tbs.	3 tbs. 5½ tsp.
Ortho-Klor 44 Spray	. /2 415.	7/2 (03.	372 tsp.
(Chlordane)	1½ qts.	8 tbs.	2 tbs.
Ortho-Klor 8 Spray			
(Chlordane) Orthophos 4 Spray	Consult la	ibel	
(Parathion)	3 oz.		
Parzate (Zineb)	1 lb.	5 tbs.	4 tsp.
Phosdrin EC-2	⅓ pt.		. 15p.
Phygon-XL	6 oz.	3 tbs.	21/4 tsp.
Red Arrow Insect SprayRHothane WP-50	1 pt.	8 tsp.	2 tsp.
RHothane 25% Emulsion	1¼ lbs. 1 pt.	8 tbs.	2 tbs.
Spergon Wettable	2 lbs.	8 tsp. 11 tbs.	2 tsp. 8 tsp.
Sulfocide	1 qt.	5 tbs.	4 tsp.
Systox	$\frac{3}{4}$ pt.	2 tbs.	1½ tsp.
Tedion 25-W	1/4 lb.	2 tbs.	1½ tsp.
Terraclor 75-W	2½ lb.	8 tbs.	2 tbs.
Trithion 25-W	1 pt. 1 lb.	8 tsp. 5 tbs.	2 tsp.
Trithion 4 Flowable	½ pt.	4 tsp.	4 tsp. 1 tsp.
Triton B-1956	3 tbs.	3/4 tsp.	20 drops
Triton X-100	3 tbs.	3/4 tsp.	20 drops
Vapophos 15 (Parathion)	½ lb.	*********	*********
Vapotone-XX Spray (TEPP) Volck Isotox Spray	4 oz.	20 46-	P .1
Volck Oil Spray	1 gal. Consult la	20 tbs.	5 tbs.
Volck Soluble Oil	½ gal.	10 tbs.	21/2 tbs.
Zerlate	1 іь.	5 tbs.	4 tsp.
Zinc Sulphate	1 1Ь.	3 tbs.	2 tsp.

INSECT AND DISEASE RECOMMENDATIONS

For detailed information on the Miller Amendment governing the use of pesticides on crops, see page 61.

BEANS— INSECTS		Coined Common Name	
Leaf miners	Kilphos-Sulphur Dust No. 1 Vapophos Wettable 15% Diazinon 25-W Diazinon-Sulphur Dust No. 1	Parathion	1 lb., 100 gals. 1 lb., 100 gals.
Bean leaf rollers	Kilphos-Sulphur Dust No. 1 Kilcide-Sulphur Dust No. 2 DDT Wettable 50%	DDT	20-35 lbs., acre 2 lbs., 100 gals.
Army wormsCutworms	Kilchlor-Kilphene Poison Bait No. 11 Kilphene Poison Bait No. 9 Kilchlor Poison Bait No. 8	I oxaphene	15-30 IDS., acre
Mexican beanbeetles	Kilphos-Sulphur Dust No. 1 Rotenone Dust No. 1	Parathion	20-30 lbs., acre 20-30 lbs., acre
DISEASES Rust	Sulphur Dust Sulphur—Wettable Maneb-Sulphur Dust No. 2		20-40 lbs., acre 10-16 lbs., 100 gals.
Powdery mildew	Maneb 80% W		1½ lbs., 100 gals.



Adult BEAN LEAF HOPPER

BROCCOLI, CABBAGE, CAULIFLOWER-

INSECTS			
	Phosdrin EC-2Kilphos Dust No. 4		I pt., 100 gals.
•	Kilphos Dust No. 4	Parathion	20-30 lbs., acre
Aphids	Vapotone XX Spray	TEPP	I pt., 100 gals.
	Vanonhos Wettable 15%	Parathion	I Ib., 100 gals.
	Orthophos 4-Spray	Parathion	1/3 pt., 100 gals.
	Kilphene 10 Dust	Toxaphene	25-35 lbs., acre
	Marlate 50	Methoxychlor	3 lbs., 100 gals.
	Marlate Dust No. 1	Methoxychlor	25-35 lbs., acre
Cabbage worms	Phosdrin EC-2		1 pt., 100 gals.
	Rotenone Dust No. 1		25-35 lbs., acre
	25% DDT Emulsion		1 qt., 100 gals.
	Endrin 1.6 Spray		1 qt., 100 gals.
	Kilphos Dust No. 4	Parathion	25-35 lbs., acre
Leaf miners	Vapophos Wettable 15%	Parathion	1 lb., 100 gals.
	Diazinon 25-W		1 lb., 100 gals.
Cutworms			
Army worms	Kilchlor-Kilphene Poison	Chlordane	15-30 lbs., acre
Harlequin cabbage bug	Bait No. 11	I oxaphene	15 30 !!
Grasshoppers	, Klipnene roison bait 140. 10	r Uxapnene	13-30 103., acre
Mole crickets		Chlordane	15-30 lbs., acre
DISEASES			
For Seedbeds	Spergon Dust No. 1	Chloranil	20-25 lbs., acre
Downy mildew	Spergon Wettable	Chloranil	4 lbs., 100 gals.
For Crops in Field			
Alternaria leaf spot	Nabam plus zinc sulphate		2 qts. plus $\frac{3}{4}$ lb., 100 gals
Downy mildew	(7ingh 65% W		2 lbs., 100 gals.



CABBAGE LOOPER



Adult HARLEQUIN CABBAGE BUG

CELERY—

Cutworms	Alltox 40 W	Toxaphene3 lbs., 100 gals.					
Army worms	Kilphene Poison Bait No. 9.	Toxaphene15-30 lbs., acre					
Celery leaf tiers Celery loopers	25% DDT Emulsion	Toxaphene20-30 lbs., acre 1 qt., 100 gals Methoxychlor3 lbs., 100 gals. Methoxychlor25-35 lbs., acre					
AphidsThrips	Kilphos Dust No. 4	Parathion20-30 lbs., acre 1 qt., 100 gals. Parathion1 lb., 100 gals.					
DISEASES	SALL I C. J.	2 ats plus 3/4 th 100 gale					
Early blightLate blight	Maneb 80% W	2 qts. plus ¾ lb., 100 gals. Copper 4 lbs., 100 gals. 1½ lbs., 100 gals. 2 lbs., 100 gals. pper-Zinc-Chromate 2 lbs., 100 gals.					
Bacterial leaf spotin seedbeds	Agrimycin 25-50 ppm plus 4 lbs. neutral copper, 100 gals. Strentomycin 25-50 ppm plus 4 lbs. neutral copper, 100 gals.						



IMPORTED CABBAGE WORM



CELERY LEAF-TIER

INSECT AND DISEASE RECOMMENDATIONS (Continued)

For detailed information on the Miller Amendment governing the use of pesticides on crops, see page 61.

Coined Common

CORN—		Name	<i>on</i>
INSECTS	Kilphos Dust No. 4	Parathion	30-35 lbs., acre
Corn earworms	Kilphos-Kilcide Dust No. 4	DDTDDT	30-35 lbs., acre 30-35 lbs., acre 2 lbs., 100 gals. 2 lbs., 100 gals.
Corn lantern flies Corn silk flies	Kilphos Dust No. 4 Vapophos Wettable 15%	Parathion	2 lbs., 100 gals.
Cutworms	Alltox 40 W Ortho-Klor 8	Chlordane	Broadcast 1 gal., acre
wire worms	Heptachlor Emulsion	***************************************) broadcast 2 gais., acre
DISEASES Corn leaf blight	Maneb 80%		1 ½ lbs., 100 gals.
	Maneb Dust No. 1		



CORN EARWORM (TOMATO FRUITWORM)



CO	W	/P	EΑ	۱S–
----	---	----	----	-----

INSECTS	Kilphene 5—Sulphur Dust	Toxaphene	25-35 lbs., acre
Pod weevils	Kilcide-Sulphur Dust No. 2	DDT	20-35 lbs., acre
DISEASES Scab and leaf blight	Kilcop Dust No. 1 Nabam plus zinc sulphate		



CUCHMBER SOLIASH CANTALOUPE

CUCUMBER, SQUAS	H, CANTALOUPE—		
•	Kilphos Dust No. 1		
INSECTS	Kilgam Dust No. 2	Lindane	25-35 lbs., acre
	Vapophos Wettable 15%	Parathion	1 lb., 100 gals.
Aphids	Vapotone XX	TEPP	1 pt., 100 gals.
	J Isotox Wettable No. 25	Lindane	1 lb., 100 gals.
	Kilphos Dust No. 2	Parathion	30-35 lbs., acre
Leaf miners	Vapophos Wettable 15%		
	Diazinon 25-W		1 lb., 100 gals.
•	Kilgam Dust No. 2	Lindane	25-35 lbs., acre
Cucumber beetles	Kilphos Dust No. 1	Parathion	25-35 lbs., acre
Pickle worms	Marlate 50	Methoxychlor	3 lbs., 100 gals.
Melon worms	Marlate Dust No. 1	Methoxychlor	25-35 lbs., acre
	Isotox Wettable No. 25		
	J Vapophos Wettable 15%		
DISEASES	Zineb Dust No. 1		25-35 lbs., acre
Downy mildew	Nabam plus zinc sulphate		2 qts. plus ¾ lb., 100 gals.
<u>-</u>	J Maneb 80%		1½ lbs., 100 gals.
Powdery mildew	Karathane 25-W		6 to 8 oz., 100 agls.
Powdery mildew	Karathane Dust No. 1		20-30 lbs acre
)		,
Angular leaf spot	Orthocide 50-W	Captan	4 lbs., 100 aals.
Anthracnose	Zineb		2 lbs., 100 ggls.
) :		



Adult STRIPED CUCUMBER BEETLE

EGGPLANT-

INSECTS			
Aphids	Kilphos Dust No. 2	Parathion	20-60 lbs., acre
Thrips		Parathion	1 lb., 100 gals.
Leaf Miners	Vapoptone XX	TEPP	1 pt., 100 gals.
	Kelthane W		2 lbs., 100 gals.
Red spiders	Vapotone XX	TEPP	1 pt., 100 gals.
Ned spiders	Kilphos Dust No. 4		
	Kelthane Dust No. 1		25-50 lbs., acre
Cutworms	Kilchlor-Kilphene Poison Bait No. 11	Chlordane	15-30 lbs acre
Mole crickets	Poison Bait No. 11	Toxaphene	13-30 1b3., dcie
Fruitworms	Kilphene 10 Dust		30-35 lbs., acre



SQUASH BUG

FLOWERS—

INSECTS			
Caterpillars	Kilgore's General Purpose Dust for		
Beetles	Flowers and Shrubs		
Aphids	Isotox Garden Spray	Lindane	1 qt., 100 gals.
Thrips	Red Arrow Insect Spray	Pyrethrum	
Cutworm, mole crickets	Kilgore's Poison Bait No. 11	Toxaphene- Chlordane	15-30 lbs., acre
Snails and Slugs	Kilgore's Slug Dust		15-30 lbs., acre
DISEASES	Kilgore's General Purpose Dust for Flowers and Shrubs		See Label
Leaf Spot	Dithane Z-78	Zineb	2 lbs., 100 gals.
Powdery Mildew	Orthocide 50-W	.Captan	4 lbs., 100 gals.




SQUASH VINE BORER

INSECT AND DISEASE RECOMMENDATIONS (Continued)

For detailed information on the Miller Amendment governing the use of pesticides on crops, see page 61.

FRUIT	TREES-
--------------	--------


INSECTS		Coined Comm	
AphidsSpider MiteWhite FlyScale InsectsRust Mite	Malathion 50 Spray Zineb 65% Volck Oil Spray	Name	2 qts., 100 gals. 2 lbs., 100 gals. See Label
Mealybugs Orange Dog Grasshoppers	Malathion 50 Spray Gamtox 10-W D.D.T. 50-W	ВНС	2 qts., 100 gals. 2½lbs., 100 gals. 4 lbs., 100 gals.
DISEASES Melanose Citrus Scab Anthracnose Greasy Spot	Kilcop "53" Copper	Copper	1½-3 lbs., 100 gals.
Mineral Deficiency	} Es-Min-El	**********	4 lbs., 100 gals.



GREEN PUMPKIN BUG (SOUTHERN STINK BUG)

GLADIOLUS—


Thrips 1 pt., 100 gals. DDT 50% Wettable 4 lbs., 100 gals. Army worms Dieldrin Emulsion 2 pt., 100 gals.	
Thrips DDT 50% Wettable 4 lbs., 100 gais.	
Army worms Dieldrin Emulsion	
(1 5 the technical per ad) 1 Dts., IUU dals.	
Signature Description 75 35 lbc dete	
Riphos Dust No. 2. Parathion 23-35 loss, detection of the parathion of the	
Red spider	
Kelthane W	
DISEASES	
Curvularia Orthocide 50-W Captan 4 lbs., 100 gals.	
Botzwiis Naham plus zinc sulphate 2 qts. plus ¾ lb., 100 g	jals.
Botrytis	_



PEPPER WEEVIL

LAWNS-

NSECTS	Kilgore's 25% DDT Emulsion	1 qt. to 1000 sq. ft.
	Kilgore's Kil-Chinch	1 qt. to 1000 sq. ft. Treat infested areas with 2 lbs., 100 sq. ft.
Chinch bugs	Trithion 2-E	qt. to 3000 sq. ft.
	Diazinon 25 Emulsion	1 pt. to 1300 sq. ft. 1/4 pt. to 1000 sq. ft.
Army worms	Saba Chinch	Treat intested areas
Mole crickets	Kilgore's 25% DDT Emulsion	1 qt. to 1000 sq. ft.
Ants	.) Kilchlor Dust No. I } Kilgore's Nema-Kil	Chlordane3 lbs. per 1000 sq. ft. Nemagon1 qt. to 600-800 sq. ft.
Nematodes	. Kilgore's Nema-Kil	Nemagon



Adult
COLORADO POTATO BEETLE

LETTUCE, ESCAROLLE—

Mole crickers	Kilchlor-Kilphene Poison Bait No. 11	
Aphids and loopers	Vapophos Wettable 15%	Parathion
DISEASES Downy Mildew	Zineb 65%	2 Ibs., 100 gals. 1½ Ibs., 100 gals.



INSECTS	Variations VV	TEPP	1 pt., 100 gals,
Aphids	Vapotone XX Kilphos Dust No. 1	Parathion	25-35 lbs., acre
Leaf miners	Vapophos Wettable 15%	Parathion	1 lb., 100 gals.
Pumpkin bugs	Kilphene 10 Dust	Toxaphene	25-35 lbs., 100 gals.
Okra caterpillars	Alltox Wettable 40	Toxaphene	$2\frac{1}{2}$ lbs., 100 gals.



Wingless Adult

APHID

ONION-

INSECTS	Gamtov Wettable 10	BHC	2½ lbs., 100 gals.
Thrips	Gamtox Wettable 10 DDT 25% Emulsion Vapophos Wettable 15%	Parathion	1 qt., 100 gals. 1 lb., 100 gals.
DISEASES			
Alternaria Leaf Blight. Downy Mildew	Zineb 65%		2 lbs., 100 gals.


INSECT AND DISEASE RECOMMENDATIONS (Continued)

For detailed information on the Miller Amendment governing the use of pesticides on crops, see page 61.

INSECTS		Coined Commo Name	on
	Vapotone XX Vapophos 15% TEPP 40	Parathion	1 lb., 100 gals.
Sod webworms Army worms Grasshoppers	Kilnhene 10-Dust*	Tovanhone	20-30 lbs acre

* CAUTION: Keep cattle off pastures for 2 weeks after applying these insecticides. Do not use in pastures for dairy cows or animals being finished for slaughter.

PEANUTS—		
INSECTS	Kilcide-Sulphur Dust No. 2DDT	25-35 lbs., acre
	Kryocide—Sulphur	
caterpillars	Dust No. 4	25-35 lbs., acre
DISEASES Leaf spot	- Dusting Sulphur	15-25 lbs., acre
Lour Spor	- Pasting Julphar	13-23 lbs., acre



RED SPIDER

CUTWORM



TOMATO PINWORM



Adult MOLE CRICKET



LEAF-FOOTED PLANT BUG

PEPPER—

INSECTS			
	Kilphos Dust No. 1 Vapophos Wettable 15% TEPP 40	Parathion	1 lb., 100 gals.
Fall army worms Blister beetles	Kilhex Dust No. 7 DDT 25% Emulsion. Gamtox Wettable 10 Marlate 50	BHCBHCBHC	25-30 lbs., acre 1 qt., 100 gals. 2½ lbs., 100 gals. 3 lbs 100 gals.
DISEASES			•
Alternaria leaf spot	Improved C-M-Z Spray Kilcop Dust No. 3 Kilcop "53" Zineb 65-W Nabam plus Zinc Sulphate Zineb 65-W Maneb 80%	Copper	4 lbs., 100 gals. 2 lbs., 100 gals. 2 qts. plus ¾ lb., 100 gals. 2 lbs., 100 gals
Bacterial leaf spot	Agrimycin (100 PPM)plus 4 Ib Agrimycin 500	s. neutral copper	, 100 gals. 5 lbs., 100 gals.
Damp-off	Orthocide 50-W	Captan	4 lbs., 100 gals. 4 lbs. each, 100 gals.
Blossom-end rot	Calcium Chloride		4 lbs., 100 gals.

POTATOES—

INS	EC	TS
-----	----	----

AphidsLeaf Miners	Thiodan EC-2. Kilphos Dust No. 1. Vapophos Wettable 15% Phosdrin EC-2.	Parathion Parathion	1 qt., 100 gals. 20-40 lbs., acre 1 lb., 100 gals.
_	Diazinon 25-W		1 lb., 100 gals.
Army worms	Kilphene 5 Dust Alltox Wettable No. 40 DDT Emulsion 25%		
DISEASES Late blight	Zineb Dust No. 2 ≻Nabam plus zinc sulphate Zineb 65-W		2 ats plus 3/4 lb 100 cale

POTATOES (SWEET)—

ľ	N	S	E	C	T	S	
---	---	---	---	---	---	---	--

Caterpillars	Kilphene-10 Dust Alltox Wettable No. 40	Toxaphene 25-35 lbs acre
Grasshoppers	Alltox Wettable No. 40	Toxaphene2½ lbs., 100 gals.
Sweet Potato Weevil	Dieldrin Dust No. 1	75 lbs., acre
Bweer rolling weeviling	Heptachlor Dust No. 1	75 lbs acre

RADISH, TURNIP, MUSTARD-

KADISII, IQ	MINI / MINISTAND—	
INSECTS	Kilphos Dust No. 1	Parathion20-40 lbs., acre
Aphids	Kilgam Dust No. 2	Lindane 20-40 lbs., acre
Foliage worr		
	msPhosdrin EC-2	½ pt., 100 gals.

INSECT AND DISEASE RECOMMENDATIONS (Continued)

For detailed information on the Miller Amendment governing the use of pesticides on crops, see page 61.

Caland Common

SHRUBS—

INSECTS	'	Coined Common Name	•
Aphids	Kilgore's General Purpose Dust for		
Spider Mites	Flowers and Shrubs	**********************	See Label
Scale Insects	Malathion 50 Spray	*****	2 qts., 100 gals.
Thrips	Flowers and Shrubs Malathion 50 Spray	Lindane	1 qt., 100 gals.
Caterpillars)		
Mealybug	Volck Isotox Spray	Oil-Lindane	See Label
White Fly	Volck Isotox Spray		
DISEASES	Kilgore's General Purpose Dust for		
Petal Blight	Flowers and Shrubs	***************************************	See Label
Leaf Spot	Kilgore's General Purpose Dust for Flowers and Shrubs	Zineb	2 lbs. 100 gals.
Iron deficiency (soil)) 101		Soo Labol
Iron and Other	Kilgore's Improved CM7 Garden S		5 lbs 100 anls
Mineral deficiencies	Kilgore's Iron Aid Kilgore's Improved CMZ Garden S	pi ay	J 103., 100 gais.





STRAWBERRIES—

INSECTS			
-	Kilhex-Sulphur Dust No. 2	BHC	25-30 lbs., acre
	Vapophos Wettable 15%	Parathion	2 lbs., 100 gals.
Apinus	Malathion 25-W		2 lbs., 100 gals.
Thrips	Kilphos-Sulphur Dust No. 6	Parathion	20-25 lbs., acre
Red Spiders	Tepp 40%		34 pt., 100 gals.
	tobb da to		05 50 11

	Kelthane Dust No. 1		25-50 lbs., acre
Pameras Flea beetles	Alltox 40-W Kilcide-Sulphur Dust No. 1 Kilgam-Sulphur Dust No. 1	Toxaphene	3 lbs., 100 gals. 20-25 lbs., acre 20-25 lbs., acre
Mole crickets	Kilchlor Poison Bait No. 8 Kilchlor-Kilphene Poison	Chlordane	15-25 lbs., acre
Field crickets			20-25 lbs., acre
Bud worms(Lesser cornstalk	DDT 25% Emulsion	Parathion	2 qts., 100 gals. 2 lbs., 100 gals.

(Lesser cornstalk borer)	Vapophos Wettable 15%	Parathion2 lbs., 100 gals.	
•	Hentachlor	Broadcast 2 gals., acre	
Wireworms		Broadcast 2 gals., acre	
11 11 0 11 V1 1113	Ortho Vice 8	Chlordone Broadcast 1 gal., acre	



Common Leaf Spot Kilcop Dust No. 2	
Kilcon "53" Copper4 lbs., 100 gais.	
Leaf Scorch	
Leai Bright Maneb 80% 1/2 lbs., 100 gais	i.
Fruit Rots Orthocide 50-W Captan 4 lbs., 100 gals.	




Adult THRIP

TOBACCO—

INSECTS			
Aphids	Kilphos Dust No. 1	Parathion TEPP	1 lb., 100 gals. 1 pt., 100 gals.
Bud worms	RHothane Dust No. 1	TDE-DDD	15-35 lbs., acre 2 lbs., 100 gals. 2 lbs., 100 gals.
Horn worms	RHothane Dust No. 1	TDE-DDD Cryolite TDE-DDD	25-30 lbs., acre 15-35 lbs., acre 2 lbs., 100 gals.
Mole crickets	*CAUTION: Do not use directly	Chlordane	15-20 lbs., acre

Scatter in paths or walk ways.



P1021020 1	
Blue mold. Zineb Dust No. 2. 1½ to 4 lbs. to 100 so 2 2 lbs. 100 gals. 4 lbs., 100 gals. 5 lbs., 100 gals.	į. yds.



Adult CHINCH BUG



Nymph CHINCH BUG

The Kilgore Seed Company processes many combinations of insecticides and fungicides which are too numerous to recommend in our General Catalog. If you do not find the combination you want, write our General Office in Plant City or call at your nearest Kilgore Store; it is likely we will have a dust in stock to fill your requirements.



INSECT AND DISEASE RECOMMENDATIONS (Continued)

For detailed information on the Miller Amendment governing the use of pesticides on crops, see page 61.

TOMATOES—		Coined Common Name	(F)
INSECTS	Kilphos Dust No. 1	Parathion25-35 lbs., acre	and the same
Aphids	Vapophos Wettable 15%	Parathion	ZEN ZEN ZEN
Thrips	Diazinon 25-W	2 lbs., 100 gals. BHC2 lbs., 100 gals.	A CONTRACTOR OF THE CONTRACTOR
•	Gamtox Wettable 10	BHC2 lbs., 100 gals.	
) Kilphos Dust No. 1	Parathion25-35 lbs., acre	A SHILL
Army worms	Vapophos Wettable 15%	Parathion	THE PLANT
Cutworms	DDT 25% Emulsion	1 qt., 100 gals.	
Tomato fruit worms	>DDT 50% Wettable		
Pin worms	RHothane WP-50		
	Kilchlor-Kilphene Poison	Chlordane20-25 lbs., acre	A. M.
	J Bait No. 11	TDE—DDD2 lbs., 100 gals. Chlordane20-25 lbs., acre Toxaphene	
Leaf miners	Kilphos Dust No. 1	Parathion 25-35 lbs., acre	*
Pumpkin bugs	Vapophos Wettable 15%	Parathion	GRASSHOPPER
Leaf-footed plant	Diazinon 25-W	2 lbs., 100 gals.	G
bugs	Phosdrin EC-2	2 lbs., 100 gals. 1 pt., 100 gais.	
) Alltox 40-W		
Horn worms	RHothane 25% Emulsion		1
TIOTH WOTHIS	Marlate 50	Methoxychlor3 lbs., 100 gals.	la la
	Marlate Dust No. 1	Methoxychlor25-35 lbs., acre	ACY
DISEASES			
Tata blimba) Maneb 80%		
Early blicks	Zineb Dust No. 2	30-40 lbs., acre	ASSOCIATION OF THE PROPERTY OF
Early blight		2 qts. plus 3/4 lb., 100 gals.	
Gray leaf-spot(Red rust)	Zineb 65%	2 lbs., 100 gals.	
(Red rusi)	J Dyrene	2 lbs., 100 gals.	
	-		

Phygon XL......Dichlone......34 lb., 100 gals.

Fungicide 658......Copper-Zinc-Chromate.....2 lbs., 100 gals.

Agrimycin (100 PPM).....plus 4 lbs. Neutral Copper, 100 gals.

......5 lbs., 100 gals.

TOMATO HORNWORM

WATERMELON—

Botrytis.....

(Gray mold)

Bacterial leaf spot.....

INSECTS AphidsLeaf miners	Kilgam Dust No. 2 Kilphos Dust No. 1 Vapophos Wettable 15% Tepp 40%	Lindane Parathion Parathion	15-40 lbs., acre 20-35 lbs., acre 1 lb., 100 gals. ½ pt., 100 gals.	
Rind worms	Kilgam Dust No. 2 Kryocide Dust No. 1 Phosdrin EC-2	Lindane Cryolite	30-40 lbs., acre 30-40 lbs., acre 1 pt., 100 gals.	
Cuiworms		Chlordane	20-25 lbs., acre	

Agrimycin 500.....

DISEASES

Cercospora Leaf Spot Downy mildew	Zineb Dust No. 2	 30-40 lbs. acre
Anthracnose	Zineb 65%	 2 lbs., 100 aals.



MELON OR ONION THRIPS

TERMITE CONTROL

Termites	Ortho-Klor 8 Spray Dieldrin 1.5 Spray	Chlordane1 qt., 25 gals. 1 qt., 15 gals.
----------	--	---



Apply to infected soil areas with low pressure spraying equipment. Thoroughly soak area. Where outside trenching operations next to foundation walls are desired, dig trench 1 to 2 ft. deep and 8" to 12" wide. Apply at the rate of one gallon per 2½ lineal feet of trench. Also treat soil as trench is being refilled. For treating fill dirt under concrete, apply 1 gal. diluted material per 10 sq. feet—Make diluted spray with 2 pints Ortho-Klor 8 or 3-1/3 pints of Dieldrin 1.5 spray to 25 gals. water. If above recommendations are followed, present results show that the protection period should be at least five years.

INSECT AND DISEASE RECOMMENDATIONS (Continued)

For detailed information on the Miller Amendment governing the use of pesticides on crops, see page 61.

SOIL FUMIGATION

DD Soil Fumigant, see page 76 Nematodes Methyl Bromide, see page 76 As preplanting Soilfume (EDB), see page 76 treatment VPM, see page 76 Kilgore's Nema-Kil, see page 76 Nematodes On established plants.

VC-13, see page 76 Nema-Kil® 17.5 G, see page 76

Wire worms and other Aldrin 2 Spray......2 gals., acre in 100 gals. water soil insects..... (Soil application only)

WIRE WORM

Dosage Guide for Bed-Over and Under Row Treatment


Row Spacing	12"	24"	30′′	36"	40′′	42''	48''	54"	60''	66''	72''
Lineal Ft./A.	43,560	21,780	17,424	14,520	13,068	12,445	10,890	9,680	8,712	7,920	7,260
Oz./100 Lin. Ft.	,	,.		•	,	, c	Gallons Pe	r Acre/Ro	w Treatm	ent	
4 (1/4 pi.)	13.6	6.8	5.4	4.5	4.1	4.0	3.4	3.0	2.7	2.5	2.3
8 (½ pt.)	27.2	13.6	10.9	9.0	8.2	7.8	6.8	6.0	5.4	4.9	4.5
12 (3/4 pt.)	40.8	20.4	16.3	13.5	12.2	11.7	10.2	9.1	8.2	7.4	6.8
16 (1 pt.)	54.4	27.2	21.7	18.0	16.3	15.6	13.6	12.1	10.9	9.9	9.1
20 (1½ pt.)	68.1	34.0	27.2	22.6	20.4	19.4	17.0	15.1	13.6	12.4	11.3
24 (1½ pt.)	81.7	40.8	32.6	27.1	24.5	23.3	20.4	18.1	16.3	14.8	13.6

LIVESTOCK PESTS

DAIRY CATTLE-

Coined Common Name

DuPont Dairy Cattle Insecticide......Methoxychlor.... Follow label directions



ADULT CATTLE FEVER TICK

DAIRY BARNS AND		Coined Common Name
Flies	Diazinon 25-E	

CAUTION: All feed troughs, feed and utensils should be covered before spraying.

RANGE CATTLE-

SPRAYS	Coined Common Name		
Mosquitoes	DDT 50% Wettable	BHC	3¾ lbs., 100 gals.
Screw worms	"Smearex," Screw Worm Preparat "Ortho 1038 Screw Worm Contr Ortho EQ 335 Screw Worm Rem Barry's Screw Worm Bomb (Aeros	ion ol" edy Follow	Label Directions
DIPS*	Ortho Healthy Herd Wettable Powder	BHC & DDT	10 lbs., 100 gals.
Flies Ticks	and Dip No. 30"	BHC & DDT	11 lbs., 100 gals.
Lice Mange mites	I Turkho Kiloon Stock Spray	Lindane and Toxaphene	



ADULT HORN FLY

*CAUTION: In using dipping vats, all traces of arsenicals, oils or sludge from such materials must be thoroughly removed before charging with Gamtox, Isotox, "Marlate" or DDT: A mixture of arsenicals or oil solutions with Gamtox, Isotox, "Marlate," or DDT is highly toxic to animals and may cause death. Directions on all containers should be carefully followed.

Lindane and See label for

PESTICIDES

For household and garden insecticides, see Page 81

When ordering pesticides allow enough for postage because all prices quoted are f.o.b. Plant City or any Kilgore store unless otherwise specified. Inquire of your postmaster for proper parcel post rates.

All pesticides not used for commercial agriculture purposes are subject to 3% State Sales Tax.

All prices quoted are subject to change without notice.

INSECTICIDES

ALDRIN 2 SPRAY—A liquid formulation containing 2 lbs. aldrin per gallon. 1 gal. \$4.69; 5-gal. can \$22.20; 30 gals. \$130.20; 53 gals. \$224.72.

ALDRIN 4 SPRAY. Contains 4 lbs. Aldrin per gallon. 5 gal. can \$34.75; 55-gal. drums \$371.25.

ARAMITE—15-W. An effective miticide. Used on citrus, ornamentals and vegetables. Controls red spider mites. Use 1½ lbs. per 100 gals. Price 4-lb. bag \$3.25; 50 lbs. \$38.50.

ARSENATE OF LEAD POWDER. Use 3 lbs. to 100 gals. water. 1 lb. canister 98c; 4 lbs. \$1.45.

BENZENE HEXA-CHLORIDE PREPA-RATIONS.

BHC 10% WETTA-BLE—Use 2½ lbs. per 100 gals. water. 5-lb. bag \$1.30.

KILHEX DUST — No. 7 (Code No. G-3) — (BHC) Contains $1\frac{1}{2}$ % gamma isomer. 1 lb. 15c; 25 lbs. \$1.80; 50 lbs. \$2.55; 100 lbs. \$5.10.



KILHEX-SULPHUR DUST NO. 1—(Code No. GS-1)—(BHC) Contains 1% gamma isomer. Lb. 15c; 25 lbs. \$2.05; 50 lbs. \$3.05; 100 lbs. \$6.10.

BLACK LEAF 40. For the control of aphids, thrips. Use from 1 to 2 pts. per 100 gallons water. Is effective as a spray against poultry lice. 2 oz. \$1.19; 6 oz. \$2.39; 12 oz. \$3.49; 5 lbs. \$8.75; 10 lbs. \$16.85. Prices quoted are prepaid.

CHLORDANE PREPARATIONS

KILCHLOR 40-W. (Contains 40% technical chlordane.) Use $2\frac{1}{2}$ lbs., per 100 gals. of water. 1-lb. can 90c; 5-lb. bag \$2.65; 10-lb. bag \$5.30; 50 lbs. \$24.50.

ORTHO-KLOR "44"—(Containing 44% liquid chlordane.) Readily emulsifiable in water. For control of ants, mole crickets, cutworms, army worms, grasshoppers, roaches and many other insects. 4 oz. 79c; 8 oz. \$1.19; pt. \$1.98; qt. \$2.98.

ORTHO-KLOR-8 SPRAY. (Contains 72% chlordane.) 1 gal. \$10.15; 5 gals. \$49.50. Use 2 qts. per 100 gallons for soil applications.

KILCHLOR DUST NO. 1 (Code No. H-1)—(Contains 5% chlordane.) Lb. 20c; 5 lbs. 80c; 10 lbs. \$1.50; 25 lbs. \$2.80; 50 lbs. \$4.60; 100 lbs. \$9.20.

DDT PREPARATIONS

KILCIDE 50W (50% DDT). Use 2 lbs. to 100 gals. water. 1-lb canister 75c; 3-lb. bag 78c; 4 lbs. \$1.04; 50-lb. bag \$11.50.

DDT 25% EMULSION. Use 2 qts. per 100 gals. water. Qt. \$1.35; gal. \$3.15; 5 gals. \$11.25; 30-gal. drum \$63.00; 53 gals. \$106.00.

KILCIDE 5 DUST—(Code No. D-3)—(Contains 5% DDT). Lb. 15c; 25 lbs. \$1.75; 50 lbs. \$2.45; 100 lbs. \$4.90.

KILCIDE-SULPHUR DUST NO. 2 — (Code No. DS-2)—(Contains 5% DDT). Lb. 15c; 25 lbs. \$2.15 50 lbs. \$3.25; 100 lbs. \$6.50.

KILCIDE 10 DUST — (Code No. D-4) — (Contains 10% DDT). Lb. 15c; 25 lbs. \$2.25; 50 lbs. \$3.45; 100 lbs. \$6.90.

KILCIDE MANGANESE-SULPHUR DUST NO. 2— (Code No. DXS-2)—(Contains 5% DDT). Lb. 15c; 25 lbs. \$2.30; 50 lbs. \$3.55; 100 lbs. \$7.10.

DIAZINON 25-E. An excellent chinch bug control. Use 4 oz. in 25 gals. water per 1000 sq. ft. Also used as a long lasting residual spray for flies and roaches. 1 pt. \$3.50; 1 gal. \$18.00.

DIAZINON 25-W. Use 1 lb. per 100 gals. for leaf miners and worms on tomatoes and an excellent control of aphids, moths, beetles, and pumpkin bugs on many other crops. 4 lb.-bag \$6.00; 50-lb. drum \$62.50.

DIELDRIN EMULSION. (Contains 1.5 lbs. technical dieldrin per gal.) 1 gal. \$6.45; 5 gals. \$31.00; 53 gals. \$318.00.

DIELDRIN—10% Granular. Used for control of flies, mosquitoes, beetles, etc. Consult label for complete information. 5-lb. bag \$3.00; 50-lb. bag \$20.65.

DIELDRIN—50% wettable. For the control of many insects. Consult label for complete information. 5 lbs. \$10.20.

ENDRIN—1.6 spray. Use strictly in accordance with label cautions, warnings and directions. 1 gal. \$11.15; 5 gals. \$54.50.

HEPTACHLOR—2 Spray. For control of wire worms and many soil borne pests. Consult label. 1 gal. \$4.69; 5 gals. \$22.20; 53 gals. \$227.37.

KELTHANE EC EMULSION. New effective miticide for use on flowers and ornamentals. Use 1 pint to 1 qt. in 100 gals. water, or 1 to 2 teaspoonfuls per gallon. 1 gal. \$10.00; 5 gals. \$45.00.

KELTHANE-W—Wettable powder. Use 1 to 2 lbs. per 100 gal. water, 4 to 8 level tablespoonfuls in 3 gallons water. 4 lbs. \$4.40; 50 lbs. \$52.50.

KILGORE'S SLUG DUST. For controlling slugs in nurseries, greenhouses, etc. Use 2 lbs. to 1000 sq. ft. 1-lb. canister \$1.00; 6-lb. bag \$3.30; 50-lb. drum \$25.00.

KRYOCIDE (Natural Cryolite). A fluorine spray for chewing insects. Use 3 pounds Kryocide to fifty gallons of water. 1-lb. can 45c; 4-lb. bag \$1.10; 50-lb. bag \$10.65.

KRYOCIDE DUST No. 1 (Code No. K-10)—(Contains 27% Kryocide). A fluorine dust especially recommended for worms. Lb. 20c; 25 lbs. \$2.85; 50 lbs. \$4.65; 100 lbs. \$9.30.

KRYOCIDE-SULPHUR DUST No. 4 (Code No. KS-4) — (Contains 27% Kryocide.) Lb. 25c; 25 lbs. \$3.10; 50 lbs. \$5.20; 100 lbs. \$10.40.

LINDANE PREPARATIONS

ISOTOX SPRAY—No. 200 (Contains 20% gamma isomer). Use 1 pt. per 100 gals. water. 4-oz. bottle 95c; 16 oz. bottle \$2.25; 1 gal. \$12.20.

ISOTOX—No. 25-W (Contains $25\,\%$ gamma isomer). Use 1 lb. per 100 gals. water. For cucumbers, squash, melons, etc. 1-lb. bag \$1.67; 4-lb. bag \$6.52; 20 lbs. \$31.80.

INSECTICIDES (Continued)

LINDANE PREPARATIONS-(Continued)

KILGAM DUST—No. 1 (Code No. Q-4)—(Isotox) Contains 1% gamma isomer of Lindane. Lb. 20c; 25 lbs. \$2.45; 50 lbs. \$3.85; 100 lbs. \$7.70.

KILGAM DUST—No. 2 (Code No. Q-5)—(Isotox) Contains 11/2 % gamma isomer of Lindane. Lb. 20c; 25 lbs. \$3.00; 50 lbs. \$4.95; 100 lbs. \$9.90.

KILGAM-SULPHUR DUST No. 1 (Code No. QS-1) (Isotox) Contains 1% gamma isomer of Lindane. Lb. 20c; 25 lbs. \$2.85; 50 lbs. \$4.70; 100 lbs. \$9.40.

MALATHION 50% EMULSION. Use 1 qt. per 100 gals. water. 4 oz. \$1.19; 1 pt. \$2.89; 1 qt. \$4.39; 1 gal. \$10.60.

MALATHION. 25% Wettable Powder. Use 2 lbs. per 100 gals. water. 1 lb. 60c; 5 lbs. \$2.75; 50 lbs. \$25.50.

"MARLATE" 50—A residual insecticide with exceptionally low toxicity to animals and humans. There is only a minimum hazard from the spray residue. Especially recommended for worm control on cucumbers, squash, and melons-will not harm plants. 4-lb. bag \$3.65; 50-lb bag \$39.00.

"MARLATE" 2-MR—(Emulsifiable)—residual insecticide for control of leafhoppers and other pasture insects. Also used for chinch bugs, fleas, soil insects and lawns, and insect and fly control of beef cattle as spray or in "back-rubber." 1 gal. \$5.15; 5 gals. \$22.20; 30 gals. \$127.20.

MARLATE DUST NO. 2 (Code No. W-2) (Contains 10% Methoxychlor). 1 lb. 40c; 25 lbs. \$5.05; 50 lbs. \$9.05; 100 lbs. \$18.10.

OIL EMULSION N-P 90. For the control of scales, mealy bugs, red spiders, etc. Dilute with water 3 qts. to 50 gals. Gal. 80c; 5 gals. \$3.25; 54 gals. \$22.00. Deposit on containers.

PARATHION PREPARATIONS. Parathion products are highly toxic to man and warm blooded animals. It is urged that all precautions and recommendations stated on labels be carefully followed.

PARATHION WETTABLE 15%. A wettable powder containing 15% parathion. Use at the rate of 1 lb. to 100 gals. of water. 1 lb. 45c; 3-lb bag \$1.08; 4-lb. bag \$1.44; 5-lb bag \$1.80; 100-lb drum \$34.00.

PARATHION 4 FLOWABLE. 1 gal. \$8.25; 5 gals. \$34.75; 30 gals. \$205.50.

ORTHOPHOS 4 SPRAY. (Contains 4 lbs. parathion per gal.) Use 1/3 pt. per 100 gallons water. 1 gal. \$15.25; 5 gals. \$75.00.

KILPHOS DUST NO. 1. (Code No. E-1)—Contains 1% parathion. 50 lbs. \$2.35; 100 lbs. \$4.70. KILPHOS-SULPHUR DUST NO. 3 (Code No.

ES-3)—Contains 1% parathion and wettable sulphur. 50 lbs. \$3.15; 100 lbs. \$6.30.

KILPHOS DUST NO. 4 (Code No. E-4)—Contains 2% parathion. 50 lbs. \$3.30; 100 lbs. \$6.60.

KILPHOS-KILPHENE-SULPHUR DUST NO. 2 (Code No. EIS-2)—Containing 11/2% parathion and 5% toxaphene. 50 lbs. \$4.60; 100 lbs. \$9.20.

PARIS GREEN. 1/4 lb. 45c; 1 lb. \$1.15; 5 lbs. \$4.50.

PERTHANE WP-50. For the control of chewing insects on lettuce, spinach, and cole crops. Use 2 lbs. to 100 gals. water. 4 lbs. \$2.80; 50 lbs. \$31.50.

PERTHANE EC (EMULSION). (Contains 4 lbs. per gal.) 5 gals. \$27.25; 30 gals. \$154.50.

PHOSDRIN EM-2. (2 lbs. per gallon). A new organic systemic phosphatic poison. Controls aphids, thrips, and certain caterpillars on cole crops. Use 11/2 pts. per 100 gals. 1 gal. \$18.50.

POISON BAITS. Ready for use to control crickets, cutworms, grasshoppers and field crickets. Also ideal for airplane application. Do not use poison baits No. 10 and 11 on cucumbers, squash and melons.

POISON BAIT NO. 8 (Code No. B-8)—(Contains chlordane). 1 lb. 20c; 25 lbs. \$2.50; 50 lbs. \$3.95; 100 lbs. \$7.90.

POISON BAIT NO. 10. (Code No. B-10)—(Contains toxaphene). 1 lb. 15c; 25 lbs. \$2.20; 50 lbs. \$3.35; 100 lbs. \$6.70.

POISON BAIT NO. 11. (Code No. B-11)—(Contains chlordane and toxaphene). 1 lb. 20c; 25 lbs. \$2.70; 50 lbs. \$4.40; 100 lbs. \$8.80.

RED ARROW GARDEN SPRAY. Safe to use on vegetables up to harvest. 1 oz. 49c; 4 oz. \$1.19; pint \$3.49; gal. \$25.00.

ROTHANE PREPARATIONS (TDE)
ROTHANE WP-50. Contains 50% of the technical material. Use at the rate of 2 lbs. to 100 gals. water. 4-lb. bag \$2.44; 50-lb. bag \$28.50.

RHOTHANE EMULSION. (Contains 25% of the technical material.) 5 gals. \$14.75; 30-gal. drum \$84.00; 55-gal. drum \$148.50.

ROTENONE DUST NO. 1. (Code No. R-6)—Safe to use up to harvest on vegetable crops for control of worms, aphids, thrips, etc. 1 lb. 25c; 25 lbs. \$3.30; 50 lbs. \$5.60; 100 lbs. \$11.20.

SYSTOX-A systemic insecticide (which acts from within the plant) effective against aphids and red spider mites. 1-gal can \$32.00.

TEDION 25-W. An ovacide and miticide for use on citrus. 21/2-lb bag \$4.15; 5-lb bags \$8.25.

THIODAN EC-2. For control of aphids, leafhoppers, and potato beetles. Use 1 qt. per 100 gallons. 1 gal. \$10.20; 5 gals. \$49.75.

TOXAPHENE PREPARATIONS

TOXAPHENE 40W. (Contains 40% toxaphene). Use from 3 to 4 lbs. per 100 gals. of water. 6-lb. bag \$1.50; 50-lb. bag \$10.50.

TOXAPHENE SPRAY NO. 4.8. (Contains 4.8 lbs. toxaphene per gal.) Use from 1 to 1½ qts. per 100 gals. of water. Gal. \$3.95; 5 gals. \$18.00.

TOXAPHENE 6 SPRAY. (Contains 6 lbs. Toxaphene per gal.) Use 1 qt. per 100 gals. water. 1 gal. \$4.10; 5 gals. \$19.00; 53 gals. \$190.80.

KILPHENE 5-DUST. (Code No. I-3)—(Contains 5% toxaphene.) 1 lb. 15c; 25 lbs \$1.70; 50 lbs. \$2.40; 100 lbs. \$4.80.

KILPHENE 10-DUST. (Code No. I-2)—(Contains 10% toxaphene.) Lb. 15c; 25 lbs. \$2.20; 50 lbs. \$3.40; 100 lbs. \$6.80.

KILPHENE 5-SULPHUR DUST. (Code No. IS-2)—(Contains 5% toxaphene.) Lb. 15c; 25 lbs. \$2.15; 50 lbs. \$3.25; 100 lbs. \$6.50.

TEPP PREPARATIONS

VAPOTONE XX SPRAY (20%) for control of aphids, thrips, red spiders. Use from ½ to 1 pt. per 100 gals. water. 1 gal. \$12.00; 5 gals. \$58.50.

ORTHO TEPP SPRAY (40%). Use ½ to 2/3 pt. per 100 gals. water per acre. For control aphids, thrips, red spiders and leaf hoppers. 1 gal. \$12.00; 5 gals. \$58.75.

TRITHION 25-W. An insecticide and miticide for use on fruit & vegetables. 5-lb. bag \$7.50; 50-lb. drum

TRITHION 2-E. An excellent chinch bug control. Use 2/3 pint in sufficient water to cover 1000 sq. ft. 1 qt. \$4.95; gal. \$18.00.

TRITHION 4 FLOWABLE. 1 gal. \$23.60; 5 gal. \$116.75; 30 gal. \$697.00; 55 gal. \$1,273.25.

VOLCK OIL SPRAY. A liquid oil spray and spray carrier, furnished in the following sizes only, 4 oz. 39c; 16 oz. 79c; 32 oz. \$1.49; 1 gal. \$3.19.

VOLCK SOLUBLE OIL 97%. A new, improved oil spray, primarily for use on Citrus. Use ½ gal. to 50 gals. water. 1 gal. \$1.25; 54 gals. \$36.18. \$3.00 deposit on 54-gal. containers.

FUNGICIDES

ANTIBIOTIC SPRAYS



Agrimycin 100 is now being used successfully for the control of bacterial spot on tomatoes and peppers, blackleg and soft rot of potatoes, wildfire and blue mold of tobacco. Experimental work is being continued on many other plants. Agrimycin 100 gives best results when used at 100 ppm in combination with neutral copper for diseases on celery, pepper and tomatoes. General recommenda-

tion is 100 ppm. One 9.34-oz. bottle of Agrimycin 100 in 100 gallons of water makes a 100 ppm spray.

AGRIMYCIN 100—9.34-oz. bottle \$6.60; 25-lb. drum (equivalent to $42\frac{1}{2}$ bottles) \$219.00.

AGRIMYCIN 500. A combination of neutral copper and Agri-Mycin 100. 5 lbs. of Agri-Mycin 500 is equivalent to 100 ppm Agri-Mycin and 4 lbs. Neutral Copper in 100 gals. 5 lbs. \$7.60; 25 lbs. \$37.00.

ARASAN-75. (75% Thiram). An effective treatment for vegetable seed. Reduces loss from seed decay and damping-off. 1-oz. pkg. 60c; 8-oz. can \$1.30; 25-lb drum \$40.30; 100-lb. drum \$156.25.

CERASAN (New Improved). Destroys seed-borne diseases by direct contact with the spores or penetrating vapor fumes. Ceresan has been used successfully in treating Gladiolus bulbs for control of Fusarium Rot before planting. Use ½ lb. Ceresan, 5 tablespoons of DuPont Spreader-Sticker to 25 gallons water. Dip bulbs in solution, let remain 15 minutes. Bulbs can be planted immediately after treatment. 1 lb. \$1.95; 100 lbs. \$110.00.

COPPER COMPOUNDS

BORDEAUX MIXTURE 1 lb. 98c; 4 lbs. \$2.00.

BLUESTONE (copper sulphate) Granular or Snowform. 1 to 10 lbs. 25c per lb.; 25 lbs. \$5.00; 50 lbs. \$9.50; 100 lbs. \$18.00.

COPPER-A COMPOUND. — An excellent Copper fungicide for spraying purposes containing 45% metallic copper. Formulated especially for greater safety on copper sensitive plants. Use 4 lbs. per 100 gals. of water. 6-lb. bag \$2.97; 50-lb. bag \$23.70.

KILCOP DUST NO. 1 (Code No. C-1)—5.3% metallic copper). Lb. 15c; 25 lbs. \$2.35; 50 lbs. \$3.70; 100 lbs. \$7.40.

KILCOP DUST NO. 2 (Code No. C-2)—(6.89% metallic copper.) Lb. 20c; 25 lbs. \$2.70; 50 lbs. \$4.35; 100 lbs. \$8.70.

KILCOP-SULPHUR DUST NO. 1. (Code No. CS-1) — (6.75% metallic copper.) Lb. 20c; 25 lbs. \$2.95; 50 lbs. \$4.90; 100 lbs. \$9.80.

KILCOP "53." (Contains 53% metallic copper.) Kilcop is extremely fine, which is essential for good coverage and effective control of fungus. Use 4 lbs. per 100 gals. water. 1-lb. can 85c; 2-lb. bag 90c; 6-lb. bag \$2.46; 50-lb. bag \$19.50.

KILCOP DUST NO. 3 (Code No. C-3)—(7.4% metallic copper.) Lb. 20c; 25 lbs. \$2.80; 50 lbs. \$4.55; 100 lbs. \$9.10.

KILCOP DUST NO. 4 (Code No. C-4)—(9% metallic copper.) Lb. 20c; 25 lbs. \$3.00; 50 lbs. \$5.00; 100 lbs. \$10.00.

DITHANE PREPARATIONS



DITHANE D-14 (Nabam). This fungicide has given excellent control on a variety of crop diseases such as late and early blight of potatoes and tomatoes, leaf blights of peppers, curvularia leaf spot of glads, anthracnose of beans and downy mildew of cucumbers, squash and melons and helminthosporium leaf blight of corn. Use 2 qts. Dithane D-14, ¾ lb. zinc sulphate to 100 gals. water. 1 gal. \$2.10; 5 gals. \$8.75; 30 gals. \$49.50.

DITHANE Z-78 (Zineb) (Wettable). Effective in the control of down mildew and anthracnose of cucumbers, squash, melons and blue mold on tobacco. Helminthosporium leaf blight of corn. Use 2 lbs. to 100 gals. 1-lb. canister \$1.50; 3-lb. bag \$2.46; 5-lb. bag \$4.10; 50-lb. bag \$39.50.

DITHANE M-22 (Maneb). Controls a number of diseases on tomatoes, celery, peppers, eggplant and potatoes. Particularly effective where manganese deficiency exists. Use 1½ lbs. per 100 gals, water. 3-lb. bag \$3.36; 4½-lb. bag \$5.04; 50-lb. drum \$55.00.

MANEB DUST NO. 1 (Code No. Ma-1)—(Contains 5.6% Maneb). Lb. 25c; 25 lbs. \$3.50; 50 lbs. \$6.00; 100 lbs. \$12.00.

ZINEB DUST NO. 1 (Code No. M-1)—(Contains 4% Zineb). Lb. 20c; 25 lbs. \$2.55; 50 lbs. \$4.10; 100 lbs. \$8.20.

ZINEB DUST NO. 2 (Code No. M-2)—(Contains 6% Zineb). Lb. 25c; 25 lbs. \$3.15; 50 lbs. \$5.25; 100 \$10.50.

ZINEB-SULPHUR DUST NO. 1 (Code No. MS-1)— (Contains 4% Zineb). Lb. 20c; 25 lbs. \$3.00; 50 lbs. \$4.95; 100 lbs. \$9.90.

DYRENE. A fungicide for the control of gray leaf spot and botrytis on tomatoes and early blight on celery. 50-lb. drums \$92.50.

FERMATE® (Ferbam) Fungicide for control of tobacco blue mold, early and late blight of celery; downy mildew of cabbage, damping-off in celery and lettuce seedbeds. Use 2 lbs. per 100 gals. water. 3-lb. bag \$1.77; 50-lb. bag \$27.50.

FUNGICIDE 658. Copper, zinc, chromate complex. A fungicide for tomatoes, potatoes, celery and pepper. Use 2 lbs. per 100 gals. 4 lbs. \$3.12; 6 lbs. \$4.68; 30 lbs. \$22.50.

KARATHANE 25-W. A specific fungicide for the control of powdery mildew on cantaloupes, cucumbers, squash and ornamentals. Use ½ lb. to ¾ lb. per 100 gals. water. 3-lb. bag \$5.10.

KARATHANE DUST NO. 1 (Code No. Z-1). 1 lb. 20c; 25 lbs. \$2.80; 50 lbs. \$4.60; 100 lbs. \$9.20.

MANZATE®—(Maneb) for control of tomato diseases such as late blight and gray leaf spot. Provides protection without yellowing or stunting of plants.

Also controls major foliage diseases of gladioli, cucumbers, potatoes, celery, and other crops. Use 1½ lbs. per 100 gallons of water. 3-lb. bag \$3.36; 4½-lb. bag \$5.04; 50-lb. drum \$55.00.

ORTHOCIDE 50-W (Captan). For control of damping-off in pepper seedbeds and other fungus diseases. For Irish potato seed treatment see page 52. Use 4 lbs. per 100 gals. water. 8-oz. canister \$1.39; 4-lb. bag \$3.04.

ORTHOCIDE 75 SEED PROTECTANT. Contains 75% captan. A seed treatment to prevent seed rot and damp-off diseases. 4½ oz. shaker can \$1.19; 20-lb. drum \$33.00.

ORTHO SEEDGUARD. A combination insecticide and fungicide to protect germinating seeds against soil insects and damping-off fungi. 20-lb. drum \$83.00.



FUNGICIDES (Continued)

PARZATE®—Zineb (Wettable)—Effective in the control of downy mildew, and anthracnose of cucumbers, cantaloupes, squash, and watermelons.

For control of helminthosporium leaf blight of corn, early and late blight of potatoes, blights and gray leaf spot of tomatoes, blue mold of tobacco, and leaf spots of many other crops. Use 2 lbs. per 100 gals. of water. 3-lb. bag \$2.46; 5-lb. bag \$4.10; 50-lb. bag \$39.50.

PARZATE®—Nabam—Used as an effective fungicide wherever Nabam or Zineb is recommended. Especially useful for potato blights and helminthosporium leaf blight of corn. Use 2 qts. "Parzate" and ¾ lb. zinc sulphate per 100 gals. 5-gal can \$8.75; 30-gal drum \$49.50.

PHYGON-XL (Wettable). A fungicide for the control of leaf blights of tomatoes, beans and various other plant diseases. Use ½ to 1 lb. per 100 gals. of water. 1 lb. \$2.50; 4 lbs. \$8.20; 100 lbs. \$200.00.

SEMESAN. An effective treatment for vegetable and flower seeds to control damping-off fungi in the soil. Also repels ants. Directions on package. 4-oz. pkg. \$2.50; 12-ozs. \$5.95; 25 lbs. \$105.00.

SEMESAN BEL. For treatment of potatoes, controlling certain diseases of this crop. Directions on package. 3 oz. \$1.45; 1 lb. \$3.30; 4 lbs. \$10.65; 40 lbs. \$99.80.

SPERGON. Very valuable for treating seeds, especially beans and peas, to control soil diseases for better stands and increased yields. Use 2 ounces per bushel of seed with beans and peas. 1 lb. \$2.40; 50 lbs. \$100.00.

SPERGON (Wettable). Valuable for spraying to control diseases—especially downy mildew in cabbage seedbeds. Use at the rate of 2 lbs. to 100 gals. of water. 1 lb. \$2.00; 5 lbs. \$6.25; 50 lbs. \$57.50.

SPERGON DUST NO. 3 (Code No. Y-3)—Contains 5% Spergon—Wettable.) For the control of downy mildew on cabbage. 25 lbs. \$3.45; 50 lbs. \$5.90; 100 lbs. \$11.80.

SULFOCIDE. A vegetable spray which is also valuable for citrus fruits. Checks mildew, rust and anthracnose on beans. Also used in washing water before shipping beans, tomatoes, eggplants and peppers to check rot and other diseases while in transit. Use 2 qts. to 100 gals. of water. ½ pt. 75c; pt. \$1.25; qt. \$1.75; gal. \$4.00; 5 gals. \$15.00.

SULPHUR (Dusting). Lb. 15c; 25 lbs. \$1.30; 50 lbs. \$1.60; 100 lbs. \$3.15; ton \$62.00.

SULPHUR (Wettable). 50 lbs. \$1.70; 100 lbs. \$3.35; ton \$65.50.

TERRACLOR WP 75% (PCNB). For the control of certain soil-borne diseases. Use 5 lbs. per 100 gals. 5-lb. bag \$4.20; 50-lb. bag \$40.50.

TERSAN® 75—A lawn and turf fungicide for control of "brown-patch" and "dollar spot." Use two table-spoonfuls to $1\frac{1}{2}$ gal. water to 60 sq. ft. 8-oz. cans \$1.95; 3-lb. bag \$5.50.

THYLATE 65W—A fungicide used primarily for the control of celery seedbed diseases. Use $1\frac{1}{2}$ lbs. per 100 gals. water. 5-lb. bag \$4.10.

ZERLATE® Ziram—Specific for control of "frog-eye" spot on peppers and leaf blights on celery. Use two pounds per 100 gals. 3-lb bag \$2.22; 50-lb. bag \$35.00.

Insecticide and Fungicide Combinations



Prices quoted are f.o.b. Plant City or Kilgore stores

Prices subject to change without notice

The Kilgore Seed Company processes many combinations of insecticides and fungicides which are too numerous to list in our General Catalog. If you do not find the combination you want, write our General Office in Plant City or call at your nearest Kilgore Store and it is likely we will have a dust in stock which will fill your requirements.

KILCIDE-KILCOP DUST NO. 1 (Code No. DC-1)—(Contains 3% DDT plus 5.3% metallic copper.) Use where DDT is needed with a neutral copper. Lb. 20c; 25 lbs. \$2.65; 50 lbs. \$4.30; 100 lbs. \$8.60.

KILCIDE-KILCOP DUST NO. 2 (Code DC-2)—(Containing 5% DDT, 7.4% metallic copper, a combination dust for control of insects and diseases. 1 lb. 25c; 25 lbs. \$3.20; 50 lbs. \$5.35; 100 lbs. \$10.70.

KILCIDE-ZINEB DUST NO. 4 (Code No. DM-4)— (Contains 5% DDT and 6½% Zineb.) For use where DDT is desirable for general insect control with Zineb. 1 lb. 30c; 25 lbs. \$3.75; 50 lbs. \$6.50; 100 lbs. \$13.00.

KILHEX—KILCOP DUST NO. 2 (Code No. GC-2)— (Contains 1½% gamma isomer and 6.89% metallic copper.) A practical insect and disease control combination at a minimum cost for cabbage, celery, tomatoes, peppers and eggplants, for application before fruits or edible portions are well developed. 1 lb. 25c; 25 lbs. \$3.15; 50 lbs. \$5.30; 100 lbs. \$10.60.

KILGAM-KILCOP DUST NO. 1 (Code No. QC-1)— (Contains 1½% lindane [essentially pure gamma isomers of benzene hexachloridel and 5.3% metallic copper.) An excellent combination designed for use on cucumbers, squash, and melons, as well as for cabbage, celery, tomatoes, peppers and eggplants, 1 lb. 30c; 25 lbs. \$4.10; 50 lbs. \$7.15; 100 lbs. \$14.30.

KILPHENE-KILCOP DUST NO. 2. (Code No. IC-2)—A convenient insect and disease control combination containing 10% toxaphene and 6.36% metallic copper. (Under no circumstances use this material on cucumbers, squash, or melons.) 1 lb. 25c; 25 lbs. \$3.45; 50 lbs. \$5.85; 100 lbs. \$11.70.

KILPHOS-ZINEB DUST NO. 1. (Code No. EM-1)— (Contains 1% Parathion and 6% Zineb). Can be used on most crops for most insect and disease control. 50 lbs. \$6.15; 100 lbs. \$12.30.

KRYOCIDE-KILCOP DUST NO. 2. (Code No. KC-2)—(Contains 27% Kryocide plus 5.3% metallic copper.) A good insecticide and fungicide combination for most vegetable crops. Lb. 30c; 25 lbs. \$3.85; 50 lbs. \$6.65; 100 lbs. \$13.30.

ZINEB KILGAM DUST NO. 3. (Code No. MQ-3)— (Contains 1½% Lindane and 6% Zineb.) This combination is recommended for purposes where Zineb has been proved more effective than copper. Refer to Kilgam-Kilcop Dust No. 1 for further important details. 1 lb. 35c; 25 lbs. \$4.70; 50 lbs. \$8.40; 100 lbs. \$16.80.

ZINEB-KRYOCIDE DUST. (Code No. MK-1)—(Contains 4% Zineb and 27% Kryocide.) This combination is recommended for purposes where Zineb has been proved more effective than copper. Lb. 30c; 25 lbs. \$4.15; 50 lbs. \$7.25; 100 lbs. \$14.50.



SOIL FUMIGANTS

Nematodes are present in most of our soils, and seriously affect most of our vegetable crops and reduce yields. Nematodes can be controlled by the proper use of soil fumigants, either solid treatment or row treatment, which is applied in the row that is to be planted. This method has become very popular, especially with small growers, as cheap handmade applicators with gravity flow can be used.

The cost of this method of treatment is only about one-third the cost of solid treatment, and has proved very satisfactory. "D-D" and Dowfume W85 are the materials most extensively used in commercial treatment at this time. Information on soil fumigation can be obtained from any of the Kilgore stores or write The Kilgore

Seed Company, Plant City, Florida.

D-D Soil Fumigant. Use 30 gals. per acre broadcast or 8 gals. per acre in 4-foot rows. Gallon \$2.00; 30-gal. drum \$45.00; 54-gal drum \$81.00. Not mailable.

KILFUME 80-20 (EDB). Use 48 gals. per acre broadcast or 12 gals. per acre in 4-foot rows. 1 gal. \$1.65; 30-gal. drum \$46.50; 50-gal. drum \$75.00.

KILFUME 60-40 (EDB). Use 23 gals. per acre broadcast or 6 gals. per acre in 4-ft rows. 5 gals. \$14.70; 50 gals. \$117.50.

DOWFUME W-85 (EDB). Use 6 or 7 gals. per acre broadcast or 1¾ gals. per acre in 4-foot rows. 5 gals. \$30.00; 30 gals. \$175.50.

V-C 13 NEMACIDE. For the control of nematodes affecting established plants. Use 1 pt. in 50 gals. water to 180 sq. ft. For control of chinch bugs use 1 pt. to 1250 sq. ft. 8 oz. bottle \$1.75; pt. \$3.25; qt. 6.00; 1 gal. \$20.00; 50 lbs. (5 gals.) \$75.00.

NEMA-KIL® 17.5 G (A Soil Fumigant in Granules). Is a nematocide especially effective against root knot nematodes. NEMA-KIL 17.5 G is effective as a preplanting application for cotton, strawberries, certain vegetables, nurseries, and ornamentals. Also useful as Post-Planting or side-dressing applications for cotton, carrots, perennials, strawberries and certain ornamental shrubs such as boxwood, etc. Useful in relatively dry soils. NEMA-KIL 17.5 G has long residual qualities. Waiting period between application and planting—2 to 3 weeks. 1 lb. 98c; 5 lbs. \$3.95. KILGORE'S NEMA-KIL (Contains Nemagon). nematocide which kills nematodes in the soil. Can be used around many established plants. See page 77 for more detailed information. Pt. \$1.35; qt. \$2.25; gal. \$6.75; 4 gals. \$26.60; 30 gals. \$165.00.

VPM SOIL FUMIGANT. For control of certain weeds, soil-borne fungus diseases, nematodes and garden centipedes, etc. Cultivate soil to promote good penetration of the liquid and maximum distribution of vapors. Do not apply to dry soil, or when soil temperature is less than 60°F. Best results are obtained when the liquid is injected into soil to a depth of 6-10 inches, directly under the row, covered and packed immediately. Apply at rate of 1 pint per 100 lineal feet of row. Allow 14 days before seeding or transplanting. 1 gal. \$3.95; 5-gal. drum \$18.00; 30-gal. drum \$96.00.

METHYL BROMIDE (with 2% chloropicrin). For treatment of nursery seedbeds and greenhouse soil. 1-lb. can 73c.

JIFFY APPLICATORS. For 1-lb. methyl bromide cans \$3.35; with 10' tubing \$5.25.

NOTE—For plastic covers used in treating seedbeds with methyl bromide, see page 82.

GENERAL FUMIGANTS

CARBON DI-SULPHIDE (Hi-life). Recommended for ridding grain of weevils, and killing ants, etc., and for destroying vermin. Sold in gallon can only. Per gallon \$2.25; 5 gals. \$11.25. (Not mailable.)

CYANOGAS ANT-KILLER. Cyanogas Ant-Killer kills ants with the least trouble. Not a bait. It's the gas that

kills 'em. 4-oz. can 50c. (Not mailable.)

CYANOGAS A-DUST. For killing rats in lumber piles; on poultry farms; in grain elevators and stockyards. Mice (field mice and pine mice), moles, gophers. 1-lb. tin \$1.25; 5-lb. can \$4.25; 25-lb. can \$13.75. (Not mailable.)

CYANOGAS G-FUMIGANT. Comparable to sea sand in size of particles. Used for fumigating greenhouses, warehouses, buildings and for grain fumigation. 4-oz. can 50c; 5-lb. can \$4.25; 25-lb. can \$13.75; 100-lb. drum \$43.35. (Not mailable.)

PYRENONE, GRAIN PROTECTANT. A contact material for the protection of corn and other grains stored in open cribs or bins. Offers freedom from toxic hazards to man and animals. Effective against weevils and other storage insects. Prices: 50-lb. bag \$12.50, f.o.b. Plant City.

VERMICIDES RODENTICIDES AND

CARBOLINEUM. A wood preservative and vermicide. Prevents damage by termites. Kills and protects against chicken mites and fowl ticks. ½-gal. can \$1.80; 1 gal. \$2.60; 5 gals. \$12.00; 35 gals. \$80.50; 55 gals. \$121.00; f.o.b. Plant City.

COLONIAL FLEA and TICK KILLER. For cats and

dogs. For dogs 4½-oz. shaker-top can \$89c; 1-lb. can \$1.50; for dip 4-oz. bottle \$1.50.

For cats 4½-oz. can 89c.

COLONIAL-42.—An effective bait containing Warfarin for the control of all species of rats and mice. 4-oz. pkg. 35c; 10 oz. 60c; 1 lb. 90c; 2 lbs. \$1.65.

D-CON. A ready mixed, ready to use rat and mouse bait. Contains Warfarin. For indoor and outdoor use. 1 lb. \$1.69; 3 lbs. \$3.95.

DIXIE KILLER. Exterminates rats, mice, roaches, waterbugs, land crabs, cutworms and other destroyers of food and property. 2-oz. jar 35c; 1 lb. \$1.25; 2 lbs. \$2.00. Prepaid.

GATOR ROACH HIVES. Small size (3 baits) 20c;

large size (3 baits) 40c. Prepaid.

HARRIS ROACH TABLETS. 2-oz. pkg. 30c; 4-oz. pkg. 55c.

KILGORE'S RAT AND MOUSE BAIT. A Warfarin bait especially designed for the control of rats and mice in fields and groves. 10 lbs. \$3.50; 25 lbs. \$8.00; 50 lbs. \$15.00.

MAGIKIL JELLY ANT BAIT. Kills ants and is harmless to plants. Ant trap 25c; Garden size 40c; Estate tube \$1.00; 7½-oz. Jumbo size \$1.95; 10 lbs. \$23.00. Prepaid.

MELON SEED TREATMENT. This treatment will effectively protect your vine seeds from birds and rodents as well as acting as an insecticide, and fungicide. 5-lb. kit (treats 5 lbs. seed) \$2.50; 10-lb. kit \$4.00; 20-lb. kit \$7.00.

PESTEX. A cresol repellent used as a protectant of seed from birds and insects. 1 lb. 40c.

TERRO ANT KILLER. Will rid your place of ants quickly. 1¼-oz. bottle 39c; 3-oz. bottle 59c. Prepaid.

ZINC PHOSPHIDE. A sure poison for field rats. To 40 lbs. of scratch feed add 4 oz. mineral oil, then add 4 oz. Zinc Phosphide. Place small quantities at 15 to 20 ft. intervals. 1-lb. can \$2.95. (Not mailable.)

Spreaders, Stickers, Emulsifiers, Etc.

DU PONT SPREADER-STICKER. Use 1 to 4 oz. per 100 gals. Oz. 59c; pt. \$1.40; gal. \$8.15; 5 gals. \$39.00.

ORTHO SPRAY STICKER—½ pt. to 100 gals. Gal. \$3.75; 5 gals. \$17.50.

PLYAC 25%. Emulsifiable polyethylene wax. ½ to 1 pint per 100 gals. Gal. \$8.25; 5 gals. \$39.25.

TRITON B-1956. An efficient emulsifier and spreader for insecticides and fungicides. A combined spreader, sticker and deposit builder. Soluble in water and oil. Use at the rate of 2 to 4 ounces per 100 gallons. Gal. \$7.70; 5 gals. \$37.75; 55 gals. \$407.00.

TRITON X100—An excellent spreader and wetting agent. Particularly good with the newer hydrocarbon insecticides. 1 gal. \$6.50; 5 gals. \$26.00; 55 gals. \$217.50.

WEED KILLERS

Note: Read labels carefully—they are put on the package for your protection. Use Weed Killers wisely and you will be using them safely. Be sure to clean spray equipment thoroughly according to approved methods after using weed control materials of any kind. f.o.b. Plant City.

AMMATE®—For complete kill of undesirable brush, trees, poison oak, poison ivy, and many other types of weeds "Ammate" provides the safest way to control these weeds that may adjoin cropland. There are no vapors to drift onto sensitive crops or foliage. "Ammate" is non-volatile, non-inflammable and can be used without hazard to humans or animals. Use 60-75 lbs. per 100 gals. plus Spreader-Sticker at 4 oz. per 100 gal. 2-lb. bag \$1.35; 6-lb. bag \$3.50.

DOWPON—85% Sodium Salt. This material is absorbed through the foliage killing as it travels through the plant, and so is not affected by weather. For the control of all perennial grasses, use 5-10 lbs. per acre in sufficient water to cover thoroughly. Repeat application in two weeks and disc thoroughly after another two week interval. Grass should be treated before seed has formed. Wait at least 30 days from last application date before planting. 1 lb. \$2.33; 5 lbs. \$6.85; 50-lb. drum \$57.50.

KARMEX® (Diuron). For selective weed control in such crops as sugar cane, pineapple, gladiolus and other bulb ornamentals. Dosage rates vary from 1-4 lbs. per acre, depending upon the crop treated.

Use at 1 lb./1000 sq. ft. wherever it is desired to kill all vegetation for a prolonged period of time (9-12 mos.)—ditches, around storage yards, barns, packing houses, fence rows, etc.

Do not apply on or near valuable trees or other plants or on areas into which their roots may extend. Do not use on lawns, walks, driveways, tennis courts, or similar areas.

2-lb. can \$9.00; 10-lb. drum \$38.50; 50-lb. drum \$175.00.

RANDOX. A selective herbicide for the control of weeds in beans, cabbage, celery, corn and onions. Consult the Experiment Station in your area for specific recommendations. 1 gal. \$12.60; 1 case (4 1-gal. cans) or 1 5-gal. can, \$12.10 per gal.; 15-gal. drum or 3 5-gal. cans, \$11.60 per gal.

SODIUM ARSENITE—8 lbs. per gal. A highly concentrated, black opaque viscous Sodium Arsenite solution. A non-selective weed killer used for killing weeds on ditch banks, fence rows, etc. Is especially recommended for killing off potato tops before harvest. Permits earlier harvest, toughens the skin and decreases mechanical injury. Use 1 gallon of Sodium Arsenite to 100 gallons of water, applied at the rate of 100 to 150 gallons per acre. Comply with cautions on label. 15-gal. drums only \$30.00.

POLYBOR-CHLORATE. A non-selective herbicide for the prolonged control of weeds about brick walks and patios, fence lines, driveways, or tennis courts, around stores, industrial properties, etc. It is non-selective in its action and will kill any plant with which it comes

in contact. Dissolve four pounds in one or two gallons of water to be applied to 100 sq. ft. Price: 5-lb. pkg. \$2.25; 50-lb. bag \$10.25.

TCA (90% Sodium TCA). A grass killer for nonselective control in areas not planted to crops. For maximum control of Para grass along ditch banks. Several applications of TCA in small amounts will produce better results than one application at a higher dosage. 10 lbs. \$5.25; 50 lbs. \$22.85; 100 lbs. \$44.25.

"TELVAR"® (Monuron). An effective general herbicide for the control of many annual and perennial grasses and herbaceous weeds on non-crop land areas where bare ground is desired. The degree of control and duration of effect will vary with the amount of chemical applied. Use at the rate of 20 to 80 lbs. per acre. 2 lbs. \$8.60; 10 lbs. \$38.00; 50 lbs. \$175.00.

TRIOX. An arsenical weed killer. Toxic to the soil and kills vegetation. Prevents plant growth for one to two years. A liquid, easy to apply with watering can. Do not use on lawns. Consult label. Qt. \$1.19; gal. \$3.50; 5 gals. \$13.50; 30-gal. drum \$72.00.

VEGADEX—A highly effective weed killer (Kills Annual Bluegrass, Bullgrass [Goose Grass], Crabgrass, Green Foxtail, Yellow Foxtail, Giant Foxtail, Pigweed [Careless Weed], Barnyard Grass [Water Grass], Henbit [Blueweed], and Purslane), for use on Collards, Mustard Greens, Turnip Greens, Broccoli, Cabbage, Snap Beans, Soybeans, Celery, Sweet Corn, Kale, Spinach, Hanover Salad, Brussels Sprouts, Cauliflower, Lima Beans, Garden Beets, Lettuce, and Field Corn. This herbicide works through the soil killing germinating weeds when the roots absorb the chemical, leaving your crop unharmed. CONSULT LABEL AND LOCAL AUTHORITIES FOR SPECIFIC RECOMMENDATIONS ON YOUR CROP IN YOUR AREA. 1 gal. \$14.50; 1 case (4 1-gallon cans) or 1 5-gallon cans, \$13.50 per gallon.

WEED-B-GON—64. An amine salt of 2, 4-D. Excellent for killing water hyacinth and most broad leaf plants. Apply as per directions on container. Caution—do not use on or let material drift on plants you do not want to kill. Price, 1 gal. \$4.35; 5 gals. \$20.75; 30 gals. \$121.50; 50 gals. \$197.50.

WEEDONE BRUSH KILLER—32. Contains 2, 4, 5-T, plus 2,4-D. Kills poison ivy, poison oak stumps and other woody plants. Use 1 to 1½ gals. to 100 gals. water or ½ to 1 pt. to 5 gals. water. Caution—do not use this material on plants you do not want to kill. Price: qt. \$3.75; gal. \$7.25; 5 gals. \$34.50.

NEW IMPROVED WEEDONE. A selective weed killer containing 2,4-D plus 2,4,5-T. For complete kill, all the foliage must be sprayed as there is little crossmovement within the stems. Do not use Weedone on St. Augustine type grasses. 8 oz. can \$1.00; 1-qt. can \$2.75; 1-gal. can \$6.75; 5-gals. \$24.75.

A number of pre-emergence and post-emergence weed control materials are now being offered to truck crop growers. As most of these materials have not been tried sufficiently under Florida conditions we hesitate to make specific recommendations in this catalog. We suggest that growers interested should consult local authorities or a Kilgore representative.

WE WILL MAKE EVERY EFFORT TO SUPPLY WEED CONTROL MATERIALS OF THIS NATURE RECOMMENDED BY THE FLORIDA EXPERIMENT STATION.

LIVESTOCK PREPARATIONS

Prices are subject to change without notice and are t.o.b. Plant City unless otherwise specified.

DDT 50% Weitable. For spray on range cattle. To control horn flies, lice and mosquitoes use 20 lbs. per 100 gals. water. Do not use on dairy cattle. 1-lb. canister 75c; 3-lb. bag 78c; 50-lb. bag \$11.50.



BHC Wettable. (10% Gamma isomer of BHC). For spray on range cattle to control horn flies, lice and mosquitoes use 3¾ lbs. per 100 gals. water. Do not use on dairy cattle. 5-lb. bag \$1.30; 50 lbs. \$11.00.

"Marlate" 50% Technical Methoxychlor Insecticide Wettable Powder. For spray on range cattle to control flies and lice. For lice use 25 lbs. per 100 gals. water. For flies use 8¾ lbs. per 100 gals. water. 4-lb. bag \$3.62; 50-lb. bag \$39.00.

Cooper-Tox Livestock Spray (45% Toxaphene and 2% BHC). For the control of horn flies, ticks, and lice. Use 1 gal. to 150 gals. water. 1 gal. \$7.10; 5 gals. \$33.75.

Diazinon 25E (25% Emulsifiable Solution.) A 25% emulsifiable solution for use in overall residual fly sprays, "spot" sprays and liquid baits for application in dairy barns and other farm buildings except poultry houses. Gives control of flies, including resistant strains, for periods up to 8 weeks. It is equally effective against fly maggots. Use 1 pint to 6 gallons of water. Consult label for more detailed information. Price—1 Pint Bottle \$3.50; gal. \$18.00.

DuPont Dairy Barn Insecticide (50% Methoxychlor and 3% Lindane) for the control of house flies and table flies in dairy barns, sheds, and stables. Use 1 lb. in 3 gals. water. 1 lb. \$1.60; 4 lbs. \$5.40.

DuPont Dairy Cattle Insecticide. (Methoxychlor, 50% wettable).

As a Spray or Dip—(except dairy animals)—for controlling hornflies, lice and fleas, use 2 to 4 lbs. per 25 gals. of water. As a spray, make sure animals are wetted to the skin; wetting is improved by adding ½-oz. DuPont Spreader-Sticker per 25 gals. of spray. Usually, one treatment will be effective for about 3 weeks.

Direct Dusting—(Dairy Cattle and Sheep)—Use one heaping tablespoonful per animal. Sprinkle on poll, neck (top and sides) and back and upper sides; rub in lightly by hand to carry powder beneath the hair. One treatment protects cattle against hornflies for about 3 weeks. Application on sheep, following shearing, gives temporary relief from sheep ticks (keds).

Farm Buildings — For controlling stable flies houseflies, and fleas, use 1 lb. in 2½ gallons of water. Apply one gallon of spray mixture to approximately 500 sq. ft. of surface—treating walls, floors, cracks, crevices, baseboards, pet bedding, coops, stalls and pens. Such heavy deposits usually are effective for 3 to 6 weeks, unless removed by weathering or other means. The use of any residual fly spray should be supplemented with proper sanitary measures, including spraying or removal of breeding sources such as manure in and around buildings. Both the buildings and the animals should be treated. Price: 1 lb. \$1.32.



Isotox Dairy Spray. (Contains lindane). Spray for killing flies in dairy barns and milk sheds. Do not spray on dairy cows. 1 lb. \$2.19; 4 lbs. \$8.60.

Ortho Healthy Herd. A wettable powder. (Contains 45% DDT and 2.5% gamma isomer of BHC and 12.5% other isomers of BHC.) Use as spray or dip on cattle, horses, hogs, sheep, and goats to control lice, ticks, horn flies and mosquitoes. Use 10 lbs, per 100 gals. water. (Not for use on dairy cattle or in dairy barns.) 5-lb. bag \$3.25; 50-lb. bag \$25.50.



Ortho Kleen Stock. An emulsifiable liquid containing toxaphene and lindane for the control of ticks, lice, flies, etc. on beef cattle and hogs. One pt. makes 4 to 12½ gals. finished spray depending on use. Directions on container. (Not for use in dairy barns or on dairy cattle.) Price 16-oz. bottle \$1.59; 32-oz. bottle \$2.69; 1 gal. \$7.29; 5 gals. \$28.98.

DRY FLY BAITS

For use against flies and their maggots in dairy and livestock barns, stables, poultry houses, feed rooms, kennels, manure piles, etc. Do not apply where poultry can feed upon the poison.

Fly Flakes. For adult flies apply 4 oz. per 100 sq. ft. For maggots, use $1\frac{1}{2}$ lbs. per 100 sq. ft. 1-lb. shaker-top can 98c; 2-lb. 89c; 5-lb. bag \$1.79; 10-lb. bag \$3.25; 25-lb. bag \$6.00.

Ortho Fly Killer Dry Bait. For adult flies use 4 oz. to 500 sq. ft. 1-lb. canister 98c; 5-lb. bag \$2.19.

SCREW WORM PREPARATION

Smearex. ½ pt. 50c; pt. 90c; qt. \$1.75; gal. \$6.00

Ortho 1038 Screw Worm Control. 4-oz. bottle 90c; 16 oz. \$2.75; 32 oz. \$5.00; 1 gal. \$12.00. 11 oz. bomb (Aerosol) \$1.69.

Barry's Screw Worm Bomb (Aerosol). 6-oz. Bomb \$1.25.

Note: Follow label directions on all screw worm preparations.



BIG STINKY OUTDOOR FLY TRAPS

The action of Big Stinky's Control Fluid with water on the starting bait produces a potent scent which lures flies in the vicinity into the trap. Big Stinky actually lures flies away from houses, barns, stock pens and other buildings where you don't want them—it's irresistible—to filthy flies.

1-gal. size with 8 oz. Control fluid	84.95
"Little Stinky" with 3 oz. control powder	2.95
"Wee Stinky" with 1½ oz. control powder	1.69
8-oz. bottle control fluid	1.50
3-oz. tube control powder	.89

POULTRY PREPARATIONS

Germozone. For use in drinking water. Valuable as preventive of simple bowel trouble, also ocular roup. 4 fluid oz. 50c; 12 fluid oz. 90c.

Kilcide 10 Powder. (Containing 10% DDT.) (Code No. D-4.) For treating chickens against lice. Apply by dusting it well into the feathers in order to get the dust down to, or near, the skin surface. 1 lb. 15c; 25 lbs. \$2.25.

FERTILIZERS

All prices quoted are f.o.b. and subject to change without notice.

CAMELLIA AND AZALEA PLANT FOOD

A special prepared fertilizer for camellias and azaleas. Containing the three primary plant foods, plus secondary elements. Price 5 lbs. 70c; 10 lbs. \$1.25; 25 lbs. \$1.95; 100 lbs. \$5.60; f.o.b. Plant City.

GRANULAR AERO CYANAMID. A fertilizer carrying both nitrogen 22% and lime 70%. It is also valuable in helping to control soil disease organisms, especially nematodes. Packed in convenient, even weight, paper lined, 100-lb. sacks. 100 lbs. \$5.00 f.o.b. Plant City; ton \$95.00.

Cyanamid—Lawn and Garden 10-lb. bag \$1.85; 25-lb. bag \$3.75; 50-lb. bag \$6.50

DAIRY ORGANIC COMPOST

High in bacteria. Supplies humus and organic plant food so greatly needed in most Florida soils. Excellent for starting lawns, flowers, shrubs and garden vegetables. Odorless, and will not burn. Weed seed is destroyed in processing.

Use 20 to 25 lbs. per 100 square ft. 5-lb. bag 59c; 25-lb. bag \$1.60; 50-lb. bag \$2.50; ton \$80.00.

PULVERIZED SHEEP MANURE

Sheep manure is highly concentrated, with all weed seeds killed. It supplies the humus and natural plantfood so greatly needed by Florida soils. For lawns, flowers and shrubs use 10 to 12 lbs. per 100 sq. ft. For commercial vegetable growing use 1,000 to 1,500 lbs. per acre drilled in the row. 25 lbs. \$1.00; 50 lbs. \$1.75; 100 lbs. \$3.50; ton \$67.50.

VERTAGREEN is an high quality, scientifically blended plant food, for lawns, flowers, shrubs, trees, vegetables. Contains nitrogen, phosphorous and potash, besides several other important plant food elements. Clean, odorless, easy to apply. Used with outstanding success.

Price per lb. 15c; 5 lbs. 60c; 10 lbs. \$1.10; 25 lbs. \$1.75; 50 lbs. \$2.75; 100 lbs. \$5.15; f.o.b. Plant City, or Kilgore Stores.



VIGORO—Complete balanced plant food supplies the nutrient elements plants must get from the soil for best growth and development. Vigoro is odorless, sanitary and easy to apply. Made for Florida soils and conditions. Use it on everything you grow. 1 lb. 15c; 5 lbs. 60c; 10 lbs. \$1.10; 25 lbs. \$1.75; 50 lbs. \$2.75; 100 lbs. \$5.15; f.o.b. Plant City or any Kilgore store. Write for prices on larger quantities.

SOLUBLE PLANT FOOD

HY-GRO. A soluble plant food containing the three major plant food elements, nitrogen, phosphoric acid and potash, plus minor elements. An excellent starter solution for potted plants, seedbeds and transplanting. Use at the rate of two tablespoonfuls dissolved in one gallon water or 1 teaspoonful to quart. Lawns can be easily watered and fertilized at the same time by attaching a Hy-Gro Gat sprayer (see page 84) to your garden hose. Hy-Gro can be used on growing and fruiting plants as

of growing and fruting plants as a plant stimulant spray, at the rate of 4 lbs. to 100 gals. of water.

3 oz. **3**5c; 1 lb. \$1.00; **2** lbs. \$1.45; 5-lb. bag \$2.85; 10-lb. bag \$5.25; 100 lbs. \$25.00, f.o.b. Plant City.

KILGORE'S GRO-RITE. A liquid plant food containing nitrogen, phosphorous and potash, needed to produce healthy plant growth. Used as ground application, starter solution and foliage spray.

Compatible with most insecticides and fungicides. Excellent for airplane spraying. It is a liquid and won't clog nozzles.

Use 4 tablespoonfuls per gallon water or 1 gal. per 100 gallons.

10-10-10. 5 gal. \$6.75; 30 gal. \$39.00. 10-20-10. 5 gal. \$9.75; 30 gal. \$57.00.

CYPRESS GARDENS PLANT FOOD. A soluble plant food—analysis 8-7-20, plus ten trace elements. Excellent for hydroponic gardening and foliar feeding.

4 oz. 39c; 1 lb. \$1.00; 4 lbs. \$3.20; 10 lbs. \$6.95; 25 lbs. \$13.75.

HYPONEX—Valuable in seedbeds and to start plants after transplanting. Comes in a powder which makes liquid plant food containing all plant food elements plus vitamin B¹. Dilute 1 oz. in 6 gals. water.

2/3 oz. 15c; 2 oz. 29c; 5 oz. 59c; 10 oz. \$1.00; 10 lbs. \$9.00, f.o.b. Plant City.

NUGREEN. A soluble plant food containing 45% nitrogen in the form of urea. Gives excellent results when quick acting nitrogen is needed. Can be applied as a foliage spray in combination with insecticides and fungicides at the rate of 4 to 5 lbs. per 100 gals. Can also be applied through irrigation or as top dresser. 80-lb. bag \$5.85; ton \$143.00.

NUTRI-LEAF 60. A soluble fertilizer containing 20% nitrogen, 20% phosphoric acid and 20% potash plus minor elements. Compatible with most insecticides and fungicides as a foliage spray for plant stimulant. Increases growth and size of fruit and prolongs the life of the plant and picking period. Use 5 lbs. per 100 gals. water. 5-lb. bag \$1.45; 25-lb. bag \$6.90.



ORTHO-GRO. A liquid plant food for potted plants, flowers, lawns and gardens. 4-oz. bottle 39c; 16 oz. \$1.00; 32 oz. \$1.69; 1 gal. \$4.98.

PLANTABBS FOR HOUSE PLANTS. Plantabbs are a complete, balanced plant food of the high analysis 11-15-20. Use Plantabbs for all indoor potted plants and flowers, window and balcony boxes. Also for garden flowers and vegetables. Plantabbs with vitamin B¹ make vastly stronger root growth. 20 tablets 19c; 50 tablets 39c; 90 tablets 59c; 200, \$1.00; 1000, \$3.50.

VHPF "VIP"—A soluble fertilizer—containing the three major plant foods, nitrogen, phosphoric acid and potash plus essential minor elements. Used as a starter solution or a nutritional spray, it has proved very successful in combination with fungicide and insecticide sprays on growing and fruiting plants as a stimulant. Use 1 lb. to 20 gals. or 6 lbs. to 100 gals. water.

3-lb. bag 75c; 25-lb. bag \$6.00, f.o.b. Plant City or Kilgore stores.



NUTRIENTS



ACTIVO. A compost activator. Converts compost quickly and easily into available plant food.

No. 2 Size for 400 lbs. of Compost, \$1.39. No. 7 Size for 1700 lbs. Compost, \$2.95.

ADCO—A clean odorless powder for converting leaves, grass clippings, etc. into compost. 5 lbs. makes 200 lbs. organic compost. 5-lb. bag \$1.35; 20-lb. bag \$3.45.

ALUMINUM SULPHATE. For increasing acidity of soils where aluminum toxicity is no factor. 10c per lb.; 100 lbs. \$5.00.

CALCIUM CHLORIDE - FLAKE. Recommended for control of blossom-end rot on peppers and tomatoes if caused by calcium deficiency. Use 4 lbs. per 100 gal. water per acre. 4 to 5 applications may be necessary. Sold in 100 lb. bags only. 100-lb. bag \$5.00.

ES-MIN-EL. For soil application. Contains many essential mineral elements in proper proportions. Add to ordinary fertilizers at rate of 1 lb. or 2 lbs. to 10 lbs. regular fertilizer or 100 to 200 lbs. per ton. 5-lb. bag \$1.10; 100-lb. bag \$11.50, f.o.b. Plant City.

ES-MIN-EL. For spray or dust. For spray use 4 lbs. to 100 gals. water. For dust use 16 lbs. Es-Min-El mixed in 84 lbs. of talc, clay or other inert ingredients. Price: 4-lb. bag \$1.75; 50-lb. bag \$16.00, f.o.b. Plant City.

KILGORE IMPROVED C-M-Z Spray. A combined nutritional and fungicidal spray for tomatoes, potatoes, peppers and other vegetables. Do not use on sweet corn. Apply at the rate of 6 lbs. per 100 gals. of water. 6-lb. bag \$2.30.

KILGORE'S C-M-Z BEAN SPRAY—For use on beans for the correction of copper, manganese, and zinc deficiencies and for the control of rust and powdery

Use 10 pounds per 100 gallons of water at intervals of 10 days. Where rust is a serious problem apply every 5 days. The addition of spreaders is not necessary. Price: 10-lb bag \$1.95; 50-lb. bag \$7.45. KILGORE'S IRON-AID. (Code No. X-5). Contains Versen-ol, a newly developed chelated iron compound, which is completely available to plants under most soil conditions. — Designed for the correction of iron chlorosis wherever symptoms show up in flowers or shrubs. These symptoms are a paling of the areas between leaf veins, while the veins themselves still retain their original green color. We suggest the use of Iron-Aid even in questionable borderline cases. Consult label for specific directions before applying.



1 lb. 45c; 5 lbs. \$1.00; 10 lbs. \$1.90; 50 lbs. \$6.50 100 lbs. \$13.00 f.o.b. Plant City

LIME, HYDRATED. Use a half ton to a ton per acre, broadcast, before breaking the ground. If soil is new and sour, the application should be heavier. Also used for the neutralization of manganese, zinc and copper sulphates. 10-lb. bag 30c; 50-lb. bag 95c; ton \$34.00; ½ ton at ton rate, f.o.b. Plant City.

MANGANESE SULPHATE (75-78% spray grade). A quick acting material for applying manganese to plants in the form of a spray when plants are turning yellow on very alkaline soils. Apply at the rate of 2 lbs. to 50 gallons of water. 1 lb. 10c; 50-lb. bag \$2.85, f.o.b. Plant City.

MANGANESE-SULPHUR DUST NO. 1. (Code No. XS-1)—For manganese deficiency in beans, peppers, etc. 50 lbs. \$2.65; 100 lbs. \$5.30.

MAGNESIUM-SULPHATE (Epsom Salts)—For magnesium deficiency. 100 lbs. \$6.50; less than 100 lbs. 10c per lb.

VERSEN-OL—A chelated iron compound for soil application—soluble in water. Not recommended for foliar applications. 1-lb. can \$2.98; f.o.b. Plant City.

ZINC SULPHATE. For zinc deficiency in beans, tomatoes, etc. 100 lbs. \$10.95; less than 100 lbs. 13c per lb.

HORMONES

GIBBERELLIC ACID. This material is used primarily to break dormancy and stimulate emergence of potato seed pieces. For green seed use 1 ppm solution as an instant dip. If eyes are colored, use 1/2 ppm. When eyes have sprouted, use 1/4 ppm.

Concentrate	pt.	qt.	gal.
1,000 ppm	\$ 4.50	\$ 8.25	\$ 32.00
4,000 ppm	\$15.00	\$27.75	\$110.00
10,000 ppm	\$35.00	\$68.00	\$270.00

HORMODIN POWDERS. Hormodin is a hormone-like plant growth substance. Its principal use is to stimulate rapid growth on cuttings. It has wide-spread use in propagating cuttings. Many cuttings formerly impossible to propagate may now be rooted with the aid of Hormodin. Easy to use. Simply dip moistened stems in the powder and then plant.

No. 1—For Chrysanthemums, Poinsettias, Roses, etc. $\frac{1}{4}$ oz. 25c; $\frac{1}{4}$ oz. 65c; 1 lb. 4.00.

No. 2—For semi-woody plants and most shrubs. 134 oz. 95c; 1 lb. \$5.95.

No. 3-For resistant species, such as Evergreens. 1 oz. \$1.30; ½ lb. \$5.95.

Combination Package, powder Nos. 1, 2 and 3, in carton, 95c.

FRUITONE is a wettable powder containing several hormones plus an effective wetting agent for securing a thorough coverage. Fruitone when sprayed on plant blossoms and young set fruit will help to prevent premature dropping. Cut flowers and twigs dipped in Fruitone will hold up longer.

Use 1 level teaspoonful to 1 gal.; 2/5-oz. to 5 gals.; 2 oz. to 25 gals.; 8 oz. to 100 gals. of water.

Price: 2/5-oz. pkg. 35c; 2-oz. can \$1.25; 12-oz. can \$5.00.

ROOTONE. A hormone powder to hasten the rooting of cuttings. It is easy, economical, safe and sure treatment for rooting cuttings. Low cost—1/30 to 1/50 cent per cutting. ¼ oz. 35c; 2 oz. \$1.25; 1 lb. \$5.00; prepaid.

ROOTONE NO. 10. Recommended especially for rooting azaleas and camellias. 2 oz. jar \$2.25; 1 lb. jar \$10.00.

TRANSPLANTONE supplies the plant with chemicals which stimulate root formation and growth after transplanting ½ oz. makes five gallons solution. ½ oz. 35c; 3-oz. can \$1.25; 1-lb. can \$4.00.



HOUSEHOLD AND GARDEN PESTICIDES

READY-TO-USE

DUST

el GORE

GENERAL

PURPOSE

DUST

COMPANY PLANT CHE

KILGORE'S C-M-Z GARDEN SPRAY. A combined nutritional and fungicidal spray for shrubs, fruit trees and vegetables. Use two tablespoons per 1 gallon of water on Avocados, Mangos and shrubs, 134 tablespoons on citrus and 2½ tablespoons on vegetables. No wetting agent required. 1-lb. can 75c.

KILGORE'S 25% DDT EMULSION. For the control of insects on truck crops, shrubs, etc. Qt. \$1.35; gal. \$3.15.

KILGORE'S FLEA-KILL. (Code No. GV-1.) Apply dust under houses and other buildings, around kennels and on bare sandy areas at the rate of 2 lbs. per 100 sq. ft. Dust lawns at the rate of 2 lbs. per 250 sq. ft. Be sure grass is dry before applying dust. 5 lbs. \$1.65; 10 lbs. \$3.00; 25 lbs. \$6.50; 50 lbs. \$12.00; 100 lbs. \$24.00.

KILGORE'S GENERAL PURPOSE DUST FOR FLOWERS AND SHRUBS. Kilgore's General Purpose Dust for flowers and shrubs is formulated especially to control all types of insects and diseases on flowers and shrubs in Florida. It controls aphids, thrips, scale insects, mites, caterpillars as well as leaf spots, blight, powdery mildew and many other fungus diseases attacking ornamentals. Dust gun complete with dust \$1.00; 1-lb. bag \$0c; 5-lb. bag \$2.00; 10-lb. bag \$3.75.

KILGORE'S GENERAL PURPOSE DUST FOR VEGETABLES. Kilgore's General Purpose Dust for vegetables is formulated especially to meet the needs of the small home gardener. Safe for use on all types of vegetables grown in Florida. A truly General Purpose Dust that controls both chewing and sucking

insects, as well as fruit rots, downy mildew, anthracnose, early and late blight, cercospora leaf spot and many other fungus diseases. It has been formulated particularly to reduce any residue hazard from highly poisonous materials. Safe to apply and easy to use. Dust gun complete with dust \$1.00; 1-lb bag 60c; 5-lb. bag \$2.50; 10-lb. bag \$4.50.

ISOTOX GARDEN SPRAY—An effective spray containing 5% lindane, for control of insects on roses, camellias, azaleas, ixoras, hibiscus, bougainvillea, and other ornamentals, house and garden plants. Use four teaspoons per gallon of water. Price: 4-oz. bottle \$1.00; 8-oz. \$1.85; 16-oz. \$2.95.

KILCHINCH DUST. (Code No. VHDG-1.) For chinch bug control in St. Augustine type grasses. 5-lb. bag \$1.25; 10-lb. bag \$2.40; 50-lb. bag \$10.10; 100-lb. bag \$20.20.

KILGORE'S KILCHLOR. A residual household spray. Contains 2% chlordane plus isobornyl thiocyanoacetate, for control of roaches, ants, and most other household insects. Qt. 65c; gal. \$2.15.

KILCHLOR-KILPHENE POISON BAIT NO. 11. (Code No. B-11)—(Contains chlordane and toxaphene). For control of mole crickets, cutworms, and many soil insects. 1 lb. 20c; 25 lbs. \$2.70; 50 lbs. \$4.40; 100 lbs. \$8.80. KILCOP "45"—(Contains 45% metallic copper.) Kilcop is extremely fine which is essential for good coverage and effective control of fungi. Consult label for dosages. 1-lb. can 90c.

KILGORE'S KILSPRAY. An excellent indoor space spray for flies, mosquitoes, roaches, ants, fleas and other household insects. Not a residual spray. It should be used when insects are present. Qt. 65c; gal. \$2.15.

KILSPRAY—Air Freshener, Bombs—Stops all offensive odors. Quick and easy to use. Non-toxic, non-inflammable. Comes in two odors—Bouquet and Pine Needle. Price: 10-oz. Bomb, 97c each.

KILSPRAY INSECT BOMBS—Kills flies, mosquitoes, gnats, flying moths, roaches, ants and silverfish. Kills quickly. Easy to use. Price: 12-oz. Bomb 97c, prepaid.

MALATHION 50% Emulsion. For control of scale and many other lawn and garden insects. Use 2 tablespoons per gal. of water. 4 oz. \$1.19; 8 oz. \$1.89; pt. \$2.89; qt. \$4.39.

KILGORE'S NEMA-KIL. (Contains Nemagon). A new

nematocide, which kills nematodes, including rootknot, sting nematodes, burrowing nematodes, stubby-root nematodes, etc. Can be used safely on many established plants, such as hibiscus, roses, gardenias, azaleas, ixoras, as well as lawn grasses and most flowers without harm to plants or grasses. NEMA-KIL is emulsified in water and easy to apply. It is the answer to a long-felt problem as to how to get rid of nematodes in your yard without digging up lawns, shrubbery, etc. One quart of NEMA-KIL will treat from 600 to 800 square feet of surface. Price: pt. \$1.35; qt. \$2.25; 1 gal. \$6.75.



NEMA-KIL 17.5 G GRANULES. 1 lb. 98c; 5 lbs. \$3.95.

NO NIB'L RABBIT REPELLENT.—Recommended as a rabbit repellent for flowers, shrubs and other ornamentals. Non-injurious to birds and other animals. It is easy to use and is effective either as a dust or as a spray. Price: 6 oz. shaker-top can \$1.29.

ORTHO-KLOR "44" SPRAY—For control of ants, mole crickets, cutworms, army worms, roaches and many other pests. For Termite Control see page 58. 4 oz. 79c; 8 oz. \$1.20; 16 oz. \$1.98; 32 oz. \$2.98.

ORTHOL GARDEN SPRAY. An oil emulsion containing nicotine and DDT for the combined control of aphids, red spiders, mealy bugs, thrips and many other insects. Use at the rate of 6 tablespoonfuls to 1 gal. of water. 4 oz. 35c; 16 oz. 75c; 32 oz. \$1.25; 1 gal. \$2.85.

RED ARROW INSECT SPRAY. Kills aphids, worms, leaf-hoppers, white flies, red spiders, caterpillars, thrips, celery leaf tier. Average dilution 1 to 400 parts water. 4 oz. \$1.20; pint \$3.49; gal. \$25.00.

KILGORE'S ROSE DUST combines the latest materials for the control of all types of rose diseases, including powdery mildew, black spot and many other fungus diseases, as well as controlling aphids, mites, thrips, and many chewing insects. Contains three different fungicides to give the best disease control possible. Also contains two insecticides and one miticide to give control of the many insects attacking roses in Florida. Can be used as a general purpose dust for almost any type of flower or shrub grown in Florida. Dust gun complete with dust \$1.00; 1-lb. bag 55c; 5-lb. bag \$2.25; 10-lb. bag \$4.00.



SABA CHINCH. (Code No. S-2.) For control of chinch bugs, army worms, sod webworms, and ants. Use 4 lbs. per 100 sq. ft. Lb. 20c; 5 lbs. 75c; 10 lbs. \$1.40; 25 lbs. \$2.90.

KILGORE'S SLUG DUST—A 15% Methaldehyde dust providing effective control of slugs in nurseries, greenhouses, etc. Use at the rate of 2 lbs. per 1000 square feet. Usually three applications are required. 1-lb. Canister \$1.00; 6-lb. bag \$3.30; 25-lb. drum \$15.00; 50 lb. drum \$25.00.

VOLCK ISOTOX SPRAY. For use on ornamental plants against red spiders, certain scale insects, mealy bugs, white flies and certain other garden pests. Use 5 tablespoonfuls per 1 gal. water. Don't use more than three applications per season. 4 oz. 49c; 8 oz. 69c; 16 oz. \$1.19; 32 oz. \$1.98.

VOLCK OIL SPRAY. 4 oz. 39c; 16 oz. 79c; 32 oz. \$1.49; 1 gal. \$3.19.



SEED INOCULANTS POLYETHYLENE FILM

All prices quoted are postpaid

When planting a legume feed or cover crop, such as clovers, cowpeas, beggarweed, crotalaria, soy beans and velvet beans, first inoculate the seed. This helps the crop to fix the maximum amount of valuable nitrogen plant food material from the air in your soil. A



good leguminous cover crop can take 50 to 150 pounds of nitrogen per acre from the air—if inoculated. For your soil's sake, inoculate all legume seed.

Group A—Alfalfa and Sweet Clover	} 100 lb. size 70c
Group B—Clovers White Special	100 lb. size 70c
Group C—Vetch, Garden and Sweet Peas	100 lb. size 55c
Group D—Beans—all varieties	1 lb. size 60c
Group E—Peanuts, Cow- peas, Velvet Beans, Crotalaria, Beggar- weed and Alyce Clover	}100 lb. size 55c
Group L—Lespedeza	$\}$ 100 lb. size 60c
Lupines—all varieties	} 100 lb. size 55c
Special Cultures: Lima Bean Trefoil, Dalea or Woods Clover, Crown Vetch, Chick Peas, Hairy Indigo and Sesbania	1/8 bu. size 25c 1 bu. size 70c
Garden Size, for Garden)

GER-PAK®

A durable, versatile plastic film for thousands of uses around the home and farm. Prices quoted are f.o.b. Plant City or any Kilgore Store.

11/2 Mil Thickness	To 25 M	25 M To	50 M To
	SQ. FT.	50 M	100 M
3' x 500' Black Only	8.85	8.25	7.65
4 Mil Thickness			
6' x 100' Clear Only	9.00	8.40	7.80
8' x 100' Clear Only	12.00	11.20	10.40
10' x 100' Black or Clear	15.00	14.00	13.00
12' x 100' Black or Clear	18.00	16.80	15.60
14' x 100' Black Only	21.00	19.60	18.20
16' x 100' Black or Clear	24.00	22.40	20.80
20' x 100' Black or Clear	30.00	28.00	26.00
40' x 100' Black or Clear	60.00	56.00	52.00
6 Mil Thickness			
6' x 100' Clear Only	13.50	12.60	11.70
10' x 100' Clear or Black	22.50	21.00	19.50
12' x 100' Clear Only	27.00	25.20	23.40
16' x 100' Clear or Black	36.00	33.60	31.20
20' x 100' Clear or Black	45.00	42.00	39.00
40' x 100' Clear or Black	90.00	84.00	78.00
2" Black Waterproof Ta	pe (180 F	t. Rolls)	4.95 Ea.

POULTRY SUPPLIES

All prices f.o.b. Plant City or Kilgore Stores and subject to change without notice.

A more complete line of poultry supplies is carried in all Kilgore stores.

HUDSON HANDY FILL CHICK FOUNTAIN



No. A-4—Capacity one gallon	\$1.65
No. A-8—Capacity two gallons	2.05
No. C-3—Capacity three gallons	3.30
No. K-5—Capacity five gallons	6.30

MASON JAR FOUNTAIN AND FEEDER

No.	O-25	Plastic	Pan	to	fit	Ma	son
ja	r					\$.20

DIXIE INOCULANT

Lupines.

Peas, Sweet Peas, All Beans, Soy Beans and

Made In Florida Especially For Florida Soils

Each package will inocu-

late 12 lbs. seed—25c

		1
	(size)	Price
Soy Beans	2 bushels5 bushels	.\$.30 60
Lespedeza, Alyce Clover Hairy Indigo	100 pounds	.50
Cowpeas, Crotalaria, Lima Beans, Peanuts and Velvet Beans	}100 pounds	.50
All Vetches, Austrian Peas, Wild Winter Peas, Garden and Sweet Peas	}100 pounds	50
Lupines, all varieties	}100 pounds	50
Clovers, Red Crimson, Ladino, White and Louisiana White	}100 pounds	.70
Alfalfa, Hubam, Floranna, Sweet and Bur Clovers	}100 pounds	70
Garden Size	} 8 pounds	20

REEL TOP FEEDERS

No.	118	Metal	Feeder,	18-inch	length	\$.55
No.	224	Metal	Feeder,	24-inch	length	.85
No.	336	Metal	Feeder,	36-inch	length	 1.15

HUDSON "KLEEN-EZY" REEL TOP FEEDERS No. 6360 A-36-inch length-Reel Top\$2.05

KEY RING LEG BANDS. Colors: red, yellow, green, white and blue. Sizes for baby chick, pigeon, bantam, medium, intermediate and large. Specify size and color. 1c each; 75c per 100.

		Ea		Doz.	
NEST	EGGS—china	\$.05	\$.50

THERMOMETERS

Black oxidized brass scales, white filled figures and

graduations, mercury tubes.	
Weather thermometer, No. 5152, 8-in. wood	.80
Weather thermometer, No. 5140, 71/8-in, wood case 1.	.30
Weather thermometer, No. 5132, 7-in. Metal Utility 1.	.90

KNAPSACK AND HAND DUSTERS

THE CHAMPION NO. 100 DOUBLE ACTION DUSTER is carried like a knapsack. It fits comfortably on the back and leaves both hands free for efficient operation. The easily operated bellows throws a screen of dust exactly where wanted within a range of 30 feet. The complete duster weighs only 14 pounds empty and will hold as much as 17 pounds of fluffy powder. Can be converted to single ac-

tion by removing upper intake valve. Price\$29.35,Prepaid.



Champion No. 100 Double Action Duster

SMALL HAND DUSTERS

ROOT HAND DUSTER 33C

(Less "Y" Attachment)

Price \$35.80, Postpaid

ROOT HAND DUSTER


MODEL 33C

with 98Y Attachment

\$38.80, Postpaid

f.o.b. Plant City







HUDSON ADMIRAL NO. 766-C

CHAMPION HAND DUSTER - MODEL 103



Champion Hand Duster Model 103

Simplicity of design and ease of operation make the Champion Hand Duster ideal equipment for flowers, vegetables, shrubs and small trees. It is used extensively in home gardens and small estates as well as in experimental and show gardens; very efficient and economical for spot dusting s m a l l plants in the field such as watermelons, etc. It is of highest quality construction. All parts

are enclosed and the steel tank is lead coated for long life. It is extremely light in weight and will carry up to eight pounds of powder, although it operates effectively with only a fraction of an ounce of powder in the tank. Price \$9.85, Prepaid.

LITTLE CHAMP NO. 902. Rotary Hand Duster. Made especially for home garden. Sturdy built, 8" tall. Wt. 234 lbs. Capacity 2 lbs. Easy to turn. Puts out a strong blast of dust. \$9.85, Postpaid.

HUDSON HAND DUSTERS

Made of aluminum sheet and castings for lightest weight. The Hudson Roto-Power "806" is designed for growers who want the easiest turning and lightest

ca pr ar sa bl pl ag fr pe

carrying duster, one that produces the high velocity and volume of air necessary to break up dust and blast it right into the plant. Holds 14 lbs. average density dust. Applies from 5 to 45 lbs. of dust per acre. High velocity: 1 crank turn produces

60½ fan revolutions. Price with "Y" attachment \$46.35; without "Y" attachment \$42.50. Postpaid.

Hudson No. 806C Roto-Power Duster (Aluminum) f.o.b. Plant City or Kilgore stores

RESPIRATORS

Postpaid

Approved for parathion dustingeach	\$6.25
Replacement cartridges, No. 11-A (4 to box)Box	2.65
Replacement filters, No. R-490 (10 to box)Box	1.60

GOGGLES

Postpaid

Monogoggles	No.	93A	Clear	Lens		2.75
Monogoggles	No.	93A	Green	Lens	***************************************	2.90

FLAME SPRAYERS


Postpaid

POWER AND HAND SPRAYERS

Prices quoted are f.o.b. Plant City unless otherwise indicated

HUDSON POWER SPRAYER

High pressure sprayer recommended for estates, small groves, dairy barn and cattle spraying, and landscape gardeners. Handles all insecticides, fungicides, weed control chemicals and whitewash.



No. 31030 Matador Cushion Tires, 30-gal. tank...\$318.55 No. 31015 Matador Cushion Tires, 15-gal. tank... 278.00

COMPRESSION AIR SPRAYERS

Prepaid

Compressed air sprayers are strong, efficient and easy to use. Ideal for dozens of spraying jobs in homes, truck gardens, dairy barns, poultry houses, etc.

Brown Compressed Air, Galv. Open-Head No. 64-D (4-gals.)	\$21.20
Hudson Compressed Air	
Perfection Sprayer (4-gals.) No. 310	19.95
Hudson Bugwiser No. 220 (2½-gals.)	10.50
Hudson Leader No. 10 (3-gals.)	12.95
Hudson Climax No. 335 (3-gals.)	17.95
Dobbins 44-A (4-gals.)	20.70
Dobbins No. 21-A (3½-gal. capacity)	14.25
Dobbins No. 24-A Budget (3½-gal. capacity)	11.55
SPRAYER HOSE	



Leader No. 293-B

%" Hose. Price per ft.25

CHAMPION SPRAYER MODEL-1



This all-purpose portable sprayer is tops in its field. The Champion combines in one unit all the desirable qualities of unusual power, simple operation, effective control, continuous agitation, and long life. Uniform pressure of 200 lbs. in the sturdy pressure cylinder is pumped as needed

by the operating lever. This lever also operates the brass agitator which keeps 4½ gallons of any liquid well mixed. Price \$36.00, Postpaid.



HOUSEHOLD SPRAYERS

No. 208, Hudson Nebu-lizer, Super Cloud 8 oz. capacity	\$.65
No. 333, Hudson Nebu-lizer, Super Fog 10 oz. capacity	
No. 444, Hudson Nebu-lizer, Super Misty 27 oz. capacity	

HOUSEHOLD CONTINUOUS SPRAYERS

TROMBONE TYPE SPRAYERS

Ken Sprayall No. 202	89.95
Redisprayer No. 214 w/extension	7.95
	4.95

GAT SPRAYER. Attaches to garden hose for dissolving, mixing and spraying Hy-Gro and other soluble plant foods.

Empty 2-lb. Hy-Gro jar with Gat Sprayer\$2.50



BRADSON SPRAY GUNS



Pest-Gun

For use with standard garden hose. Pest-Gun, Gard-N Gun and Bug-Gun for spraying insecticides and fungicides. Gro-Gun for application of soluble fertilizers and plant foods



Bug-Gun

Pest-Gun No. 106	34.95
Gard-N-Gun No. 102	2.69
Gro-Gun No. 103	2.95
Bug-Gun No. 104	1.98

LAWN SPRINKLERS

Rain King Model-K Adjustable	\$9.95
Rain King Model-K-GSS, Rust resistant	10.25
	5.95
Rain King Model N1 Garden hose nozzle	1.50
Park Sprinkler Model 911	2.95







Model D

FARM AND GARDEN TOOLS

All prices quoted are f.o.b. Plant City or Kilgore stores



No. 4-D Seeder—Shipping weight 41 lbs. Price....\$35.70 No. 4 Seeder—Shipping weight 50 lbs. Price.......... 44.35 No. 300-A Seeder—Shipping weight 55 lbs. Price 45.50

PLANET JR. NO. 219 FERTILIZER DISTRIBUTOR



No. 8558X, Covers for No. 215. Shipping weight 2 lbs. Price.... 1.95

PLANET JR. NO. 220



No. 220 Fertilizer distributor\$72.90 Shipping Weight 115 lbs.

PLANET JR. NO. 13 DOUBLE AND SINGLE WHEEL HOE



No. 13 Double and Single Wheel Hoe—Shipping weight 27 lbs. Price\$21.75

NO. 17 SINGLE WHEEL HOE




NO. 19 WHEEL HOE



NO. 119 HIGH WHEEL PLOW Shipping wt. 18 lbs.\$12.50


GARDEN TRACTORS



CYCLONE SEEDER

The Cyclone will sow many acres per hour, adjustable to any amount per acre.

In ordering seeder to be used with power take-off, specify make and model of tractor to be used. Electric motor model operates from 6 volt battery from jeep or truck.



INSCO SOIL FUMIGANT APPLICATOR

A 4-gal. tank gravity flow applicator for single row application. Can be attached to horse drawn plow or small tractors. Price\$15.00

FARM AND GARDEN TOOLS (Continued)

Prepaid unless specified. Prices subject to change without notice.

ECLIPSE POWER MOWER



ECLIPSE builds precision power mowers with extra features for unmatched performance in low-cost maintenance of beautiful lawns. Balanced design, durable construction, dependable power, self-propelled, precise cutting action and effortless handling make lawn mowing easier, faster and better.

REEL TYPE

Lark 18-in. Recoil Starter	124.50
Rocket 20-in. Recoil Starter	
Parkhound 21-in. Recoil Starter	172.50
Sharpeners for reel type mowers available	

BOTARY TYPE

NOIANI IIFL	
Duo-Master (Hand propelled) 17-in\$	79.95
Duo-Master (Hand propelled) 19-in.	
Duo-Master (Hand propelled) 20-in.	99.95
Leaf Mulcher for Rotary Models	3.00
Recoil Starters for any Power Mower	5.00
Electric Starters for any Power Mower	35.00
Prices f.o.b. Plant City or Kilgore Stores	

ECLIPSE HAND MOWERS

Eclipse	"L"—16-inch	 36.00
Eclipse	"L"—18-inch	 37.50

Prices f.o.b. Plant City or Kilgore stores

TRUMP GARDEN **TOOLS**

Sturdy, attractive hand made tools finished in baked green enamel. Set includes trowel, fork and cultivator.



No.	800	Trowel250	each
No.	801	Fork	each
No.	802	Cultivator250	each
No.	804	Transplanter25c	each
No.	805	Weed Cutter25c	each
No.	811	3-piece Set85c p	er set

WISS GARDEN TOOLS

Postpaid

FH-4 Flower Cutter Holder, Chrome Plated 6½....\$2.75



No. 808 Draw Cut Pruning Shears

No. 808 Draw Cut	\$3.95
No. 607-R Pruning Shears	2.95
No. C-1 Orange Clippers	2.75

WISS GARDEN TOOLS (Continued)



No. 81/2-E Hedge Shears

٧o.	8½-E Hedge Shears\$	5.75
Vo.	9-A Hedge Shears	5.00
٠oľ	9½-E Hedge Shears	6.25



No. 224-T Lopping Shears

No.	321-T	21	inch		6.50
-----	-------	----	------	--	------



No. 5600 Forged Grass Shears


No.	5600	Forged Grass Shears	\$2.95
		Grass Shears	
No.	900	Grass Shears—Long Handle	4.95
No.	801	"Quick Trim"	. 1.98
		Grass Shears	

LAWN FERTILIZER DISTRIBUTORS

f.o.b. Plant City or Kilgore stores



Fox Spreader J-20 inch-50-lb. capacity all steel construction with tubular steel handle, rubber handle grip. Steel wheels with rubber tires. For the customers who want "something better" in a small spreader. Price \$13.95.



JACKSON Handi-Cart

—The Handi-Cart has an outstanding reputation among users as a sturdy, dependable tool for hundreds of moving jobs around the home and garden.

3 cu. ft. Handi-Cart, wt. 25 lbs	.\$10.20
4 cu. ft. Deluxe Cart, wt. 33 lbs.	12.50



FARM AND GARDEN SUPPLIES

Prices subject to change without notice.

BIRD NEPONSET PAPER POTS



Nutrient Treated

Made of tough waterproof paper, adapted for growing plants from seed to transplanting age. Plants get no setback when transplanted. Performs the major function of clay pots at a fraction of the cost.

Red Paper Pots

Neponset Red Paper pots are designed for shipping plants which have been grown in clay pots or in other containers. Shipping costs are lower by using Neponset Red Pots because there is con-

siderable less weight and less breakage, and the clay pots can be saved for further use.

By mail post 100 pots			f.o.b. Pla 1000	ant City pots	
Vita	Red	Vita	Wt.	Red	Wt.
2"\$2.65	\$1.65	\$16.50	2 0 lbs.	\$ 9.90	10 lbs.
2½" 2.90	1.75	17.05	31 lbs.	10.00	16 lbs.
3" 3.25	1.80	19.15	43 lbs.	10.60	22 lbs.
3½" 4.25	2.35	25.00	60 lbs.	14.10	32 lbs.
4" 4.80	2.90	28.30	38 lbs.	17.05	21 lbs.
5" 6.30	3.65	36.75	60 lbs.	20.90	33 lbs.
6" 8.20	4.35	48.50	48 lbs.	25.75	47 lbs.

PLANT BANDS		f.o.b.
Vita-Band "10"	Postpaid	Plant City
Contains sufficient fertilizer for 10 weeks	per 100	per 1000
134 x 134 x 2½ inches	.\$ 1.05	\$5.50
2 x 2 x 2½ inches		5.80
2 x 2 x 3 inches	. 1.26	6.35
2½ x 2½ x 3 inches		6.90
3 x 3 x 3 inches		7.60
4 x 4 x 3 inches	. 1.98	9.90

Vita-Band "D"

A nutrient treated band which disintegrates in the soil.

1½ x 1½ x 2½	1.05 1.20 1.45	f.o.b. Plant City per 1000 \$5.35 5.70 5.90 6.95 10.20
TOM THUMB FLATS No. 1 (for bands 1¾ x 1¾ x No. 2 (for bands 2 x 2 x 2½)	f.o.b. per 100 2½)\$5.50 6.10	f.o.b. per 1000 \$49.20 54.60

"JACK POT" PEAT MOSS POTS

These are sturdy wall pots made from pressed peat moss. Plants started in these pots never receive a transplant set-back. Plant pot and all. Peat holds moisture and fertilizer within reach of roots.

		f.o.b.	Plant City
	per 100	per 1000	wt.
134" Square	\$1.50	\$ 7.00	30 lbs.
21/4" Round		7.25	32 lbs.
21/4" Square	2.20	11.00	32 lbs.
3" Square		18.25	35 lbs.
3" Round		13.25	30 lbs.
4" Standard		29.25	30 lbs.
4" Azalea		28.75	28 lbs.

Standard packaging: 1¾4"—2,500; 2¼"—2,000; 3"—1,000; 4"—500 per case.

Write for special prices on larger quantities.

TERRA-LITE (VERMICULITE)



An excellent light weight medium for starting cuttings, seed in flats, paper pots or plant bands. Can be mixed with soil in seedbeds. Increases germination by reducing damping-off. Holds moisture for several days without rewatering. 4-qt. bag 39c; ½-bu. bag 89c; 4-cu. ft. bag \$2.35.

f.o.b. Plant City

Write for complete descriptive pamphlet on Terre-Lite.

PEAT MOSS

Good to add humus and holds moisture in seedbeds. Also an excellent poultry litter.

Georgia Peat Humus, 1½ cu. ft. bag \$1.40 each 5 cu. ft. bag \$3.50 each.

European Peat. Compressed in burlap.

Wrapped bales 20" x 20" x 36" each \$4.95

Bulk 10c per lb.

f.o.b. Plant City

SPHAGNUM MOSS

Burlap Wrapped Bales 14" x 16" x 45" each \$4.00 Bulk 50c per lb. f.o.b. Plant City

OSMUNDIA FIBRE —Used for growing Orchids.	
Per Bale\$	9.50

RUBBER STRIPS-For Budding and Grafting



Postpa Size	i d Length	Width	Approx. Pieces Per Lb.		Lb.	5 Lbs.
No. 520	5 in.	3/16 in.		.25	\$2.75	\$12.00
No. 620	6 in.	3/16 in.		.25	2.75	12.00
No. 838	8 in.	3/8 in.		.25	2.75	12.00

AIRWRAP

Plastic adapted to grove and plant propagation. Coated with nutrient and rooting hormones.

9" width—8 vd rolls \$1.95

9"	width—8 yd. rolls\$	1.95
	x 8" price per sheet	.06
9"	width—less than 8 yds. per yd	.25

Write for quotations on larger quantities

AIRLAYERING ROOTING KIT

For plant propagation. Kit contains sphagnum moss, plastic wraps and tying strips. Complete instructions in box.

Kit complete (Prepaid)\$1.95



GLOVES



Green Thumb



FARM AND GARDEN SUPPLIES (Continued)

Prices subject to change without notice

TWIST-EM PLANT TIES

Postpaid

Helps you support plants quickly, neatly, perman	nently.
Package of 35—8-inch size	15c
Package of 50—8-inch size	20c
Box of 100—8-inch size	29c
Box of 100—16-inch size	57c

Utility Twist-Ems Flat

1/4"	\mathbf{x}	4"	\$1.	15	prepaid
1/4"	\mathbf{x}	7"	1.	75	prepaid
1/4"	\mathbf{x}	8"	1.	95	prepaid
1/4"	\mathbf{x}	12"	2.	95	prepaid
1/4"	x	16"		90	prepaid
	1/4" 1/4" 1/4"	1/4" x 1/4" x 1/4" x	1/4" x 7" 1/4" x 8" 1/4" x 12"	14" x 7" 1. 14" x 8" 1. 14" x 12" 2.	1/4" x 4" \$1.15 1/4" x 7" 1.75 1/4" x 8" 1.95 1/4" x 12" 2.95 1/4" x 16" 3.90

TWIST-N-TYE ROUND

The grower's all-purpose vine and plant tie.

	Size	,	Price	
5	inch		per	\mathbf{M}
7	inch		per	\mathbf{M}
8	inch	\$1.00	per	\mathbf{M}
9	inch	\$1.15	per	\mathbf{M}
16	inch	\$1.95	per	M

Prices quoted are f.o.b. Plant City and all Kilgore stores.

TRAIN-ETTS

New weatherized trellis netting, ideal support for sweet peas and any vining flower or vegetable plants. Comes in three sizes. No. 1, 60×72 in., 75c; No. 2, 60×96 in., 95c; No. 3, 60×180 in., \$1.35, postpaid.

BAMBOO CANE STAKES

Dyed green pencil-thickness	
3 ft. Stakes, per 1000 f.o.b. Plant City	1 2 .50
3 ft. Stakes, per 100 postpaid	1.50
Less than 100 2½c each, postpaid.	
4 ft. Stakes, per 1000 f.o.b. Plant City	16.00
4 ft. Stakes, per 100, postpaid	2.00
Less than 100 3c each, postpaid	

LABELS—Plant and garden (painted)

Postpaid	Per 100	Per M
4" Pot Labels	\$.50	\$3.75
5" Pot Labels	55	4.10
6" Pot Labels	60	4.50
8" Garden Labels	1.05	9.00
10" Garden Labels	1.15	10.20
12" Garden Labels	1.90	17.50
3½" Copper Wired	60	4.50

GRAFTING WAX-Postpaid

A prepared wax recommended for all hand and brush grafting operations. 1/4 lb. 45c; 1/2 lb. 60c; 1 lb. 90c.

BUDDING and GRAFTING TAPE—Postpaid

Grafting Ta	ape ¼-in.	wide,	60-yd.	Roll		\$ 1.10
Grafting Ta	ape ½-in.	wide,	60-yd.	Roll		 1.45
Grafting Ta	ape 1-in.	wide,	60-yd.	Roll		 2.15
Plastic Bud	ding Tap	e 5⁄8-in	ı. wide	100-	yd. Roll	 .80

BUDDING KNIVES-Postpaid

 	
No. 11 Budding Knife, brass lining, 4½-in. Price \$	3.50
No. 21SH Florist Knife, 2 blades, Price	3.25
No. 8½C Grafting Blade, Price	1.75
No. 9C Budding Blade, Price	1.75

HOTKAPS

Hotkaps are "individual hothouses" for every plant—protecting them against frost, rain, wind, insects and crusting. They force plant growth, making hardier, healthier plants and producing crops two or three weeks earlier. They may be used with practically any crop that needs protection, but are especially suited for use with vine crops such as tomatoes, and where constants and water.



cucumbers, cantaloupes and watermelons.

20 pkg. Complete with Paper Setter, each\$	
100 with Fibreboard Garden Setter	4.60
250 with Fibreboard Garden Setter	7.50
1000 without Setter	22.65
5000 without Setter, per 1000	22.60
10000 without Setter, per 1000	22.55
25000 without Setter, per 1000	
Fibreboard Garden Setter, each	
Steel Setter, each	
•	

Prices quoted are f.o.b. Plant City and all Kilgore stores

LABEL PASTE

Ours is a genuine wheat paste, especially made for pasting labels on vegetable and fruit boxes. Mixes easily in cold or warm water. Will not discolor labels. Unaffected by water or atmospheric changes after drying. 28c per lb.; 100 lbs. \$25.35.

Prices quoted are f.o.b. Plant City and all Kilgore stores

TWINE

Jute—Soft vegetable 2 ply (app. 1520 ft. to lb.) tying twine, excellent quality. Used for tying tomato plants and bunching vegetables. ½-lb. ball 35c; lb. (cut) 52c; reel (50-60 lbs.) 49c per lb.

Prices goted are f.o.b. Plant City and all Kilgore stores

Cotton Twine (Variegated) — Used for tying pole beans. 2½-lb. cones. Price per lb. \$1.20.

f.o.b. Plant City

RAFFIA

The plain or natural color comes in uneven weight hanks, 1 lb. \$1.40; 3 to 7 lbs. \$1.00 per lb.; 10 lbs. \$9.00, Prepaid.

TARPAULINS

Write for prices stating size desired

HAND TALLY REGISTERS-Pocket Size

A fast and accurate way to count live stock, poultry, watermelons, plants, packed crates, etc. No. O counts to 999, \$3.80. No. 1 counts to 9999, \$4.90. Postpaid.

SUDBURY SOIL TEST KIT

Home Garden Model

Complete equipment with instructions and solutions for 20 tests, for Nitrogen, Phosphorus, Potash and Acidity. \$3.98 Postpaid.



Jr. Professional Model

Contains complete equipment and instructions—solutions for 50 tests for nitrogen, phosphorus, potash and acidity (pH). Transparent plastic case. Price \$6.98, Postpaid.

Two Minute Acidity Kit\$1.98

Write for prices on larger models.

CANNING SUPPLIES

Prices subject to change without notice

BURPEE PRESSURE COOKER



Model AR25—Holds 10 qt. or 17 pt. cans or 7 qt. or 14 pt. jars. Equipped with canning basket, recipe book, and cooking pans. \$29.95, Postpaid.

Model AR25

Model AR1525 holds 15 qt. or 25 pt. cans, or 14 qt. or 21 pt. jars. Equipped with 2 canning baskets, recipe book, and pans. \$34.95, Postpaid.

BURPEE SIMPLEX SEALER WITH CAN SAVER ATTACHMENT



Seals cans quickly and easily. Set the can in position, spin the crank with the right hand and move a lever with the left hand. Made of hardened steel for long life. Rustproof. Equipped with attachment which simplifies changing from one size can to another. Clamps on a table. Seals, opens and reflanges two sizes of cans, the No. 2 and No. 3. Tested canning recipes and illustrated instructions with every sealer. Net weight 11 lbs. Price \$19.95, prepaid.

SANITARY CANS

f.o.b. Plant City

Our sanitary open top cans are ideal for home canning. The finest fruits or vegetables and all your hard work can be wasted if the cans are not right.

Our cans for home canning are uniform, accurate, dependable, and strong. For years they have been recognized as cans of highest quality, preferred by home canners.

	No. in		
	Case	Case	1,000
No. 2 Plain Sanitary cans, wt. 26 lbs.	100	\$5.30	\$52.65
No. 2 Plain Sanitary cans, wt. 6 lbs.		1.40	56.45
No. 3 Plain Sanitary cans, wt. 36 lbs.	100	8.10	79.85
No. 3 Plain Sanitary cans, wt. 9 lbs.	24	2.10	84.95
No. 2 Sanitary lacquered			
cans, wt. 26 lbs	100	5.55	54.45
No. 2 Lacquered cans, wt. 6 lbs	24	1.45	58.25
No. 3 Lacquered cans, wt. 36 lbs	100	8.40	82.70
No. 3 Lacquered cans, wt. 9 lbs	24	2.20	87.75
			Per 100
Extra Covers for No. 2 Plain Sanitary	can	S	. \$1.35
Extra Covers for No. 3 Plain Sanitary			

FRICTION TOP CANS

f.o.b. Plant City

Friction top or syrup cans are used for packaging syrup, honey and for many other purposes.

		o. in ase	Case	1,000
11/2	Friction-top cans, wt. 8 lbs	36	\$2.25	\$ 61.85
No.	2½ Friction-top (approx. ¼		•	•
	gal.) cans, wt. 11 lbs	24	2.45	100.30
No.	5 Friction-top, holds 4½ lbs.			
	of Honey, wt. 24 lbs	50	7.25	143.40
No.	5 Friction-top cans, wt. 6 lbs.	12	1.85	151.40
No.	10 Friction-top (approx. 1			
	gal.) cans, wt. 39 lbs	50	10.90	216.45
5 ga	l. Square Honey Cans—each 9	0c; c	ase (16) \$12.40.
50-1	b. Lard Can—each \$1.00; case	of	12\$11	.20.
Cor	rugated Shipping Cartons for n 25; 10 for \$2.30; 25 for \$5.50	5-ga	al. Hone	ey Cans,

ATLAS FRUIT JARS

Atlas Square Mason Jars,

f.o.b. Plant City



Atlas Jars are made of clear crystal glass of highest quality giving a natural color to packed fruits and vegetables. Best preserving jar on the market, guaranteed to give entire satisfaction when properly used.

Per doz.

½-pint, wt. 7 lbs\$1	1.10
Atlas Square Mason Jars, pint, wt. 9 lbs.	1.20
Atlas Square Mason Jars, quart, wt. 12 lbs 1	
Atlas Square Mason, ½-gal., wt. 22 lbs	1.55
Atlas Wide Mouth, pint, wt. 10 lbs.	1.35
Atlas Wide Mouth, quart, wt. 16 lbs.	L.55
, -	
CAPS FOR ALL TYPES JARS	
f.o.b. Plant City Per	doz.
f.o.b. Plant City Per	
f.o.b. Plant City Per Atlas Seal All Standard Caps, 2-piece, wt. 9 oz\$	
f.o.b. Plant City Per	.37
f.o.b. Plant City Atlas Seal All Standard Caps, 2-piece, wt. 9 oz\$ Atlas Seal All Standard Lids only, wt. 4 oz 63 MM 2-piece Caps for Coffee and Relish Jars, wt. 8 oz.	.37 .17
Atlas Seal All Standard Caps, 2-piece, wt. 9 oz\$ Atlas Seal All Standard Lids only, wt. 4 oz 63 MM 2-piece Caps for Coffee and Relish Jars, wt. 8 oz	.37 .17 .37 .15
Atlas Seal All Standard Caps, 2-piece, wt. 9 oz\$ Atlas Seal All Standard Lids only, wt. 4 oz 63 MM 2-piece Caps for Coffee and Relish Jars, wt. 8 oz	.37 .17
Atlas Seal All Standard Caps, 2-piece, wt. 9 oz\$ Atlas Seal All Standard Lids only, wt. 4 oz 63 MM 2-piece Caps for Coffee and Relish Jars, wt. 8 oz	.37 .17 .37 .15

JELLY GLASS

### ### ##############################	
HONEY JARS f.o.b. Plant City wt. per	per
Utility Round case	doz.
½-lb., 2 doz. to case	\$.60
2-lb., 1 doz. to case	.90
No. 5 Glass Honey Pail	
No. 10 Glass Honey Pail14 lbs.	2.05
Queenline	
½-lb., 2 doz. in case 8 lbs.	\$.65
1-lb., 2 doz. in case	.85
2-lb., 1 doz. in case 9. lbs.	1.05
Comb Honey	
1 lb. Round Wide Mouth, 24 to case14 lbs	
2½ lbs. Square Wide Mouth, 12 to case. 12 lbs	s. 1.40

5 lbs. Square Wide Mouth, 6 to case10 lbs.

2.40

BEE KEEPERS' SUPPLIES

Prices subject to change without notice and are f.o.b. Plant City or any Kilgore store.

If you wish your shipment to be made by mail, please add postage when ordering. Most all Bee Supplies are exempt from State Tax except Hive Bodies and Covers. The following is only a partial list of our large and complete line of bee supplies carried in stock in all Kilgore's Florida stores and in Kilgore's mail order department at Plant City, Florida. We can always get whatever a customer wants from the factory if we happen to be out of stock, in which case the price will be f.o.b. factory, Medina, Ohio, and a 5% discount will be allowed on orders of \$100.00 or more. Allow two or three weeks for factory shipments.

We Buy Wax. Write for price
Write for complete catalog of Bee-keepers' supplies



Brood Foundation — (3-Ply) 8 1/16 x 163/4 in. — 7 sheets to pound Weight Price	Deep 10-Frame Extracting Supers C-1 1 KD Full depth super, empty wt. 12 lbs. \$ 2.98 C-2 5 KD Full depth supers, empty wt. 45 lbs. 12.95
H-1 10-sheet Carton	Shallow 10-Frame Extracting Supers C-12 1 KD Empty only, no fittings. Wt. 5 lbs. \$1.80
Medium Brood 8 1/16 x 16¾ in. — 8 sheets to pound	C-13 5 KD Empty only, no fittings. Wt. 28 lbs. 7.95
H-20 10-sheet Carton	Beeway Section Supers D-5 1 KD Without sections or foundation. Wt. 7 lbs
3 Ply Foundation	77 0 0 1881
for 2-pc. btm bar — 7 sheets to pound	List of Inside Fixtures for Section Supers
H-8 10-sheet Carton	E-1 7 Beeway Section Holders
H-10 50-sheet Carton	KD Wt. 3 lbs. \$ 1.10 E-2 50 Beeway Section Holders
Medium Brood	KD
4½ x 16% in. — 16 sheets to pound	KD Wt. 23 lbs. 11.95
Weight Price	E-15 10 Beeway Wood Separators, 1/10 in. thick Wt. 2 lbs75
H-26 10-sheet Carton 2 lbs. \$ 1.25	E-35 100 Super Springs (3 used to a
H-28 50-sheet Carton	super) Wt. 1½ lbs. 1.80
Medium Brood, Crimp-Wired	Comb Honey Sections — (Mill Run)
8½ x 16¾ in. — 7 sheets to pound	F-1 100 Wt. 7 lbs\$ 4.05
Weight Price	F-3 500 Wt. 32 lbs
D-350 10-sheet Carton	
D-000 50-Sifeet Carton	Metal Covers, 10-Frame
Medium Brood, Crimp-Wired	B-1 1 KD Metal Cover with inner
$434 \times 1634 - 12$ sheets to pound	cover Wt. 11 lbs. \$ 4.10 B-2 5 KD Metal Cover with inner
Weight Price	cover Wt. 51 lbs. 17.95
D-352 10-sheet Carton 2 lbs. \$ 1.65	
D-352 50-sheet Carton 6 lbs. 7.00	Inner Covers, 10-Frame
	B-8 1 KD Inner Cover, without bee
Thin Super	escape Wt. 3 lbs. 1.20 B-9 5 KD Inner Cover, without bee
J-1 1-lb. Carton, 3% x 15% in., 28 sheets to lb	escape Wt. 12 lbs. 5.10
J-3 5-lb. Carton, 37% x 1534 in.,	
28 sheets to 1b	Bottom Boards, 10-Frame
J-22 1-lb. Carton, $4\frac{1}{2} \times 16\frac{3}{4}$ in.,	B-12 1 KD Reversible Bottom
22 sheets to lb	Board Wt. 7 lbs. 2.40 B-13 5 KD Reversible Bottom
J-24 5-lb. Carton, 4½ x 16¾ in., 22 sheets to lb	Board Wt 37 lbs 940
22 sheets to lb	
22 sheets to lb	
J-29 5-lb. Carton, 5 x 16¾ in.,	
22 sheets to lb	
One-story standard hive with aluminum cover	
Hives in the flat are furnished with nails, reversible	
bottom boards, metal cover, inner cover, Hoffman	
frames, and complete directions for nailing. 161/4x20 in.	
A-1 Knocked down without foundation.	
Wt. 31 lbs\$10.30	Shallow Extracting Super


BEE-KEEPERS' SUPPLIES (Continued)

For information on Beekeeping in Florida, write the Florida Agricultural Experiment Station, Mailing Department, Gainesville, Florida, for free copy of Bulletin No. 151, entitled "Beginning Beekeeping." This publication is illustrated and contains much valuable information for beginners.

Frames

Furnished in the flat with nails. All frames are furnished with solid bottoms unless otherwise specified. Standard frames. Size 91/8x17 11/16. 10 50 1-piece bottom bar. Wt. 45 lbs. G-2 G-1 G-6 per 100\$2.20 \$8.75 \$16.50 Standard frames. Size 81/2x163/4. 2-piece bottom bar. Wt. 45 lbs. G-14 G-15 G-26 per 100\$2.20 \$8.75 \$16.50 Shallow frames. Size 4½x16¾. Thick top bar, 1-piece bottom bar. G-25 G-26 Wt. 38 lbs. per 100 \$7.95 \$14.95 Shallow frames. Size 4¾x16¾.

Thick top bar, 2-piece bottom bar. G-24 G-27 G-28 \$8.00 \$15.00 Wt. 37 lbs. per 100\$2.05




L-15

Honey Extractors

Bee Escape

L-15	2-frame Reversible			
	(9½ x 17 in. pocket)Wt.	100	lbs.	\$ 79.50
L-20	20-frame SimplicityWt.	300	lbs.	257.00
L-25	4-frame Automatic Reversible			
	(9½ x 17 in. pocket)Wt.	205	lbs.	172.50
L-35	3-frame, hand powerWT.	41	lbs.	41.50
L-45	45-frame Simplicity Wt.	470	lbs.	406.50



O-5 Double Bee Escape. Wt. 1 oz		\$.35
Queen Excluders		
Bees pass through rapidly, without hesitation.	Each	10
10-frame Standard ? wire	B-30	B-31

nesitation.	Lacii	10
10-frame Standard 3 wire.	B-30	B-31
Wt. 4 lbs. ea., 25 lbs. for 10	.\$1.85	\$17.95
10-frame Special 7-wire.		B-36
Wt. 4 lbs. ea., 32 lbs. for 10	\$1.80	\$17.50
Unbound Zinc Excluders.		B-41
Wt. 1 lb. ea., 8 lbs. for 10	\$1.55	\$14.85

No. 28 Tinned Wire

Standard frame takes 6½ feet of wire; Shallow frame, 3½ feet.

G-61	1/4-lb. Spool, 350 ft8 oz.	\$ 42
G-62	½-lb. Spool, 700 ft	.65
G-63	1-lb. Spool, 1400 ft	1.15
G-64	5-lb. Spool, 7000 ft7 lb.	4.60
N-65	Spur Wire Imbedder6 oz.	.45



Bristle Bee Brush N-85 Wt. 4 oz. .65
Steel Hive Tool N-81 10-in. Wt. 6 oz\$1.20
Queen and Drone Trap (10-frame) O-8 Wire Trap. Wt. 2 lbs\$1.50
Feeders

O-2 Boardman Feeder, with cap. Wt. 12 oz......\$.35

Smokers


N-3 Standard Tin Smoker, 3½x7 in. Wt. 2 lbs.\$2.55

N-7 Big Jumbo Tin Smoker, 4x10 in. with asbestos shield. Wt. 3 lbs.\$3.95



Leather and Drill Bee Gloves

Leather (N-21 medium or N-22 large). Wt. 2 lbs.....\$2.75 Drill (N-25 medium or N-26 large). Wt. 1 lb. 1.55



Veils	
N-10	Root Indestructible Bee Veil. Wt. 1 lb\$1.95
N-12	Dephos Folding Veil. Wt. 1 lb 2.45
N-15	Cotton Tulle Veil. Wt. 3 oz 1.50

INDEX

Page Page Page SUPPLIES GENERAL INFORMATION VEGETABLE AND FIELD SEED Directions to Customers Airlayering Kit _____87 Airwraps 87 Home Gardens Beans Insect and Disease 59 Beans, Velvet Bee Supplies _____90-91 Recommendations _____65-71 10 Beets _____ Budding Rubber Bands 87 Introduction of Pesticides Beets, Spinach _____ Budding Knives _____ 88 Locating the Garden Beets, Stock Making a Seedbed _____ 55 Budding Tape _____ 88 Bene or Sesame Miller Bill and Tolerances 10 Cans ____ 89 Ordering and Mailing1 Brussels Sprouts 10 Can Sealers 89 Cabbage 11-12
Cabbage, Chinese 12 Pesticide Dilution Table _____64 Pesticide Compatibility and Cane Stakes 88 Interval Chart _____62-63 Cyclone Seeders Cantaloupe 13 85 Plants per Acre _____ 59 Carrots Dusters ____ Preparing the Garden Cauliflower _____ Reference Table for Vegetable and Flower Gardens 4, 5, 6 Fertilizers 15 Celery ____ Fertilizer Spreaders Chufas _____ 55 Safe Combinations—Pesticides _____61 Clover, Alyce _____ 53 Flame Sprayers _____ 83 Sales Tax Clover, Hubam 54 Fly Baits ____ 78 60 Sanitation _____ Fly Traps 64 Seed Treatments Clover, Floranna Standard Household Measurements 64
What and When to Plant in Florida 6 Fruit Jars _____ 89 Clover, Kenland Red _____ Clover, Ladino 54 Fumigants _____ 76 Collard _____ Fungicides _____74-75 FLOWER SEED Corn, Field and Roasting Ear ____ Garden Tools _____85-86 Alyssum ____ Corn, Sweet 18 39 Garden Tractors Cowpeas, Field and Table ____ Aster Gloves _____ 55 87 Baby's Breath (Gypsophila) 39 Crotalaria _____ Cucumber _____19-20 39 Grafting Wax Bachelor Button (Centaurea) 88
 Dill
 22

 Eggplant
 21
 Balsam—Double Camelia flowered.... 39 Hand Tallys _____ 88 Bean-Scarlet Runner 39 Honey Jars _____ 89 Egyptian Wheat _____59 Bells of Ireland Endive ______ 22
Flower Seed ______ 39-50 Blanket Flower or Gaillardia Hormones _____ 80 40 Blue-Eyed African Daisy or Arctotis ... 40 Blue Lace Flower (Didiscus) Field Seed _____51-59 40 Household and Garden Pesticides Garlic ____ Calendula 40 Inoculants Grass, Lawn _____51-52 California Poppy (Eschscholtzia) Grass, Pasture _____ 53 Insco Applicators 40 Calliopsis or Coreopsis Gladiolus Blubs Insecticides ______72-73 Candytuft ____ 41 58 Carnation Hegari ____ Insecticide-Fungicide Combinations 75 Celosia (Cockscomb) Hairy Indigo 55 Jelly Glasses _____ Chinese Forget-Me-Not or Kale Kohl Rabi Lawn Mowers Cynoglossum _____ 41 Leek _____ Cleome (Spider Plant) 41 Lawn Sprinklers 84 Cosmos _____ 23 Lettuce _____ 41 Livestock Preparations 78 Lupine _____55-56 41 Nutrients _____ Delphinium (Perennial Larkspur) Millet _____ 56 42 Osmundia Fibre Mustard _____ 24 87 Floss Flower or Ageratum Paper Pots _____ 87 Globe Amaranth or Fla. Bachelor Oats ____ 56 Button (Gomphrena) 42 Okra _____ Paste ____ Gourd-Ornamental 42
Hollyhock 42 Peat Moss Onion Onion Sets _____ 26 Planet Jr. Tools Parsley 27 Larkspur 42 Plant Bands and Flats 87 Linaria (Baby Snapdragon) 43 Lupine 43 Parsnip _____ Plant Labels Peanuts56-57 Polyethylene Film _____ Marigold ______43-44 Peas, English or Garden _____ Poultry Preparations Pepper _____28-29 Morning Glory (Improved Ipomea) 44 Poultry Supplies 82 Potatoes Mourning Bride or Scabiosa 47 Nasturtium _____44-45 Pressure Cookers _____ Pumpkin ____ 30 Pansy 45
Petunia 45-46 Radish Raffia _____ 88 Rape, Dwarf Essex 57 Rodenticides _____ Rice _____Rutabaga 5747 Soil Test Kits Pinks—Chinese or Dianthus Soluble Plant Foods Portulaca or Sun Plant Rye ____ 48 Sphagnum Moss Sage _____ Queen Anne's Lace 47 Sprayers Sesbania Salvia ____ 47 Spreader-Stickers Shasta Daisy 47 Shallu _____ Snapdragon or Antirrhinum
Statice
Strawflower or Helichrysum Tarpaulins ____ Sorghum _____ 58 48 Spinach Terra-Lite Squash _______32-33 48 Thermometers _____ Strawberry Plants 50 Sunflower 59 Sunflower or Helianthus 48 Train-etts Sun Plant or Portulaca 48 Twine _____ Swiss Chard Sweet Peas Twist-Ems Tithonia or Florida Sunflower 49
Verbena 49 Tomato _____34-35 Vermicides Turnip _____ Weed Killers _____ 77 Vinca or Periwinkle Watermelon _____ 37-38 49 Wiss Shears Zinnia _____49-50 Wheat, Egyptian _____ 59



ORDER BLANK

THE KILGORE SEED COMPANY


PLANT CITY, FLORIDA
A DIVISION OF ASSOCIATED SEED GROWERS, INC.



AMOUNT ENCLOSED

Ship by Exp. or Frt. SPECIAL On item Sales Ta We war within r seeds, b	Parcel Post Express Truck Freight Total	the "State ne container rice of such COMPANY. sint against 10.00 filing	e Ce
SPECIAL On item Sales Ta We war within r seeds, b Florida Seller a fee, wit	Parcel Post Express Truck Freight Total	the "State ne container rice of such COMPANY. sint against 10.00 filing mail. Price	1
SPECIAL On item Sales Ta We war within r seeds, b Florida Seller a fee, with	IMPORTANT! Be sure address and shipping instructions are correct. Please write plainly. AL NOTICE: Remittances should be made by postal money orders, checks, or cash. DO NOT SEN ms not marked postpaid be sure your remittance includes the necessary amount for postage and also fax" on non-exempt merchandise. The arrant that the seeds, bulbs and plants we sell are, at the time of delivery, as described on the recognized tolerances; but we limit our liability on this warranty to the amount of the purchase probulbs and plants. We give no other or further warranty express or implied. THE KILGORE SEED statutes Section 578.26 requires as prerequisite to suit for defective seeds that sworn complaint alleging damages sustained be filed with Florida Commissioner of Agriculture, accompanied by \$\frac{5}{2} in the days after defect becomes apparent; and copy of complaint be sent to Seller by registered in the c	the "State ne container rice of such COMPANY. sint against 10.00 filing mail. Price	1
SPECIAL On item Sales Ta We war within r seeds, b Florida Seller a fee, wit	IMPORTANT! Be sure address and shipping instructions are correct. Please write plainly. AL NOTICE: Remittances should be made by postal money orders, checks, or cash. DO NOT SEN ms not marked postpaid be sure your remittance includes the necessary amount for postage and also fax" on non-exempt merchandise. The arrant that the seeds, bulbs and plants we sell are, at the time of delivery, as described on the recognized tolerances; but we limit our liability on this warranty to the amount of the purchase probulbs and plants. We give no other or further warranty express or implied. THE KILGORE SEED Statutes Section 578.26 requires as prerequisite to suit for defective seeds that sworn complained alleging damages sustained be filed with Florida Commissioner of Agriculture, accompanied by \$ in the days after defect becomes apparent; and copy of complaint be sent to Seller by registered in the complaint that the seeds are addressed in the complaint that the seeds are addressed in the complaint that the seeds are addressed in the complaint to the service of the complaint that the seeds are addressed in the complaint that the complaint	the "State ne container rice of such COMPANY. sint against 10.00 filing mail. Price	1
On item Sales Ta: We war within r seeds, b Florida Seller a fee, with	NOTICE: Remittances should be made by postal money orders, checks, or cash. DO NOT SEN ms not marked postpaid be sure your remittance includes the necessary amount for postage and also fax" on non-exempt merchandise. arrant that the seeds, bulbs and plants we sell are, at the time of delivery, as described on the recognized tolerances; but we limit our liability on this warranty to the amount of the purchase probulbs and plants. We give no other or further warranty express or implied. THE KILGORE SEED Statutes Section 578.26 requires as prerequisite to suit for defective seeds that sworn complaints alleging damages sustained be filed with Florida Commissioner of Agriculture, accompanied by \$100 per proposed in the days after defect becomes apparent; and copy of complaint be sent to Seller by registered in the days after defect becomes apparent; and copy of complaint be sent to Seller by registered in the	the "State ne container rice of such COMPANY. sint against 10.00 filing mail. Price	1
We war within r seeds, b Florida Seller a fee, wit	arrant that the seeds, bulbs and plants we sell are, at the time of delivery, as described on the recognized tolerances; but we limit our liability on this warranty to the amount of the purchase probulbs and plants. We give no other or further warranty express or implied. THE KILGORE SEED Statutes Section 578.26 requires as prerequisite to suit for defective seeds that sworn complaints alleging damages sustained be filed with Florida Commissioner of Agriculture, accompanied by \$ in the days after defect becomes apparent; and copy of complaint be sent to Seller by registered in the days after defect becomes apparent; and copy of complaint be sent to Seller by registered in the days after defect becomes apparent; and copy of complaint be sent to Seller by registered in the days after defect becomes apparent; and copy of complaint be sent to Seller by registered in the days after defect becomes apparent; and copy of complaint be sent to Seller by registered in the days after defect becomes apparent; and copy of complaint be sent to Seller by registered in the days after defect becomes apparent; and copy of complaint be sent to Seller by registered in the days after defect becomes apparent; and copy of complaint be sent to Seller by registered in the days after defect becomes apparent.	rice of such COMPANY. sint against 10.00 filing nail. Price	1
Quantity	ITEMS WANTED		1
Quantity	TIEMS WANTED	Dollars	Co
STITUTES:	—You may substitute the next best or most similar of any variety you cannot supply. —Please do not substitute without first notifying me.		

Name



THE KILGORE SEED COMPANY

General Offices and Mail Order Department, Plant City, Florida

A DIVISION OF ASSOCIATED SEED GROWERS, INC.